MAT250S25 Proof 1

Kishan S Patel

January 27, 2025

Let $\mathbb{R}^n$ be any Euclidian Space, and let $\vec{v}_1, \vec{v}_2, \vec{v}_3 \in \mathbb{R}^n$ be vectors such that the set of vectors $\{\,\vec{v}_1, \vec{v}_2, \vec{v}_3\,\}$ is linearly independent.

Lemma 1. Any ordered set B is Linearly Independent if it is a subset of a linearly independent set.

$$B \in L_I \iff (B \subset A) \wedge A \in L_I$$

Prove that the set of vectors $\{\,\vec{v}_1, \vec{v}_2, \vec{v}_1 + \vec{v}_3\,\}$ is linearly independent (Proposition P).

Proof. Let $\mathcal{F}$ be the set of all ordered sets that have elements $\in \mathbb{R}$, but not any ordered set with all 0 values.
Let Q be the proposition that $\{\,\vec{v}_1, \vec{v}_2, \vec{v}_3\,\}$ is linearly independent.
Assuming $Q \wedge \neg P \to \{\,\vec{v}_1, \vec{v}_2, \vec{v}_1 + \vec{v}_3\,\}$ is linearly dependent.

$$\neg P \to \exists\,\{\,x_1, x_2, x_3\,\} \in \mathcal{F} : x_1\vec{v}_1 + x_2\vec{v}_2 + x_3(\vec{v}_1 + \vec{v}_3) = \vec{0}$$

$$\sim (x_1 + x_3)\vec{v}_1 + x_2\vec{v}_2 + x_3\vec{v}_3 = \vec{0}$$
$$(x_1 + x_3)\vec{v}_1 + x_2\vec{v}_2 = -x_3\vec{v}_3$$
$$-\left(\frac{x_1}{x_3} + 1\right)\vec{v}_1 - \frac{x_2}{x_3}\vec{v}_2 = \vec{v}_3$$

$$\text{let } t = -\frac{x_1}{x_3},\, s = -\frac{x_2}{x_3}$$

$$\neg P \implies \exists t, v : t\vec{v}_1 + s\vec{v}_2 = \vec{v}_3$$

This brings a contradiction: assuming $\neg P$ implies that there exists two real numbers t, v (who are not both 0) that can be made into a linear combination with $\{\,\vec{v}_1, \vec{v}_2\,\}$ to equal $\vec{v}_3$, therefore $\neg P \implies \{\,\vec{v}_1, \vec{v}_2, \vec{v}_3\,\}$ is linearly dependent.

As shown by Lemma 1, $\{\,\vec{v}_1, \vec{v}_2\,\}$ is linearly independent as it is a subset of the linearly independent set $\{\,\vec{v}_1, \vec{v}_2, \vec{v}_3\,\}$.

This shows that $Q \wedge \neg P \implies \neg Q$, which is a contradiction which must mean that P must be true, and the set $\{\,\vec{v}_1, \vec{v}_2, \vec{v}_1 + \vec{v}_3\,\}$ is linearly independent.

QED