

MAT250S25 Proof 2

Kishan S Patel

February 14, 2025

Theorem 1 (\mathcal{L}_1 , from the **Big Theorem**TM). A set of vectors is linearly independent if and only if it is injective.

Theorem 2 (\mathcal{L}_2 , from the **Big Theorem**TM). A set of vectors $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ is linearly independent if and only if:

$$[\vec{x}_1 \ \vec{x}_2 \ \cdots \ \vec{x}_k] \vec{b} = \vec{0} \iff \vec{b} = \vec{0}$$

Given:

- let $\mathbb{R}^n, \mathbb{R}^m$ be Euclidean spaces.
- let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation with corresponding $n \times m$ matrix A .
- let $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in \mathbb{R}^m$ be vectors such that the set of vectors
- $\{T(\vec{x}_1), T(\vec{x}_2), \dots, T(\vec{x}_k)\}$ is linearly independent.

Prove that the set of vectors $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ is linearly independent.

Proof.

$$\forall X \in M_{n \times m} f_X : \vec{b} \mapsto X\vec{b}$$

$$\text{let } C = [T(\vec{x}_1) \ T(\vec{x}_2) \ \cdots \ T(\vec{x}_k)]$$

$$\text{let } B = [\vec{x}_1 \ \vec{x}_2 \ \cdots \ \vec{x}_k]$$

$$C = A \circ B$$

$$f_C = T \circ f_B$$

$$\mathcal{L}_1 \implies \forall \vec{a}, \vec{b} : f_C(\vec{a}) = f_C(\vec{b}) \iff \vec{a} = \vec{b}$$

$$\text{let } \vec{b}, \vec{c} \in \mathbb{R}^k$$

$$f_B(\vec{b}) = f_B(\vec{c})$$

$$(T \circ f_B)(\vec{b}) = (T \circ f_B)(\vec{c})$$

$$f_C(\vec{b}) = f_C(\vec{c})$$

$$\vec{b} = \vec{c}$$

$\therefore f_B$ is injective

$$\implies \forall \vec{b}, \vec{c} \in \mathbb{R}^k : B\vec{b} = B\vec{c} \iff \vec{b} = \vec{c}$$

$$\implies B\vec{b} = \vec{0} \iff \vec{b} = \vec{0}$$

$$\cdots \wedge \mathcal{L}_2 \implies \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \text{ is linearly independent.}$$

QED