

**SEARCH FOR LIGHT NEUTRAL BOSONS IN THE TREK/E36  
EXPERIMENT AT J-PARC**

A Dissertation

by

DONGWI HANDIIPONDOLA DONGWI

Submitted to the Graduate College of Hampton University in  
partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

August 2020

This dissertation submitted by Dongwi Handiipondola Dongwi in partial fulfillment of the requirements for the degree of Doctor of Philosophy at Hampton University, Hampton, Virginia is hereby approved by the committee under whom the work has been completed.

---

Michael Kohl, Ph.D.  
Committee Chair

---

José L. Goity, Ph.D.

---

Michael Hasinoff, Ph.D.

---

Alberto Accardi, Ph.D.

---

Narbe Kalantarians, Ph.D.

---

Michelle Penn-Marshall, Ph.D.  
Dean, The Graduate College

---

Date

Copyright by  
DONGWI HANDIIPONDOLA DONGWI  
2020

# ABSTRACT

Search for Light Neutral Bosons in the TREK/E36 Experiment at J-PARC

(August 2020)

Dongwi Handiipondola Dongwi, B.A., Westminster College;

Ph.D. Hampton University

Chair of Advisory Committee: Dr. Michael Kohl

The Standard Model (SM) represents our best description of the subatomic world and has been very successful in explaining how elementary particles interact under the influence of the fundamental forces. Despite its far reaching success in describing the building blocks of matter, the SM is still incomplete; falling short to explain dark matter, baryogenesis, neutrino masses and much more. The E36 experiment conducted at J-PARC in Japan, allows for sensitivity to search for light  $U(1)$  gauge bosons, in the muonic  $K^+$  decay channel. Such  $U(1)$  bosons could be associated with dark matter or explain established muon-related anomalies such as the muon  $g - 2$  value, and perhaps the proton radius puzzle. A realistic simulation study was employed for these rare searches in a mass range of 20 MeV to 100 MeV. Currently, about  $\sim 8\%$  of the data has been analyzed and upper limits for the  $A'$  branching ratio  $\mathcal{Br}(A')$  have been extracted at 95% CL.

Dedicated to my parents.

## ACKNOWLEDGEMENTS

I would like to acknowledge...

# TABLE OF CONTENTS

Chapter	Page
<b>ABSTRACT</b> . . . . .	<b>iv</b>
<b>DEDICATION</b> . . . . .	<b>v</b>
<b>ACKNOWLEDGEMENTS</b> . . . . .	<b>vi</b>
<b>TABLE OF CONTENTS</b> . . . . .	<b>vii</b>
<b>LIST OF TABLES</b> . . . . .	<b>viii</b>
<b>LIST OF FIGURES</b> . . . . .	<b>ix</b>
<b>1 Introduction</b> . . . . .	<b>1</b>
<b>REFERENCES</b> . . . . .	<b>3</b>

# LIST OF TABLES

Table

Page



## LIST OF FIGURES

Figure

Page

# CHAPTER 1

## INTRODUCTION

The TREK/E36 experiment was conducted at the Japan Proton Accelerator Research Complex (J-PARC), with physics data-taking occurring from September to December 2015, and has been decommissioned as of March 2016. TREK/E36 was part of the TREK program at J-PARC, where TREK stands for Time Reversal Experiment with Stopped Kaons. The primary goal of the TREK/E36 experiment was to provide a high precision electroweak measurement in order to test lepton universality, which is expressed as an identical coupling constant of the charged lepton family ( $e$ ,  $\mu$ , and  $\tau$ ). Lepton universality is a staple of the Standard Model (SM) and any violation of this would be clear evidence of New Physics (NP) beyond the SM [1].

Data analysis of two-body leptonic decays of stopped kaons  $K_{l2}$  is currently being performed in order to provide a precise measurement of the decay width ratio  $R_K = \Gamma(K^+ \rightarrow e^+\nu)/\Gamma(K^+ \rightarrow \mu^+\nu)$  to test lepton universality. Due to the  $V - A$  structure of charged current couplings, the ratio  $R_K$  is helicity-suppressed in the SM and is therefore sensitive to beyond SM physics. The  $K_{l2}$  decay width is calculated as

$$\Gamma(K_{l2}) = g_l^2 \frac{G^2}{8\pi} f_K^2 m_K m_l^2 \left(1 - \frac{m_l^2}{m_K^2}\right)^2 \quad (1.0.1)$$

where  $g_l$  is the coupling constant of the lepton current,  $G$  is the Fermi constant,  $f_K$  is the kaon form factor and  $m_K$  and  $m_l$  are the kaon and lepton masses respectively. The SM value for  $R_K$  is very precise because to a first approximation the strong interaction dynamics from equation 1.0.1 cancel

$$\begin{aligned}
R_K^{SM} &= \frac{\Gamma(K^+ \rightarrow e^+ \nu)}{\Gamma(K^+ \rightarrow \mu^+ \nu)} \\
&= \frac{m_e^2}{m_\mu^2} \left( \frac{m_K^2 - m_e^2}{m_K^2 - m_\mu^2} \right)^2 (1 + \delta_r) \\
&= (2.477 \pm 0.001) \times 10^{-5}
\end{aligned} \tag{1.0.2}$$

where  $\delta_r$  represents radiative corrections, detailed calculations of which were carried out in. Thus the SM value for  $R_K^{SM}$  has been calculated to high accuracy ( $\Delta R_K/R_K \sim 0.4 \times 10^{-4}$ ) thereby making it possible to search for NP effects by conducting a precise measurement of  $R_K$  [2].

## REFERENCES

- [1] K. Paton, et al. Measurement of T-violation Transverse Muon Polarization in  $K^+ \rightarrow \pi^0 \mu^+ \nu$  Decays.
- [2] M. Kohl. The TREK/E36 experiment at J-PARC. *AIP Conf. Proc.*, 1563:147–150, 2013.