



## COVID-19 type III hypersensitivity reaction



To the editor,

The outbreak of the novel COVID-19 (coronavirus disease 2019) centered in Hubei Province of the People's Republic of China and spread to many other countries in the world and on thirty of January 2020, the WHO declared a global health emergency in the world [1]. Tyrell and Bynoe in 1966 were first described and cultivated coronaviruses as enveloped, positive single-stranded large RNA viruses that infect humans and animals [2]. The median incubation period after infection is three days ranged from 0 to 24 days [3]. The clinical manifestation of the disease consisting of fever, dry cough, nasal congestion, fatigue, headache and the disease progress to dyspnoea and pneumonia [4]. Some reports describe gastrointestinal, fulminant myocarditis [5]. COVID-19 infects lung alveolar epithelial cells using receptor-mediated endocytosis through coronavirus S (spike) protein contains a ligand binding domain with the angiotensin-converting enzyme II (ACE2) [6]. After virus entry; the first line of defence mechanism (Innate) is recognition of invaded virus by pathogen associated molecular patterns (PAMPs) via endosomal RNA receptors, TLR3, TLR7 and the cytosolic RNA sensor, RIG-I/MDA5. This recognition leads to activation of the downstream signaling cascade NF- $\kappa$ B and IRF3 that induce expression of type I IFN and other pro-inflammatory cytokines that activates the JAK-STAT pathway that initiate the transcription of IFN type 1 [7]. There is also increased in number of neutrophils and decrease in number of lymphocytes (lymphopenia) which is the second line of defense mechanism (Adaptive) and elevated of inflammatory markers like C-reactive protein and proinflammatory cytokines like IL-2, IL-7, IL-10, G-CSF, IP-10, MCP-1, MIP-1A, and TNF $\alpha$  which is correlate with disease severity and death [8]. The virus then directly infects macrophages and T cells [9]. The macrophages will present viral antigenic peptide to T cells that differentiate into follicular helper cells (T<sub>FH</sub> cells), activated T cells CD4, CD8 + T cells and antibody secreting cells (ASC) (CD3-CD19 + CD27hiCD38hi ASC and CD4 + CXCR5 + ICOS + PD-1 + cTFH cell). This will recruitment of different immune cell populations ASCs, TFH cells and activated CD4 + and CD8 + T cells with IgM and IgG SARS-CoV-2-binding antibodies in the patient's blood [10]. The COVID-19 viral antigens lead to stimulate antibodies formation of IgM in acute phase and IgG type in chronic phase which is facilitate viral entry and fusion with infected cell through uptake of the virus-IgG complex via the Fc family of receptors and later viral fusion with antigen presenting cells like macrophages, B cells, monocytes via FcR family, and vascular endothelium through the neonatal Fc receptor (nFcR) instead of antibodies induced viral agglutination and this is known as antibody dependent enhancement (ADE) [11–13]. This Ag-Ab (IgG1 and IgG3) immune complex that deposit in the lung will also end in complement activation by classical pathway of the complement (membrane attack complex MAC) end in formation

C3a, C5a and C5b67 that leads to vasoactive amines release from mast cell and basophile and platelets aggregation leads to microthrombus formation which block coronary blood vessels of the heart and also leads to neutrophils infiltration and aggregation at site of infection in alveoli leading to sever tissue damage which is the main cell in type III hypersensitivity reaction and the patient showed increased in neutrophils number which is the same immune mechanism in rheumatoid arthritis [14]. Thus function of complement activation instead of viral clearance and antigen presentation, phagocytosis now act as other route for viral infection [15]. In addition to that; the immune complex Anti-S protein:CoV immune complexes leads to increased cytokine release like MCP-1 and IL-8 in lung macrophages which is known as cytokine storm syndrome while in the rheumatoid arthritis known as cytokine soup [16]. Some trails used the same monoclonal antibodies that used in treatment of rheumatoid arthritis like Tocilizumab which is monoclonal antibodies against IL-6R [17].

### Funding source

None.

### Prior presentation

None.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.mehy.2020.109763>.

### References

- [1] Velavan TP, Meyer CG. The COVID-19 epidemic. *Trop Med Int Health* 2020;25:278–80.
- [2] Tyrrell DA, Bynoe ML. Cultivation of viruses from a high proportion of patients with colds. *Lancet* 1966;1:76–7.
- [3] Guan W, Ni Z, Yu H, et al. Clinical characteristics of 2019 novel coronavirus infection in China. medRxiv preprint posted online on Feb. 9, 2020; <https://doi.org/10.1101/2020.02.06.20020974>.
- [4] Chan JF, Yuan S, Kok KH, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;S0140–6736(20):30154–9. [https://doi.org/10.1016/S0140-6736\(20\)30154-9](https://doi.org/10.1016/S0140-6736(20)30154-9).

- [5] SARS-CoV-2: a potential novel etiology of fulminant myocarditis. Chen C, Zhou Y, Wang DW. *Herz*. 2020 Mar 5. doi: 10.1007/s00059-020-04909-z. Online ahead of print.
- [6] Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of the SARS-CoV-2 by full-length human ACE2 [Journal Article]. *Science*. 2020; Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32132184>.
- [7] de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into emerging coronaviruses. *Nat Rev Microbiol* 2016;14:523–34.
- [8] COVID-19, ECMO, and lymphopenia: a word of caution. Henry BM. *Lancet Respir Med*. 2020 Mar 13:S2213-2600(20)30119-3. doi: 10.1016/S2213-2600(20)30119-3. Online ahead of print.
- [9] Perlman S, Dandekar AA. Immunopathogenesis of coronavirus infections: implications for SARS. *Nat Rev Immunol* 2005;5(12):917–27.
- [10] Thevarajan I, Nguyen THO, Koutsakos M, et al. Breadth of concomitant immune responses prior to patient recovery: a case report of non-severe COVID-19. *Nat Med* 2020. <https://doi.org/10.1038/s41591-020-0819-2>.
- [11] Wang J, Zandy MS. Potential mechanisms of age related severity of covid-19 infection: implications for development of vaccines, convalescent serum, and antibody therapies. *OSF Preprints*. 2020; March 21- DOI: 10.31219/osf.io/f3pze, rev. 3.
- [12] Wan Y, Shang J, Sun S, Tai W, Chen J, Geng Q, et al. Molecular mechanism for antibody-dependent enhancement of coronavirus entry. *J Virol*. 2020;94(5). Available from: <https://jvi.asm.org/content/94/5/e02015-19>.
- [13] *Immunol Rev* 2015;268(1):340–64. <https://doi.org/10.1111/imr.12367>.
- [14] Kumar Vinay. “6”. Robbins and Cotran Pathologic Mechanisms of Disease. (8th ed.). Philadelphia: Elsevier; 2010. p. 204–5.
- [15] Tirado SMC, Yoon KJ. Antibody-dependent enhancement of virus infection and disease. *Viral Immunol* 2003;16:69.
- [16] *J Pharm Anal* 2020. <https://doi.org/10.1016/j.jpha.2020.03.001>.
- [17] New therapeutic opportunities for COVID-19 patients with Tocilizumab: Possible correlation of interleukin-6 receptor inhibitors with osteonecrosis of the jaws. Bennardo F, Buffone C, Giudice A. *Oral Oncol*. 2020 Mar 21:104659. doi: 10.1016/j.oraloncology.2020.104659. Online ahead of print.

Batool Mutar Mahdi

*Consultant Clinical Immunology, Head of HLA Research Unit, Department of Microbiology, Al-Kindy Medical College, University of Baghdad, Baghdad, Iraq*