Ramakrishna Mission Vivekananda Educational and Research Institute



PO Belur Math, Howrah, West Bengal 711 202 School of Mathematical Sciences Department of Computer Science

MSc Big Data Analytics and MSc Computer Science : Batch 2023-25 DA109: Linear Algebra and Matrix Computation
Instructor: Dr. Soumitra Samanta
TA: Rajdeep Mondal

Problem set: 1

Please try to solve all the problems ¹ alone. If you are stuck, I encourage you first consult with your batch mates and then TA. If you still need help then come to Instructor.

- 1. Answer the followings:
 - i) By which property of a vector space V you will explain that 0 + 0 = 0 where 0 is the zero vector of V.
 - ii) Prove that:
 - a) 0.x = 0, where $x, 0 \in V$ and 0 is the zero vector.
 - b) (-a)x = -(ax) = a(-x) for each $a \in F$ and each $x \in V$.
- 2. What is the Parallelogram Law of Vector Addition? Consider the vector space \mathbb{R}^2 , add the vectors (3,1) and (2,1) in \mathbb{R}^2 by this law and explain it by a picture.
- 3. Prove that the set of all $m \times n$ matrices over a field \mathbb{F} denoted by $M_{m \times n}(\mathbb{F})$ is a vector space under usual matrix addition and matrix multiplication.
- 4. Let V denote the set of ordered pairs of real numbers. If (a_1, a_2) and (b_1, b_2) are elements of V and c is an element of F. Define, $(a_1, a_2) + (b_1, b_2) = (a_1 + b_1, a_2b_2)$ and $c(a_1, a_2) = (ca_1, ca_2)$. Is V is a vector space under this operations? Justify your answer.
- 5. Consider the set of all polynomials with coefficients from a field \mathbb{F} with usual polynomial addition and multiplication denoted by $P(\mathbb{F})$. Is $P(\mathbb{F})$ a vector space over \mathbb{F} ? Justify your answer.
- 6. Define the Subspace of a Vector Space. Consider \mathbb{R}^2 , prove that $W = \{(a, a) : a \in \mathbb{R}^2\}$ is a subspace of \mathbb{R}^2 . Draw the subspace in \mathbb{R}^2 . What are the basis and dimension of W?
- 7. Which of the following sets are the subspaces of \mathbb{R}^3 ?
 - (a) $A = \{(\lambda, \lambda + \mu^3, \lambda \mu^3) : \lambda, \mu \in \mathbb{R}^2\}$
 - (b) $B = \{(\lambda^2, -\lambda^3, 0) : \lambda \in \mathbb{R}\}$
 - (c) Let $\gamma \neq 0$ be in \mathbb{R} . $C = \{(\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3 : \xi_1 2\xi_2 + 3\xi_3 = \gamma\}$
 - (d) $D = \{(\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3 : \xi_2 \in \mathbb{Z}\}$
- 8. Let W_1 and W_2 be subspaces of a vector space V. Prove that $W_1 \cup W_2$ is a subspace of V if and only if $W_1 \subseteq W_2$ or $W_2 \subseteq W_1$.
- 9. The set of all differentiable real-valued functions f on the interval (0,3) such that f'(2) = b is a subspace of $\mathbb{R}^{(0,3)}$ where $\mathbb{R}^{(0,3)}$ is the set of real-valued functions on the interval (0,3) if and only if b=0.
- 10. Let V be a vector space and $v, w \in V$. Explain why there exists a unique $x \in V$ such that v + 3x = w.
- 11. Answer the following:

¹All the problems have been selected by the TA.

- (a) What are the standard bases and dimensions for the following vector spaces: $\{0\}$, \mathbb{R}^n , $P_n(\mathbb{R})$ (set of polynomials of degree $\leq n$), $M_{m \times n}(\mathbb{R})$ (set of matrices of order $m \times n$).
- (b) How do you relate the bases of $M_{m\times n}(\mathbb{R})$ and \mathbb{R}^k for any k?
- 12. Find the bases and dimensions of the following subspaces of \mathbb{R}^n :
 - (a) $W_1 = \{(a_1, a_2, \dots, a_n) \in \mathbb{R}^n : a_1 + a_2 a_3 = 0\}$
 - (b) $W_2 = \{(a_1, a_2, \dots, a_n) \in \mathbb{R}^n : a_{11} + 2a_{21} = 0, a_{27} = a_4 = a_9\}$
 - (c) $W_3 = \{(a_1, a_2, ..., a_n) \in \mathbb{R}^n : m \text{ constraints are given, } m \leq n\}$. What happens if we assume $m \geq n$?
- 13. Let $S = \{x_1, x_2, \dots, x_n\}$ be a linearly independent subset of a vector space V over the field \mathbb{Z}_2 . How many elements are there in span(S)? Justify your answer.
- 14. If V and W are vector spaces over \mathbb{F} for which |V| = |W| then does it mean that dim(V) = dim(W)?
- 15. Choose $x = (x_1, x_2, x_3, x_4)$ in \mathbb{R}^4 . It has 24 rearrangements like (x_2, x_1, x_3, x_4) and (x_4, x_3, x_1, x_2) . Those 24 vectors, including x itself, span a subspace S. Find specific vectors x so that the dimension of S is: (a) zero, (b) one, (c) three, (d) four.
- 16. Let V be a real vector space of all polynomial functions from \mathbb{R} into \mathbb{R} of degree 2 or less. Let t be a fixed real number and define $g_1(x) = 1$, $g_2(x) = x + t$, $g_3(x) = (x + t)^2$. Prove that $B = \{g_1, g_2, g_3\}$ is a basis for V. If $f(x) = c_0 + c_1 x + c_2 x^2$ then what are the coordinates of f in this ordered basis B?
- 17. Let $v = (x_1, x_2)$ and $w = (y_1, y_2)$ be vectors in \mathbb{R}^2 such that $x_1^2 + x_2^2 = y_1^2 + y_2^2 = 1$ and $x_1y_1 + x_2y_2 = 0$. Prove that $B = \{v, w\}$ is a basis for \mathbb{R}^2 . Find the coordinates of the vector (a, b) in the basis B. Can you interpret the conditions geometrically?
- 18. Let W_1 and W_2 be two finite-dimensional subspaces of a vector space V. Then prove the following:
 - (a) $W_1 + W_2$ is the smallest subspace of V that contains both W_1 and W_2 .
 - (b) $dim(W_1 + W_2) = dim(W_1) + dim(W_2) dim(W_1 \cap W_2).$
 - (c) If $V = W_1 + W_2$ then $V = W_1 \oplus W_2$ if and only if $dim(V) = dim(W_1) + dim(W_2)$, where \oplus denote the direct sum.
- 19. Suppose V is a vector space over \mathbb{R} . Prove that V cannot be written as the set-theoretic union of a finite number of proper subspaces.
- 20. Give an example of distinct linear transformations T and U such that N(T) = N(U) and R(T) = R(U). Where R(T) and N(T) are the Range of T and Null space of T respectively.
- 21. Suppose $U = \{(x, x, y, y) \in \mathbb{F}^4 : x, y \in \mathbb{F}\}$. Find a subspace W of \mathbb{F}^4 such that $U \bigoplus V = \mathbb{F}^4$, where \oplus denote the direct sum.
- 22. Let W_1, W_2, W_3 be three distinct subspaces of \mathbb{R}^{10} with each W_i has dimension 9. If $W = W_1 \cap W_2 \cap W_3$ then prove that $7 \leq dim(W) \leq 8$.
- 23. Let V and W be finite-dimensional vector spaces and $T: V \to W$ be linear.
 - (a) Prove that if dim(V) < dim(W), then T cannot be onto.
 - (b) Prove that if dim(V) > dim(W), then T cannot be one-to-one.
- 24. Suppose $T: \mathbb{R}^3 \to \mathbb{R}^4$ and $S: \mathbb{R}^4 \to \mathbb{R}^3$ be two linear transformations such that $S \circ T = I$ on \mathbb{R}^3 . Then show that $T \circ S$ is neither one-one nor onto.
- 25. T and S be two linear operators on \mathbb{R}^n such that ST = TS = 0 and T + S is invertible. Then show that:
 - (a) Rank(T) + Rank(S) = n
 - (b) Nullity(T) + Nullity(S) = n

[Hint: Use $Rank(T+S) \leq Rank(T) + Rank(S) \leq Rank(TS) + n$]

- 26. A linear transformation T rotates each vector in \mathbb{R}^2 clockwise through an angle θ . Find the matrix of T w.r.t the standard ordered basis.
- 27. Let $T: \mathbb{R}^4 \to \mathbb{R}^5$ such that Tx = 0 iff x = 0. Find Rank(T).
- 28. Let V be a vector space and $T: V \to V$ be a linear transformation. Prove that the following two statements about T are equivalent:
 - (a) The intersection of the range of T and the null space of T is the zero subspace of V.
 - (b) If T(Ta) = 0, then Ta = 0.
- 29. Find two linear operators T and U on \mathbb{R}^2 such that TU = O but $UT \neq O$.
- 30. If W is a k-dimensional subspace of an n-dimensional vector space V, then prove that W is the intersection of n k hyperspaces in V.
- 31. Let V be a finite-dimensional vector space and $T: V \to V$ be linear. Then if V = R(T) + N(T) prove that $V = R(T) \bigoplus N(T)$, where \bigoplus denotes the direct sum.
- 32. A function $T: V \to W$ between vector spaces V and W is called additive if T(x+y) = T(x) + T(y) for all $x, y \in V$. Prove that if V and W are vector spaces over the field of rational numbers, then any additive function from V into W is a linear transformation.
- 33. Let V and W be vector spaces, and let T and U be nonzero linear transformations from V into W. If $R(T) \cap R(U) = \{0\}$, prove that $\{T, U\}$ is a linearly independent subset of $\mathcal{L}(V, W)$.
- 34. Show that $W = \{T \in \mathcal{L}(\mathbb{R}^5, \mathbb{R}^4) : Nullity(T) > 2\}$ is not a subspace of $\mathcal{L}(\mathbb{R}^5, \mathbb{R}^4)$.
- 35. Suppose that V is finite-dimensional and that $T \in \mathcal{L}(V, W)$. Prove that there exists a subspace U of V such that $U \cap N(T) = \{0\}$ and $R(T) = \{Tu : u \in U\}$.
- 36. Suppose U, V, and W are arbitrary subspaces of a finite-dimensional vector space. Then prove that:
 - (a) $U \cap (V + W) \supset (U \cap V) + (U \cap W)$.
 - (b) $(U \cap V) + W \subset (U + W) \cap (V + W)$.