

Computational Methods in Relativistic Hydrodynamics

Applications to Astrophysical Mergers and Explosions

Biswarup Biswas

Jointly with the MU Computational Maths Group

Department of Mathematics, Mahindra University



National Conference on Computational Mathematics

Special Session on Numerical Relativity

July 3, 2026

- Relativistic hydrodynamics (RHD) plays a crucial role in modeling high-energy astrophysical phenomena.
- Examples include binary neutron star mergers¹.
- The thermodynamic properties of a relativistic gas are often modeled using the Synge equation of state².

¹ L. Baiotti and L. Rezzolla. "Binary neutron star mergers: a review of Einstein's richest laboratory". In: *Reports on Progress in Physics* 80.9 (2017), p. 096901

² J. L. Synge and P. M. Morse. *The relativistic gas*. 1958

Numerical Methods

Theorem Conservation Laws

The governing equations of relativistic hydrodynamics can be written as a system of hyperbolic conservation laws:

$$\frac{\partial U}{\partial t} + \sum_{i=1}^3 \frac{\partial F^i(U)}{\partial x^i} = S(U)$$

where U is the vector of conserved variables.

Definition Primitive Variables

The primitive variables (e.g., density ρ , velocity v , and pressure p) must be recovered from the conserved variables U at each time step, typically requiring numerical root-finding methods.

Simulation & Visualization

Dynamics of RHD Shockwaves

- Relativistic shockwaves exhibit complex interface instability.
- High-order WENO schemes are employed to resolve the shocks without spurious oscillations.
- The animation on the right shows the evolution of the density field over 21 frames.




Figure: 2D Relativistic flow evolution.  **Mahindra**
University
Global Thinkers. Engaged Leaders.

- Robust numerical schemes are essential for simulating relativistic flows.
- The `mucomp` package provides support for standard theme styles, math blocks, footnoting/citations, and animations.