

THE ORGANIC CHEMISTRY OF DRUG SYNTHESIS

VOLUME 3

DANIEL LEDNICER

Analytical Bio-Chemistry Laboratories, Inc.
Columbia, Missouri

LESTER A. MITSCHER

The University of Kansas School of Pharmacy
Department of Medicinal Chemistry
Lawrence, Kansas

A WILEY-INTERSCIENCE PUBLICATION

JOHN WILEY AND SONS

New York · Chichester · Brisbane · Toronto · Singapore

Copyright © 1984 by John Wiley & Sons, Inc.

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Section 107 or 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

Library of Congress Cataloging in Publication Data:

(Revised for volume 3)

Lednicer, Daniel, 1929—

The organic chemistry of drug synthesis.

"A Wiley-Interscience publication."

Includes bibliographical references and index.

1. Chemistry, Pharmaceutical. 2. Drugs. 3. Chemistry, Organic—Synthesis. I. Mitscher, Lester A., joint author. II. Title. [DNLN 1. Chemistry, Organic. 2. Chemistry, Pharmaceutical. 3. Drugs—Chemical synthesis. QV 744 L473o 1977]

RS403.L38 615'.19 76-28387

ISBN 0-471-09250-9 (v. 3)

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

With great pleasure we dedicate this book, too, to our wives,
Beryle and Betty.

The great tragedy of Science is the slaying of a
beautiful hypothesis by an ugly fact.

Thomas H. Huxley, "Biogenesis and Abiogenesis"

Preface

The first volume in this series represented the launching of a trial balloon on the part of the authors. In the first place, we were not entirely convinced that contemporary medicinal chemistry could in fact be organized coherently on the basis of organic chemistry. If, however, one granted that this might be done, we were not at all certain that the exercise would engage the interest of others. That book's reception seemed to give an affirmative answer to each of these questions. The second volume was prepared largely to fill gaps in the coverage and to bring developments in all fields up to a common date - 1976. In the process of preparing those volumes, we formed the habit of scrutinizing the literature for new nonproprietary names as an indication of new chemical entities in or about to be in the clinic. It soon became apparent that the decreased number of drugs being granted regulatory approval was not matched by a decrease in the number of agents being given new generic names. The flow of potential new drugs seemed fairly constant over the years. (For the benefit of the statistician, assignment of new USAN names is about 60 per year.) It was thus

obvious that the subject matter first addressed in Volume 1 was increasing at a fairly constant and impressive rate.

Once we had provided the background data up to 1976, it seemed logical to keep the series current by adding discussion of newer agents. Reports of drugs for new indications as well as the occurrence of brand-new structural types as drugs made it particularly important to update the existing volumes. The five-year cycle for preparation of new volumes represents a compromise between timeliness and comprehensiveness. A shorter period would date earlier entries. This volume thus covers compounds reported up to 1982.

As has been the practice in the earlier volumes, the only criterion for including a new therapeutic agent is its having been assigned a United States nonproprietary name (USAN), a so-called generic name. Since the focus of this text is chemistry, we have avoided in the main critical comments on pharmacology. The pharmacological activity or therapeutic utility described for the agents covered is that which was claimed when the USAN name was assigned.

The changes in chapter titles as well as changes in their relative sizes in going from volume to volume constitute an interesting guide to directions of research in medicinal chemistry. The first two volumes, for example, contained extensive details on steroid drugs. This section has shrunk to about a third of its former size in this book. The section on β -lactam antibiotics, on the other hand, has undergone steady growth from volume to volume: not only have the number of entries multiplied but the syntheses have become more complex.

This book, like its predecessors, is addressed to students at the graduate level in organic and medicinal chemistry as well as to practitioners in the field. It is again assumed that the reader has a comfortable grasp of organic synthesis as well as a basic grounding in biology.

We are pleased to acknowledge the helpful assistance of several individuals in preparing this volume. Particularly, we are grateful to Mrs. Janet Gill for preparing all of the illustrations and to Mrs. Violet Huseby for long hours and careful attention to detail in preparing the final copy and several drafts.

Daniel Lednicer
Dexter A. Mitscher

Dublin, Ohio
Lawrence, Kansas
January, 1984

Contents

Chapter 1.	Alicyclic and Cyclic Compounds	1
	1. Cyclopentanes	1
	a. Prostaglandins	1
	b. Retenoids	11
	c. Miscellaneous	13
	References	16
Chapter 2.	Phenethyl and Phenoxypropanolamines	19
	1. Phenylethanolamines	20
	References	34
Chapter 3.	Arylaliphatic Compounds	37
	1. Arylacetic Acid Derivatives	37
	2. Anilines, Benzyl Amines, and Analogues	45
	3. Diarylmethane Analogues	47
	4. Stilbene Analogues	50
	References	52
Chapter 4.	Monocyclic Aromatic Agents	55
	1. Aniline Derivatives	55
	2. Benzoic Acid Derivatives	58
	3. Benzenesulfonic Acid Derivatives	61
	References	63
Chapter 5.	Polycyclic Aromatic Compounds	65
	1. Indanones	65

	2. Naphthalenes	68
	3. Tricyclic Compounds: Anthracene, Phenanthrene, and Dibenzocycloheptene	72
	References	78
Chapter 6.	Steroids	81
	1. Estranes	82
	2. Androstanes	87
	3. Pregnanes	90
	4. Miscellaneous Steroids	99
	References	107
Chapter 7.	Compounds Related to Morphine	109
	1. Bridged Polycyclic Compounds	111
	2. Piperidines	116
	3. Miscellaneous Compounds	121
	References	124
Chapter 8.	Five-Membered Heterocycles	127
	1. Pyrroles and Pyrrolidines	127
	2. Furans	129
	3. Imidazoles	131
	4. Triazoles	137
	5. Pyrazolines	137
	6. Isoxazoles	138
	7. Tetrazoles	139
	8. Miscellaneous	139
	References	141
Chapter 9.	Six-Membered Heterocycles	145
	1. Pyridines	145
	2. Pyridazines	151
	3. Pyrimidines	152
	4. Miscellaneous Heterocycles	157
	References	162
Chapter 10.	Five-Membered Heterocycles Fused to Benzene	165
	1. Indoles	165
	2. Benzimidazoles	172
	3. Benzothiazoles	178
	References	179
Chapter 11.	Benzofused Six-Membered Heterocycles	183
	1. Quinoline Derivatives	183
	2. Isoquinoline Derivatives	186

3. Benzopyran Derivatives	188
4. Benzodioxane Derivatives	191
5. Benzoxazolinone Derivatives	191
6. Quinazolinone Derivatives	192
7. Phthalazines	195
8. Benzodiazapines and Related Substances	195
9. Miscellaneous	198
References	199
Chapter 12. Beta Lactams	203
1. Penicillins	203
2. Cephalosporins	209
References	221
Chapter 13. Miscellaneous Fused Heterocycles	225
References	250
Cross Index of Drugs	253
Cumulative Index, Vols. 1-3	261
Index	279

THE ORGANIC CHEMISTRY OF DRUG SYNTHESIS

VOLUME 3

1 Alicyclic and Cyclic Compounds


1. CYCLOPENTANES

a. Prostaglandins.

Few areas of organic medicinal chemistry in recent memory have had so many closely spaced pulses of intense research activity as the prostaglandins. Following closely on the heels of the discovery of the classical monocyclic prostaglandins (prostaglandin E_1 , F_2 , A_2 , etc.), with their powerful associated activities, for example, oxytocic, blood pressure regulating, and inflammatory, was the discovery of the bicyclic analogues (the thromboxanes, prostacyclin) with their profound effects on hemodynamics and platelet function. More recently, the non-cyclic leucotrienes, including the slow releasing substance of anaphylaxis, have been discovered. The activity these substances show in shock and asthma, for example, has excited considerable additional interest. Each of these discoveries has opened new physiological and therapeutic possibilities for exploitation. The newer compounds in particular are chemically and biologically short lived and are present in vanishingly small quantities so that much chemical effort has been expended

on finding more efficient means of preparing them, on enhancing their stability, and on finding means of achieving greater tissue specificity.


In addition to its other properties, interest in the potential use of the vasodilative properties of prostaglandin E_1 , alprostadil (4), has led to several conceptually different syntheses.¹⁻⁵ For this purpose, the classic Corey process¹ has to be modified by reversing the order of addition of the side chains to allow for convenient removal of the unwanted double bond in the upper side chain. For example, Corey lactone 1 is protected with dihydropyran (acid catalysis), reduced to the lactol with diisobutylaluminum hydride, and then subjected to the usual Wittig reaction to give intermediate 2. This is esterified with diazomethane, acetylated, and then catalytically hydrogenated to give intermediate 3 in which all of the oxygen atoms are differentiated. Further transformation to alprostadil (4) follows the well-trodden path of sequential Collins oxidation, Horner-Emmons olefination, zinc borohydride reduction, deetherification with aqueous acetic acid, separ-




ation of the resulting C-15 epimers, dihydropyranylation, saponification of the ester groups, Jones oxidation (to introduce the C-9 keto group), and finally, deetherification.


The classic method for controlling stereochemistry is to perform reactions on cyclic substrates. A rather lengthy but nonetheless efficient example in the prostaglandin field uses bicyclic structures for this purpose.² Bisacetic acid derivative 5 is available in five steps from Diels-Alder reaction of trans-piperylene and maleic anhydride followed by side-chain homologation. Bromolactonization locks the molecule as bicyclic intermediate 6. Esterification, reductive dehalogenation (H_2 /Raney Ni; $Cr(OAc)_2$), base opening of the lactone, careful esterification (CH_2N_2), and dehydration with methanesulfonyl chloride gives 7. The net result is movement of the double bond of 5. Treatment of 7 with NaH gives a fortunately unidirectional Dieckmann ring closure; alkylation with methyl ω -iodoheptanoate introduces the requisite saturated sidechain; lithium iodide-collidine treatment saponifies the ester during the course of which the extra carboxy group is lost; the side-chain methyl ester linkage is restored with diazomethane and the future keto group is protected by reaction with ethylene glycol and acid to give intermediate 8. Next, periodate-permanganate oxidation cleaves the double bond and leads to a methyl ketone whereupon the requisite trans-stereochemistry is established. Diazomethane esterification followed by Bayer-Villiger oxidation introduces the future C-11 α hydroxyl group protected as the acetate. The dioxolane moiety at the future C-9 prevents β -elimination of the acetoxy group of 9. In order to shorten the three-carbon sidechain, methoxide removes the acetyl group so that t-BuOK can close the lactone ring. NaH catalyzed condensation with methyl formate produces inter-

mediate 10. Ozonization removes one carbon atom and acetic anhydride is used to form enolacetate 11, which intermediate is now ready for excision of another carbon. Periodate-permanganate oxidation followed by ethylenediamine hydrolysis produces the needed aldehyde linkage, and the remainder of the synthesis is rather straightforward. Horner-Emmons condensation produces ketone 12 which is sequentially protected with trimethylsilyl chloride, and reduced with sodium borohydride, the isomers separated, and then the blocking groups are removed by base and then acid treatment to give alprostadiol(4).






A conveniently short synthesis of alprostadiol begins with a mixed aldol assembly of the requisite cyclopentenone 13.³ This product is then oxidatively cleaved with periodate-permanganate and the alcohol moiety is protected as the tetrahydropyranyl ether (14). Aqueous chromous sulfate satisfactorily reduces the olefinic linkage and the trans stereoisomer 15 predominates after work-up. The remainder of the synthesis of 4 involves the usual steps, through 16 to 4, with the exception that the xyl tetrahydrolimonyllithium borohydride is used to reduce the C-15 keto moiety so as to produce preferentially the desired C-15S stereochemistry.




Consonant with the present interest in chiral synthesis, two additional contributions can be cited. Sih *et al.*⁴ utilized a combined microbiological and organic chemical sequence in which key chirality establishing steps include the conversion of 17 to chiral, but unstable, 18 by enzymic reduction using the fungus Diplodascus uninucleatus. Lower side-chain synthon 20 was prepared by reduction of achiral 19 with Pencillium decumbens.



Stork and Takahashi⁵ took D-glyceraldehyde synthon 21 from the chiral pool and condensed it with methyl oleate, using lithium diisopropylamide as catalyst for the mixed aldol reaction, leading to 22. The olefinic linkage is a latent form of the future carboxyl group. Protection of the diastereoisomeric mixture's hydroxyl by a methoxymethyleneoxo ether (MEMO) group and sequential acid treatments lead to β -lactone 23. This is tosylated, reduced to the lactol with dibal, and converted to the cyanohydrin (24). Ethylvinyl ether is used to cover the hydroxyl groups and then sodium hexamethyldisilazane treatment is used to express the nucleophilicity of the cyanohydrin ether, an umpohlung reagent for aldehydes that Stork has introduced. This internal displacement gives cyclopentane derivative 25. Periodate-permanganate oxidation cleaves the


olefinic linkage, the ether groups are removed by dilute acid,




and diazomethane leads to the ester. The other protecting groups are removed to give chiral 26, which was already well known in its racemic form as a prostaglandin synthon.

A significant deactivating metabolic transformation of natural prostaglandins is enzymic oxidation of the C-15 hydroxyl to the corresponding ketone. This is prevented, with retention of activity, by methylation to give the C-15 tertiary carbinol series. This molecular feature is readily introduced at the stage of the Corey lactone (27) by reaction with methyl Grignard reagent or trimethylaluminum. The resulting mixture of tertiary carbinols (28) is transformed to oxytocic carba-prost (29) by standard transformations, including separation of diastereoisomers, so that the final product is the C-15 (R) analogue. This diastereoisomer is reputedly freer of typical prostaglandin side effects than the C-15 (S) isomer.⁶

Carbaprost can be converted to the metabolically stable




prostaglandin E analogue, arbaprostil (31), which exerts anti-secretory and cytoprotective activity in the stomach following oral administration and so promotes ulcer healing. At -45°C , selective silanization of the methyl ester of carbaprost gives 30, which undergoes Collins oxidation and acid catalyzed de-blocking to produce arbaprostil (31).⁶ The stereochemical configuration of the drug was confirmed by x-ray analysis. The branched alcoholic moiety can also be introduced by suitable modifications in the Horner-Emmons reaction.⁷



Another device for inhibiting transformation by lung prostaglandin-15-dehydrogenase is introduction of gem-dimethyl branching at C-16. This stratagem was not sufficient, however, to provide simultaneously the necessary chemical stability to allow intravaginal administration in medicated devices for the purpose of inducing labor or abortion. It was found that this could be accomplished by replacement of the C-9 carbonyl group by a methylene (a carbon bioisostere) and that the resulting


agent, meteneprost (33), gave a lower incidence of undesirable gastrointestinal side effects as compared with intramuscular injection of carbaprost (29) methyl ester. The synthesis⁸ utilizes the sulfur ylide prepared from N,S-dimethyl-S-phenylsulfoxime and methyl Grignard (32a). This reacts with 16,16-dimethylprostaglandin E₂ methyl ester bis-(trimethylsilyl) ester (32). The resulting β -hydroxysulfoximine undergoes olefination on reduction with aluminum amalgam⁹ and deblocking produces the uterine stimulant meteneprost (33).



Among the other metabolic transformations that result in loss of prostaglandin activity is ω -chain oxidative degradation. A commonly employed device for countering this is to use an aromatic ring to terminate the chain in place of the usual aliphatic tail. Further, it is known in medicinal chemistry that a methanesulfonimide moiety has nearly the same pK_a as a carboxylic acid and occasionally is biologically acceptable as well as a bioisostere. These features are combined in the uterine stimulant, sulprostone (39). Gratifyingly these changes also result in both enhanced tissue selectivity toward the uterus and lack of dehydration by the prostaglandin-15-dehydrogenase.

The synthesis follows closely along normal prostaglandin


lines with the variations being highlighted here. Processed Corey lactone 34 undergoes Horner-Emmons trans olefination with ylide 35 to introduce the necessary features of the desired ω -side chain (36). After several standard steps, intermediate 37 undergoes Wittig cis-olefination with reagent 38 and further standard prostaglandin transformations produce sulprostone (39).¹⁰



Thromboxane A_2 , formed in blood platelets, is a vasoconstrictor with platelet aggregating action whereas prostacyclin, epoprostenol (43), formed in the lining cells of the blood vessels, is a vasodilator that inhibits platelet aggregation. Their biosynthesis from arachadonic acid via the prostaglandin cascade is normally in balance so that they together exert a sort of yin-yang balancing relationship fine tuning vascular homeostasis. The importance of this can hardly be overestimated. Thrombosis causes considerable morbidity and mortality in advanced nations through heart attacks, stroke, pulmonary

embolism, thrombophlebitis, undesirable clotting associated with implanted medical devices, and the like. Impairment of vascular prostacyclin synthesis can well result in pathological hypertension and excess tendency toward forming blood clots. Administering exogenous prostacyclin, epoprostenol (43), shows promise in combating these problems even though the drug is not active if given orally and is both chemically and metabolically unstable so that continuous infusion would seem to be needed for normal maintenance therapy.

The drug is conveniently synthesized from prostaglandin 12α methyl ester (40), which undergoes oxybromination in the presence of potassium triiodide to give 41. Treatment with DBN




(diazabicyclo[4.3.0]non-5-ene) gives dehydrohalogenation to enol ether 42. Careful alkaline hydrolysis gives the sodium salt of epoprostenol (43).¹¹ The free acid is extremely unstable, presumably due to the expected acid lability of enol ethers.

Much chemical attention is currently devoted to finding chemically stable analogues of 43; Volume 4 will surely have much to say about this.

b. Retenoids


The discovery that some retinoids possess prophylactic activity against carcinogenesis in epithelial tissues¹² has reawakened

interest in these terpene derivatives, particularly in 13-cis-retinoic acid (isotretinoin, 48) which is relatively potent and nontoxic. Isotretinoin also has keratolytic activity of value in the treatment of severe acne. The synthesis^{13,14} is complicated by ready isomerization, and some early confusion existed in the literature regarding the identity of some intermediates. The natural terpene β -ionone (44) is subjected to a Reformatsky reaction with zinc and ethyl bromoacetate and the resulting product is reduced to the allylic alcohol with lithium aluminum hydride and then oxidized to trans-(β -ionylidene)acetaldehyde (45). This is condensed in pyridine with β -methylglutaconic anhydride to give 46. Careful saponification gives mainly diacid 47 which, on heating with copper and quinoline, decarboxylates to isotretinoin (48).^{13,14}



The keratolytic analogue motretinide (53) is effective in treating acne and the excess epithelial growth characteristic

of psoriasis, demonstrating that an aromatic terminal ring is compatible with activity. The synthesis¹⁵ passes through the related orally active antipsoriatic/antitumor agent, etrinitate (52). These synthetic compounds have a wider safety margin than the natural materials. Etrinitate is synthesized¹⁶ from 2,3,5-trimethylanisole by sequential chloromethylation (HCl and formaldehyde) to 50 followed by conversion to the ylid (51) with triphenylphosphine. Wittig olefination then leads to etrinitate (52). Etrinitate may then be saponified, activated by PCl_3 to the acid chloride, and then reacted with ethylamine to give motretinide (53).




The retinoids share with certain steroid hormones the distinction of belonging to the few classes of substances capable of powerful positive influence on cell growth and differentiation.


c. Miscellaneous

In building their characteristic cell walls, bacteria utilize D-alanine which they must manufacture enzymatically by epimerization of the common protein constituent, L-alanine, taken up in their diet. Because mammals have neither a cell wall nor an apparent need for D-alanine, this process is an attractive target for chemotherapists. Thus there has been developed a

group of mechanism-based inhibitors of alanine racemase. The principle utilized in their design is that the enzyme would convert an unnatural substrate of high affinity into a reactive Michael acceptor which would then react with the enzyme to form a covalent bond and inactivate the enzyme. Being unable to biosynthesize an essential element of the cell wall, the organism so affected would not be able to grow or repair damage. It was hypothesized that a strategically positioned halo atom would eliminate readily in the intermediate pyridoxal complex (54) to provide the necessary reactive species. A deuterium atom at the α -carbon is used to adjust the rate of the process




so that the necessary reactions occur at what is judged to be the best possible pace. The process is shown schematically above (54 to 56) for the drug fludalanine (56). In practice, the drug is combined with the 2,4-pentanedione enamine of cycloserine. The combination is synergistic as cycloserine inhibits the same enzyme, but by a different mechanism.



One of the syntheses of fludalanine begins with base promoted condensation of ethyl fluoroacetate and ethyl oxalate to give 57. This is then converted by hydrolytic processes to the insoluble hydrated lithium salt of fluoropyruvate (58). This last is reductively aminated by reduction with sodium borodeuteride and the resulting racemate is resolved to give D-fludalanine (59).¹⁷

There is a putative relationship between the pattern of certain lipids in the bloodstream and pending cardiovascular accidents. As a consequence, it has become a therapeutic objective to reduce the deposition of cholesterol esters in the inner layers of the arterial wall. One attempts through diet or the use of prophylactic drug treatments to reduce the amount of very low density lipoproteins without interfering with high density lipoproteins in the blood. The latter are believed to be beneficial for they transport otherwise rather water insoluble cholesterol. Clofibrate, one of the main hypocholesterolemic drugs, has been shown to have unfortunate side effects in some patients so alternatives have been sought. Gemcadiol (62) is one of the possible replacements. This compound may be synthesized by alkylating two molar equivalents of the cyclohexylamine imine of isopropanal (60) with 1,6-dibromohexane under the influence of lithium diisopropylamide. The resulting dialdehyde (61) is reduced to gemcadiol (62) with sodium boro-



hydride.¹⁸ There is evidence that gemcadiol is metabolically converted to diacid (63) which is believed to be the active agent at the cellular level.


REFERENCES

1. T. J. Schaff and E. J. Corey, J. Org. Chem., **37**, 2921 (1972).
2. H. L. Slates, Z. S. Zelawski, D. Taub and N. L. Wendler, Tetrahedron, **30**, 819 (1974).
3. M. Miyano and M. A. Stealey, J. Org. Chem., **40**, 1748 (1975).
4. C. J. Sih, R. G. Salomon, P. Price, R. Sood and G. Peruzzotti, J. Am. Chem. Soc., **97**, 857 (1975); C. J. Sih, J. B. Heather, R. Sood, P. Price, G. Peruzzotti, L. F. Hsu Lee, and S. S. Lee, ibid., 865.
5. G. Stork and T. Takahashi, J. Am. Chem. Soc., **99**, 1275 (1977).
6. E. W. Yankee, U. Axen, and G. L. Bundy, J. Am. Chem. Soc., **96**, 5865 (1974).
7. E. W. Yankee and G. L. Bundy, J. Am. Chem. Soc., **94**, 3651 (1972); G. Bundy, F. Lincoln, N. Nelson, J. Pike, and W. Schneider, Ann. N.Y. Acad. Sci., **76**, 180 (1971).
8. F. A. Kimball, G. L. Bundy, A. Robert, and J. R. Weeks, Prostaglandins, **17**, 657 (1979).
9. C. R. Johnson, J. R. Shanklin, and R. A. Kirchoff, J. Am. Chem. Soc., **95**, 6462 (1973).
10. T. K. Schaff, J. S. Bindra, J. F. Egger, J. J. Plattner, J. A. Nelson, M. R. Johnson, J. W. Constantine, H.-J. Hess, and W. Elger, J. Med. Chem., **24**, 1353 (1981).
11. R. A. Johnson, F. H. Lincoln, E. G. Nidy, W. P. Schneider, J. L. Thompson, and U. Axen, J. Am. Chem. Soc., **100**, 7690 (1978).

12. D. L. Newton, W. R. Henderson, and M. B. Sporn, Cancer Res., 40, 3413 (1980).
13. C. D. Robeson, J. D. Cawley, L. Weister, M. H. Stern, C. C. Eddinger, and A. J. Chechak, J. Am. Chem. Soc., 77, 41111 (1955).
14. A. H. Lewin, M. G. Whaley, S. R. Parker, F. I. Carroll, and C. G. Moreland, J. Org. Chem., 47, 1799 (1982).
15. W. Bollag, R. Rueegg, and G. Ryser, Swiss Patent 616,134 (1980); Chem. Abstr., 93, 71312j (1980).
16. W. Bollag, R. Rueegg, and G. Ryser, Swiss Patent 616,135 (1980); Chem. Abstr., 93, 71314m (1980).
17. U.-H. Dolling, A. W. Douglas, E. J. J. Grabowski, E. F. Schoenewaldt, P. Sohar, and M. Sletzing, J. Org. Chem., 43, 1634 (1978).
18. G. Moersch and P. L. Creger, U.S. Patent 3,929,897 (1975); Chem. Abstr., 85, 32426q (1976).

2 Phenethyl and Phenoxypropanolamines


The phenylethanolamine derivatives epinephrine (1) and nor-epinephrine (2) are intimately associated with the sympathetic nervous system. These two neurotransmitter hor-



(1) R = CH_3

(2) R = H

(3) R = $\text{CH}(\text{CH}_3)_2$



(4)

mones control many of the responses of this branch of the involuntary, autonomic nervous system. Many of the familiar responses of the "fight or flight" syndrome such as vasoconstriction, increase in heart rate, and the like are mediated by these molecules. The profound biological effects elicited by these molecules have spurred an enormous amount of synthetic medicinal chemistry a better understanding of the

action of the compounds at the molecular level and aimed also at producing new drugs. The availability of analogues of the natural substances interestingly led to the elucidation of many new pharmacological concepts. In spite of the fact that they differ only by an N-methyl group, the actions of epinephrine and norepinephrine are not quite the same. The former tends to elicit a largely inhibiting effect on most responses whereas the latter in general has a permissive action. These trends were accentuated in the close analogues isoproterenol (3) and phenylephrine (4). The pharmacology that led to the division of the sympathetic nervous system into the α - and β -adrenergic branches was put on firmer footing by the availability of these two agents. It may be mentioned in passing that isoproterenol is an essentially pure β -adrenergic agonist whereas phenylephrine acts largely on the α -adrenergic system.


The search for new drugs in this series has concentrated quite closely on their action on the lungs, the heart and the vasculature. Medicinal chemists have thus sought sympathomimetic agents that would act exclusively as bronchodilating agents or as pure cardiostimulant drugs. The adventitious discovery that molecules which antagonize the action of β -sympathomimetic agents - the β -blockers - lower blood pressure has led to a corresponding effort in this field.

1. PHENYLETHANOLAMINES

As noted above, β -adrenergic agonists such as epinephrine typically cause relaxation of smooth muscle. This agent


would thus in theory be useful as a bronchodilator for treatment of asthma; epinephrine itself, however, is too poorly absorbed orally and too rapidly metabolized to be used in therapy. A large number of analogues have been prepared over the years in attempts to overcome these shortcomings. The initial strategy consisted in replacing the methyl group on nitrogen with an alkyl group more resistant to metabolic N-dealkylation. Isoproterenol (3) is thus one of the standbys as a drug for treatment of asthma.

The tertiary butyl analogue, colterol (9) is similarly resistant to metabolic inactivation. (It might be noted that there is some evidence that these more lipophilic alkyl groups, besides providing resistance to inactivation, also result in higher intrinsic activity by providing a better drug receptor interaction.) This drug can in principle be prepared by the scheme typical for phenylethanalamines. Thus acylation of catechol by means of Friedel-Crafts reaction with acetyl chloride affords the ketone 6; this is then halogenated to give intermediate 7. Displacement of bromine by means of tertiary butylamine gives the amino-ketone 8. Reduction of the carbonyl group by catalytic hydrogenation affords colterol (9).




Absorption of organic compounds from the gastrointestinal tract is a highly complex process which involves at one stage passage through a lipid membrane. Drugs that are highly hydrophilic thus tend to be absorbed very inefficiently by reason of their preferential partition into aqueous media. One strategy to overcome this unfavorable distribution consists in preparing a derivative that is more hydrophobic and which will revert to the parent drug on exposure to metabolizing enzymes after absorption. Such derivatives, often called prodrugs, have been investigated at some length in order to improve the absorption characteristics of the very hydrophilic catecholamines.

Acylation of aminoketone 8 with the acid chloride from p-toluic acid affords the corresponding ester (10); catalytic hydrogenation leads to the bronchodilator bitolerol (11)¹. An analogous scheme starting from the N-methyl ketone (12) and pivaloyl chloride gives aminoalcohol (14). This compound is then resolved to isolate the levorotatory isomer². There is thus obtained the drug dipivefrin.




A variant on this theme contains mixed acyl groups. In the absence of a specific reference it may be speculated that the synthesis starts with the diacetyl derivative (15). Controlled hydrolysis would probably give the monoacetate (16) since the ester para to the ketone should be activated by that carbonyl function. Acylation with anisoyl chloride followed by reduction would then afford nisobutanol (18).



Catecholamines are also intimately involved in cardiac function, with β -sympathetic agonists having a generally stimulant action on the heart. Some effort has thus been devoted to the synthesis of agents that would act selectively on the heart. (Very roughly speaking, β^1 -adrenergic receptor agonists tend to act on the heart while β^2 -adrenergic receptor agonists act on the lungs; much the same holds true for antagonists; see below.)


Preparation of the cardiotonic agent butopamine (23) starts with reductive amination of ketone 19. Acylation of the resulting amide (20) with hydroxyacid 21 affords the corresponding amine (22). Treatment with lithium aluminum hydride serves both to reduce the amide and remove the acetyl protecting groups. There is thus obtained butopamine³.



Drugs that block the action of α -adrenergic activation effectively lower blood pressure by opposing the vasoconstricting effects of norepinephrine. Drawbacks of these agents, which include acceleration of heart rate, orthostatic hypotension and fluid retention, were at one time considered to be due to the extension of the pharmacology of α -blockers. Incorporation of β -blocking activity into the molecule should oppose these effects. This strategy seemed particularly promising in view of the fact that β -adrenergic blockers were adventitiously found lower blood pressure in their own right. The first such combined α - and β -blocker, labetalol has confirmed this strategy and proved to be a clinically useful antihypertensive agent.

The drugs in this class share the phenylethanolamine moiety and a catechol surrogate in which the 3-hydroxyl is replaced by some other function that contains relatively acidic protons.


Synthesis of the prototype⁴ begins with Friedel Crafts acetylation of salicylamide (24). Bromination of the ketone (25) followed by displacement with amine 27 gives the corresponding aminoketone (28). Catalytic hydrogenation to the aminoalcohol completes the synthesis of labetolol (24). The presence of two chiral centers at remote positions leads to the two diastereomers being obtained in essentially equal amounts.




In much the same vein, alkylation of bromoketone (26) with amine (30) (obtained by reductive amination of the corresponding ketone) affords aminoketone (31). Catalytic reduction leads to medroxalol (32)⁵.

The methyl group on a sulfoxide interestingly proves sufficiently acidic to substitute for phenolic hydroxyl. The preparation of this combined α - and β -blocker, sulfinalol⁶, begins by protection of the phenolic hydroxyl as its benzoate ester (34). Bromination (35) followed by

condensation with amine 36 gives the aminoketone (37). Successive catalytic reduction and saponification affords the aminoalcohol (38). Oxidation of the sulfide to the sulfoxide with a reagent such as metaperiodate gives sulfinalol (39). This last step introduces a third chiral center because trigonal sulfur exists in antipodal forms. The number of diastereomers is thus increased to eight.




A phenylethanolamine in which the nitrogen is alkylated by a long chain aliphatic group departs in activity from the prototypes. This agent, suloctidil (43) is described as a peripheral vasodilator endowed with platelet antiaggregatory activity. As with the more classical compounds, preparation proceeds through bromination of the substituted propiophenone (40) and displacement of halogen with octylamine. Reduction, in this case by means of sodium borohydride affords suloctidil (43)⁷.









Pharmacological theory would predict that β -adrenergic blockers should oppose the vasodilating action of epinephrine and, in consequence, increase blood pressure. It was found, however, that these drugs in fact actually decrease blood pressure in hypertensive individuals, by some as yet undefined mechanism. The fact that this class of drugs tends to be very well tolerated has led to enormous emphasis on the synthesis of novel β -blockers. The observation that early analogues tended to exacerbate asthma by their blockade of endogenous β -agonists has led to the search for compounds that show a preference for β^1 -adrenergic sites.

With some important exceptions, drugs in this class are conceptually related to the phenylethanalamines by the interposition of an oxymethylene group between the aromatic ring and the benzyl alcohol.





Compounds are prepared by a fairly standard sequence which consists of condensation of an appropriate phenol with epichlorohydrin in the presence of base. Attack of phenoxide can proceed by means of displacement of chlorine to give epoxide (45) directly. Alternatively, opening of the epoxide leads to anion 44; this last, then, displaces halogen on the adjacent carbon to lead to the same epoxide. Reaction of the epoxide with the appropriate amine then completes the synthesis.

Application of this scheme to *o*-cyclopentyl phenol, *o*-cyclohexylphenol and *m*-cresol thus leads to respectively, penbutolol (47)⁸, exapralol (48)⁹ and bevantolol (49)¹⁰. The phenoxypanolamine tipropidil (52) interestingly exhibits much the same biological activity as its phenylethanolamine parent suloctidil (53).





The phenol (55) required for preparation of diacetolol (50)¹¹ can be obtained by Friedel-Crafts acetylation of *p*-acetamidophenol. The starting material (58) for pamatolol (51)¹² can be derived from *p*-hydroxyphenylacetone nitrile (56) by reduction to the amine (57) followed by treatment with ethyl chloroformate. Bucindolol (52) is one of the newer β -blockers designed to incorporate non-adrenergically mediated vasodilating activity in the same molecule as the adrenergic blocker. Preparation of the amine (61) for this

agent starts by displacement of the dimethylamino group in gramine (59) by the anion from 2-nitropropane. Reduction of the nitro group leads to the requisite intermediate¹³.



Synthesis of primidolol (65)¹⁴ can be carried out by a convergent scheme. One branch consists in application of the usual scheme to o-cresol (62); ring opening of the intermediate oxirane with ammonia leads to the primary amine (63). The side chain fragment (64) can be prepared by alkylation of pyrimidone (63) with ethylene dibromide to afford 64. Alkylation of aminoalcohol 62 with halide 64 affords primidolol.






It is by now well accepted that most drugs, particularly those whose structures bear some relation to endogenous agonists owe their effects to interaction with biopolymer receptors. Since the latter are constructed from chiral subunits (amino acids, sugars, etc.), it should not be surprising to note that drugs too show stereoselectivity in their activity. That is, one antipode is almost invariably more potent than the other. In the case of the adrenergic agonists and antagonists, activity is generally associated with the R isomer. Though the drugs are, as a rule, used as racemates, occasional entities consist of single enantiomers. Stereospecific synthesis is, of course, preferred to resolution since it does not entail discarding half the product at the end of the scheme.


Prenalterol (73) interestingly exhibits adrenergic agonist activity in spite of an interposed oxymethylene group. The stereospecific synthesis devised for this molecule¹⁵ relies on the fact that the side chain is very

similar in oxidation state to that of a sugar. Condensation of the monobenzyl ether of phenol 66 with the epoxide derived from D-glucofuranose (67) affords the glycosylated derivative (68). Hydrolytic removal of the protecting groups followed by cleavage of the sugar with periodate gives aldehyde 69. This is in turn reduced to the glycol by means of sodium borohydride and the terminal alcohol is converted to the mesylate (71). Displacement of that group with isopropylamine (72) followed by hydrogenolytic removal of the O-benzyl ether affords the β^2 - selective adrenergic agonist prenalterol (73).




Formal cyclization of the hydroxyl and amine functions to form a morpholine interestingly changes biological act-

ivity markedly; the resulting compound shows CNS activity as an antidepressant rather than as an adrenergic agent. Reaction of epoxide (74) with the mesylate from ethanolamine leads to viloxazine (76) in a single step¹⁶. It is likely that reaction is initiated by opening of the oxirane by the amino group. Internal displacement of the leaving group by the resulting alkoxide forms the morpholine ring.



The widely used tricyclic antidepressant drugs such as imipramine and amitriptyptiline have in common a series of side effects that limit their safety. There has thus occasioned a wide search for agents that differ in structure and act by some other mechanism. Nisoxetine and fluoxetine are two nontricyclic compounds which have shown promising early results as antidepressants. Mannich reaction on acetophenone leads to the corresponding aminoketone (78). Reduction of the carbonyl group (79) followed by replacement of the hydroxyl by chlorine gives intermediate 80. Displacement of chlorine with the alkoxide from the monomethyl ether of catechol gives the corresponding aryl

ether (81). The amine is then dealkylated to the monomethyl derivative by the von Braun sequence (cyanogen bromide followed by base) to give nisooxetine (82). Displacement on (80) with the monotrifluoromethyl ether from hydroquinone followed by demethylation leads to fluoxetine (84)¹⁷.



REFERENCES

1. M. Minatoya B. F. Tullar and W. D. Conway, U.S. Patent 3,904,671; Chem. Abstr. 84, 16943e (1976).
2. A. Hussain and J. E. Truelove, German Offen. 2,343,657; Chem. Abstr. 80, 145839s (1974).
3. J. Mills, K. K. Schmiegel and R. R. Tuttle, Eur. Patent Appl. 7,205 (1980); Chem. Abstr. 93, 94972 (1980).
4. L. H. C. Lunts and D. T. Collin, German Offen. 2,032,642; Chem. Abstr. 75, 5520c (1971).
5. J. T. Suh and T. M. Bare, U.S. Patent 3,883,560; Chem. Abstr. 83, 78914j (1975).
6. Anon. British Patent 1,544,872; Chem. Abstr. 92, 163686s (1980).
7. G. Lambelin, J. Roba, and C. Gillet, German Offen. 2,344,404; Chem. Abstr. 83, 97820 (1975).
8. G. Haertfelder, H. Lessenich and K. Schmitt, Arzneim. Forsch. 22, 930 (1972).
9. M. Carissimi, P. Gentili, E. Grumelli, E. Milla, G. Picciola and F. Ravenna, Arzneim. Forsch. 26, 506 (1976).
10. M. Ikezaki, K. Irie, T. Nagao, and K. Yamashita, Japanese Patent 77, 00234; Chem. Abstr. 86, 1894767 (1977).
11. K. R. H. Wooldridge and B. Berkley, South African Patent 68 03,130; Chem. Abstr. 70, 114824 (1969).
12. A. E. Brandstrom, P. A. E. Carlsson, H. R. Corrodi, L. Ek and B. A. H. Ablad, U.S. Patent 3,928,601; Chem. Abstr. 85, 5355j (1976).

13. W. E. Kreighbaum, W. L. Matier, R. D. Dennis, J. L. Minielli, D. Deitchman, J. L. Perhach, Jr. and W. T. Comer, J. Med. Chem., 23, 285 (1980).
14. J. Augstein, D. A. Cox and A. L. Ham., German Offen. 2,238,504 (1973). Chem. Abstr. 78, 136325e (1973).
15. K. A. Jaeggi, H. Schroeter, and F. Ostermayer, German Offen. 2,503,968; Chem. Abstr. 84, 5322 (1976).
16. S. A. Lee, British Patent 1,260,886; Chem. Abstr. 76, 99684e (1972).
17. B. B. Molloy and K. K. Schmiegel, German Offen. 2,500,110; Chem. Abstr. 83, 192809d (1975).

3 Arylaliphatic Compounds


The aromatic portion of the molecules discussed in this chapter is frequently, if not always, an essential contributor to the intensity of their pharmacological action. It is, however, usually the aliphatic portion that determines the nature of that action. Thus it is a common observation in the practice of medicinal chemistry that optimization of potency in these drug classes requires careful attention to the correct spatial orientation of the functional groups, their overall electronic densities, and the contribution that they make to the molecule's solubility in biological fluids. These factors are most conveniently adjusted by altering the substituents on the aromatic ring.

1. ARYLACETIC ACID DERIVATIVES


The potent antiinflammatory action exerted by many arylacetic acid derivatives has led to the continued exploration of this class. It is apparent from a consideration of the structures of compounds that have become prominent that considerable structural latitude is possible without loss of activity.

The synthesis of fenclofenac (5), a nonsteroidal antiinflammatory agent (NSAI), starts with condensation of o-chloro-

acetophenone (1) and 2,4-dichlorophenol (2) under Ullmann conditions (Cu/NaOH). The unsymmetrical diarylether (3) is subjected to the Willgerodt-Kindler reaction to give thioamide 4. This last is saponified to produce fenclofenac (5).¹




A structure more distantly related to these is amfenac (10). Like most of the others, amfenac, frequently used after tooth extraction, is an antiinflammatory agent by virtue



of inhibition of the cyclooxygenase enzyme essential for prostaglandin biosynthesis. The synthesis begins with hydrazone (7) formation between phenylacetone and 1-aminoindolin-2-one (6) by warming in acetic acid. Treatment with HCl/EtOH results in a Fischer indole rearrangement to produce 8. Ozonolysis produces unsymmetrical diarylketone 9 which is converted to an intermediate indolin-2-one by cleavage of the ester and acetamide moieties with HCl and then lactam hydrolysis with NaOH to give amfenac (10).²

A closet member of this little group, revelation of whose real nature requires metabolic transformation of an acetylenic linkage to an acetic acid moiety, is fluretofen (14). The synthesis begins by Friedel-Crafts acylation of 2-fluorobiphenyl (11) with acetic acid to give ketone 12. Heating with $\text{PCl}_5/\text{POCl}_3$ then produces the α -chlorostyrene analogue 13. Dehydrohalogenation with strong base (NaNH_2) produces the acetylene, fluretofen (14). Metabolic studies reveal facile bioconversion to the arylacetic acid analogue 15. This is an active antiinflammatory agent when administered directly and so is believed to account for the activity of fluretofen. A likely pathway for this transformation involves terminal oxygen-




ation of the acetylenic moiety, which product would then tautomerize to the ketene. Spontaneous hydration of the latter would complete the sequence.³

Phenylacetamides have a variety of pharmacological actions depending upon the nature of the amine-derived component.

Guanfacine (17), an antihypertensive agent acting as a central α -adrenergic receptor agonist, requires administration only once daily and reportedly has fewer CNS sideeffects than the somewhat related drug, clonidine. Guanfacine is prepared readily by ester-amide exchange of methyl 2,6-dichlorophenyl acetate (16) using guanidine.⁴





Use of a large, lipophilic nitrogenous component results in a lidocaine like, local anesthetic type cardiac antiarrhythmic drug, lorcainide (20). Synthesis begins with the Schiff's base (18) derived by reaction of p-chloro-aniline and borohydride followed by acylation with phenylacetyl chloride produces amide 19. Selective hydrolysis with HBr followed by alkylation with isopropyl bromide completes the synthesis of lorcainide (20).⁵

The structural requirements for such activity are not very confining, as can be seen in part by comparing the structure of lorcainide with oxiramide (21). Antiarrhythmic oxiramide (21) is made by a straightforward ester-amide exchange reaction




involving ethyl 2-phenoxyphenylacetate and 4-cis-(2,6-dimethyl-piperidino)butylamine.⁶



Introducing yet more structural complexity into the amine component leads to the antiarrhythmic agent disobutamide (24). Disobutamide is structurally related to disopyramide (25)⁷ but is faster and longer acting. The synthesis of 24 begins with sodamide induced alkylation of 2-chlorophenylacetonitrile with 2-diisopropylaminoethyl chloride to give 22. A second sodamide mediated alkylation, this time with 2-(1-piperidino)ethyl chloride, gives nitrile 23. Subsequent sulfuric acid hydration completes the synthesis of disobutamide (24).⁸










The treacheries inherent in naive attempts at pattern recognition are illustrated by the finding that ester 28, known as cetiedil, is said to be a peripheral vasodilator. Clemmensen reduction of Grignard product 26 removes the superfluous benzylic hydroxyl group and esterification of the sodium salt of the resulting acid (27) with 2-(1-cycloheptylamino)ethyl chloride produces cetiedil (28).⁹



An interesting biphenyl derivative utilizing a bioisosteric replacement for a carboxyl group is the antidiarrheal agent, nufenoxole (34). To get around addictive and analgesic side effects associated with the classical morphine based antidiarrheal agents, a different class of drug was sought. Nufenoxole has few analgesic, anticholinergic, or central effects. Reduction with a ruthenium catalyst (to prevent hydrogenolysis) converts p-aminobenzoic acid to cyclohexane derivative 29. Internal N-acetylation of the cis isomer followed by heating gives bicyclic lactam 30. Hydride reduction to the isoquinuclidine and alkylation gives 2-azabicyclo[2.2.2]octane synthon 31. This is used to alkylate diphenylacetone nitrile to give 32. Cycloaddition of sodium azide (ammonium chloride and DMF) gives the normal carboxyl bioisosteric tetrazolyl analogue 33. The synthesis of antidiarrheal nufenoxole is completed by heating





with acetic anhydride to give the 2-methyl-1,3,4-oxadiazol-5-yl analogue.^{10, 11} The mechanism of this rearrangement is believed to involve N-acetylation (35) with subsequent ring opening to the diazoalkane (36) which loses nitrogen to give carbene 37, which cyclizes to the oxadiazole (38).¹²



A phenylacetone nitrile derivative, closantel (41), is an anthelmintic agent useful against sheep liver flukes. Its patented synthesis involves a Schotten-Baumann amidation

between acid chloride 39 and complex aniline 40 to give closantel (41).¹³

A cinnamoylamide, cinromide (44), is a long-acting anti-convulsant similar in its clinical effects to phenacetamide but is less hepatotoxic. The synthesis involves the straightforward amidation of acid 42 via the intermediate acid chloride (SOCl_2) 43. It appears that the drug is mainly deethylated in vivo to give active amide 45.¹⁴



2,2-Disubstituted aryloxyacetic acid derivatives related to clofibrate have been intensively studied in an attempt to get around the side effects of the latter drug.


Ciprofibrate (48), a more potent lipid-lowering agent than clofibrate, is prepared from Simmons-Smith product 46 by Sandmeyer replacement of the amino group by a hydroxyl via the diazonium salt. Phenol 47 undergoes the Reimer-Thiemann like process common to these agents upon alkaline treatment with acetone and chloroform to complete the synthesis of ciprofibrate (48).¹⁵

Further indication that substantial bulk tolerance is available in the para position is given by the lipid lowering agent bezafibrate (50). The p-chlorobenzamide of tyramine (49) undergoes a Williamson ether synthesis with ethyl 2-bromo-

/-methylpropionate to complete the synthesis. The ester group is hydrolyzed in the alkaline reaction medium.¹⁶

Apparently a substantial spacer is also allowable between the aromatic ring and the carboxy group. Gemfibrozil (52), a hypotriglyceridemic agent which decreases the influx of steroid into the liver, is a clofibrate homologue. It is made readily by lithium diisopropylamide-promoted alkylation of sodium isopropionate with alkyl bromide 51.¹⁷


A rather distantly related analogue incorporating a β -dicarbonyl moiety as a bioisosteric replacement for a carboxyl, arildone (55), blocks the uncoating of polio virus and herpes simplex virus type I and thus inhibits infection of cells and the early stages of virus replication. Thus effective therapy would require careful timing as it does with amantidine. Alkylation of phenol 53 with 1,6-dibromohexane gives haloether




54. Finkelstein reaction with sodium iodide is followed by acylation of heptane-3,5-dione to complete the synthesis of arildone (55).¹⁸

2. ANILINES, BENZYL AMINES, AND ANALOGUES


An orally active local anesthetic agent that can be used as an antiarrhythmic agent is meobentine (57). Its patented synthesis starts with p-hydroxyphenylnitrile and proceeds by dimethyl sulfate etherification and Raney nickel reduction to 56. Alkylation of S-methyl-N,N'-dimethylthiourea with 56 completes the synthesis of meobentine (57).¹⁹




Bepridil (59) blocks the slow calcium channel and serves as an antianginal agent and a vasodilator. In its synthesis, alcohol 58 (derived from epichlorohydrin) is converted to the corresponding chloride with thionyl chloride and displaced with the sodium salt of N-benzylaniline to give bepridil (59)²⁰



A number of quaternary amines are effective at modulating nerve transmissions. They often have the disadvantage of being relatively nonselective and so possess numerous sideeffects. This contrasts with the advantage that they do not cross the blood-brain barrier and so have no central sideeffects. Clofilium phosphate (63) is such an antiarrhythmic agent. It is synthesized from ester 60 by saponification followed by Clemmensen reduction and amide formation (oxalyl chloride followed by n-heptylamine) to give 61. Diborane reduction gives secondary amine 62. Reaction with acetyl chloride followed by another diborane reduction gives the tertiary amine. Finally, reaction with ethyl bromide and ion exchange with phosphate complete the synthesis of clofilium phosphate (63).²¹



Another quaternary antiarrhythmic agent is emilium tosylate (65). It is synthesized simply by quaternization of m-methoxybenzyl chloride (64) with dimethylethylamine followed by ion exchange.²²




3. DIARYLMETHANE ANALOGUES


Prenylamine (66) was long used in the treatment of angina pectoris, in which condition it was believed to act by inhibiting the uptake and storage of catecholamines in heart tissue. Droprrenilamine (69), an analogue in which the phenyl ring is reduced, acts as a coronary vasodilator. One of several syntheses involves simple reductive alkylation of 1,1-diphenylpropylamine (67) with cyclohexylacetone (68).²³

Drobuline (71) is a somewhat related cardiac-directed drug with antiarrhythmic action. Since both enantiomers have the


same activity, it is likely that its pharmacological action is due to a local anesthetic-like action. It is synthesized by sodium amide mediated alkylation of diphenylmethane with allyl bromide to give 70. Epoxidation with m-chloroperbenzoic acid followed by opening of the oxirane ring at the least hindered carbon by isopropylamine completes the synthesis.²⁴




A slightly more complex antiarrhythmic agent is pirmentol (74). It is synthesized from 4-chloropropiophenone (72) by keto group protection as the dioxolane (with ethylene glycol and acid) followed by sodium iodide-mediated alkylation with cis 2,6-dimethylpiperidine to give 73. Deblocking with acid followed by addition of 2-lithiopyridine completes the synthesis of pirmentol (74).²⁵



For many years after the discovery of the antidepressant activity of phenothiazine, almost all synthetic activity centered about rigid analogues. Recently attention has been paid to less rigid molecules in part because of the finding that zimelidine (77) is an antidepressant showing selective inhibition of the central uptake of 5-hydroxytryptamine and that it possesses less anticholinergic activity than amitriptylene. One of a number of syntheses starts with p-bromoacetophenone and a Mannich reaction (formaldehyde and dimethylamine) to give aminoketone 75. Reaction with 3-lithio-pyridine gives tertiary carbinol 76. Dehydration with sulfuric acid gives a mixture of Z and E forms of which the Z analogue is the more active.²⁶




Pridefine (80) is a somewhat structurally related antidepressant. It is a centrally active neurotransmitter blocking agent. It blocks norepinephrine in the hypothalamus but does not affect dopamine or 5-hydroxytryptamine. Its synthesis begins by lithium amide-promoted condensation of diethyl succinate and benzophenone followed by saponification to 78. Heating in the presence of ethylamine gives N-ethylsuccinimide 79. Lithium aluminum hydride reduction completes the synthesis of pridefine (80).²⁷



4. STILBENE ANALOGUES


Cells from tissues associated with primary and secondary sexual characteristics are under particular endocrine control. Sex hormones determinethe growth, differentiation, and proliferation of such cells. When a tumor develops in such tissues, it is sometimes hormone dependent and the use of antihormones removes the impetus for the tumor's headlong growth. Many non-steroidal compounds have estrogenic activity; diethylstilbestrol (81) may be taken as an example. Certain more bulky an-



alogues are antagonists at the estrogenic receptor level and exert a second order anti-tumor response.

Nitromifene (85) is such an agent. A Grignard reaction of arylether 82 and ketone 83 leads to tertiary carbinol 84. Tonic acid dehydration leads to a mixture of Z and E stilbenes which constitute the antiestrogen, nitromifene (85).²⁸

Another example is tamoxifen (89). Its synthesis begins with Grignard addition of reagent 86 to aryl ketone 87 giving carbinol 88. Dehydration leads to the readily separable Z and E analogues of 89. Interestingly, in rats the Z form is an antiestrogen whereas the E form is estrogenic. Metabolism involves p-hydroxylation and this metabolite (90) is more potent than tamoxifen itself. In fact, metabolite 90 may be the active form of tamoxifen (89).²⁹



REFERENCES

1. D. C. Atkinson, K. E. Godfrey, B. J. Jordan, E. C. Leach, B. Meek, J. D. Nichols, and J. F. Saville, J. Pharm. Pharmacol., **26**, 357 (1974).
2. W. J. Welstead, Jr., H. W. Moran, H. F. Stauffer, L. B. Turnbull, and L. F. Sancillo, J. Med. Chem., **22**, 1074 (1979).
3. W. B. Lacefield and W. S. Marshall, U.S. Patent 3,928,604; H. R. Sullivan, P. Roffey, and R. E. McMahon, Drug Metab. Disposn., **7**, 76 (1979).
4. J. B. Bream, H. Lauener, C. W. Picard, G. Scholtysik, and T. G. White, Arzneim. Forsch., **25**, 1477 (1975).
5. H. K. F. Hermans and S. Sanczuk, U.S. Patent 4,197,303 (1975); Chem. Abstr., **93**, 132380d (1980).
6. Anon., Netherlands Patent, 6,605,452 (1962); Chem. Abstr., **66**, 104914e (1967).
7. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 2, Wiley, New York, 1980, p. 81.
8. P. K. Youan, R. L. Novotney, C. M. Woo, K. A. Prodan, and F. M. Herschenson, J. Med. Chem., **23**, 1102 (1980).
9. M. Robba and Y. LeGuen, Eur. J. Med. Chem., **2**, 120 (1967).
10. G. W. Adelstein, C. H. Yen, E. Z. Dajani, and R. G. Biandi, J. Med. Chem., **19**, 1221 (1976).
11. W. Schneider and R. Dillman, Chem. Ber., **96**, 2377 (1963).
12. R. Huisgen, J. Sauer, H. J. Sturm, and J. H. Markgraf, Chem. Ber., **93**, 2106 (1960).
13. M. A. C. Janssen and V. K. Sipido, German Offen., 2,610,837 (1976); Chem. Abstr., **86**, 55186w (1977).
14. E. M. Grivsky, German Offen., 2,535,599 (1976); Chem. Abstr., **84**, 164492x (1976).
15. D. K. Phillips, German Offen., 2,343,606 (1974); Chem.

- Abstr., 80, 133048v (1974).
16. E. C. Witte, K. Stach, M. Thiel, F. Schmidt, and H. Stork, German Offen., 2,149,070 (1973); Chem. Abstr., 79, 18434k (1973).
 17. P. L. Creger, G. W. Moersch, and W. A. Neuklis, Proc. R. Soc. Med., 69, 3 (1976).
 18. G. D. Diana, U. J. Salvador, E. S. Zalay, P. M. Carabateas, G. L. Williams, J. C. Collins, and F. Pancic, J. Med. Chem., 20, 757 (1977).
 19. R. A. Maxwell and E. Walton, German Offen., 2,030,693 (1971); Chem. Abstr., 74, 87660q (1971).
 20. R. Y. Mauvernay, N. Busch, J. Simond, A. Monteil, and J. Moleyre, German Offen., 2,310,918 (1973); Chem. Abstr., 79, 136777x (1973).
 21. B. B. Molloy and M. I. Steinberg, Eur. Pat. Appl., 2,604 (1979).
 22. R. A. Maxwell and F. C. Copp, German Offen., 2,030,692 (1971); Chem. Abstr., 74, 76156d (1971).
 23. M. Carissimi, F. Ravenna, and G. Picciola, German Offen., 2,521,113 (1976); Chem. Abstr., 34, 164388t (1976).
 24. P. J. Murphy, T. L. Williams, J. K. Smallwood, G. Bellamy, and B. B. Molloy, Life Sci., 23, 301 (1978).
 25. R. W. Fleming, German Offen., 2,806,654 (1978); Chem. Abstr., 89, 197346j (1978).
 26. B. Carnmalm, T. De Paulis, T. Hogberg, L. Johansson, M.-L. Persson, S.-O. Thorburg, and B. Ulff, Acta Chem. Scand. B, 36, 91 (1982); J.-E. Backvall, R. E. Nordberg, J.-E. Nystrom, T. Hogberg, and B. Ulff, J. Org. Chem., 46, 3479 (1981).
 27. S. Ohki, N. Ozawa, Y. Yabe, and H. Matsuda, Chem. Pharm. Bull., 24, 1362 (1976).


28. D. J. Collins, J. J. Hobbs, and C. W. Emmens, J. Med. Chem., 14, 952 (1971).
29. D. W. Robertson and J. A. Katzenellenbogen, J. Org. Chem., 47, 2386 (1982).

4 Monocyclic Aromatic Agents

The pharmacological response elicited by monocyclic aromatic agents is a function of the number and spatial arrangement of the functional groups attached to the aromatic ring; this is true of a great many drugs.

1. ANILINE DERIVATIVES


Many local anesthetics have a selective depressant action on heart muscle when given systemically. This is useful in treatment of cardiac arrhythmias, and a lidocaine-like drug with this kind of action is tocainide (2).¹




Part of the reason for ortho substitution in such compounds is to decrease metabolic transformation by enzymic

amide cleavage. Encainide (5) is another embodiment of this concept. Its published synthesis involves acetic anhydride-catalyzed condensation of α -picoline with 2-nitrobenzaldehyde to give 3. N-Methylation followed by catalytic reduction gives piperidine 4. The synthesis concludes by acylation with *p*-methoxybenzoyl chloride to give antiarrhythmic encainide (5).²

When the side chain involves an unsymmetrical urea moiety, muscle relaxant activity is often seen. One such agent, lid-amidine (6) exerts its activity as an antiperistaltic agent. Its synthesis involves the straightforward reaction of 2,6-dimethylphenylisocyanate and N-methylguanidine.³




A cyclized version, xilobam (8), is synthesized from N-methylpyrrolidone by conversion to the imine (7) by sequential reaction with triethyloxonium tetrafluoroborate and then anhydrous ammonia. When this is reacted with 2,6-dimethylphenylisocyanate, the centrally acting muscle relaxant xilobam (8) is formed.⁴




A number of muscle relaxants are useful anthelmintic agents. They cause the parasites to relax their attachment to the gut wall so that they can be eliminated. One such agent is carbantel (9). Its synthesis follows the classic pattern of reaction of 4-chlorophenylisocyanate with n-amylamine.⁵


To prepare another such analogue, N-methylation of N,N-dicarbomethoxythiourea gives 10, which itself reacts with complex aniline analogue 11 to give the veterinary anthelmintic agent felsantel (12).⁶



A simple aniline derivative acts as a prostatic antiandrogen. Its synthesis involves simple acylation of disubstituted aniline 13 with isobutyryl chloride to give flutamide (14).⁷




A phenylguanidine analogue is readily prepared by first reacting 2-imino-N-methylpyrrolidine with phenylisothiocyanate to give synthon 15. This is next S-methylated with methyl iodide to give 16 which itself, on reaction with pyrrolidine, is



converted to the antidiabetic agent pirogilride (17).⁸


Finally, in demonstration of the pharmacological versatility of this chemical subclass, ethyl lodoxamide (20) shows antiallergic properties. It shows a biological relationship with disodium chromoglycate by inhibiting the release of mediators of the allergic response initiated by allergens. It can be synthesized by chemical reduction of dinitrobenzene analogue 18 to the m-diamino analogue 19. This, then, is acylated with ethyl oxaly chloride to complete the synthesis of ethyl lodoxamide (20).⁹



2. BENZOIC ACID DERIVATIVES


It has been documented in an earlier volume that appropriately substituted molecules with two strongly electron withdrawing substituents meta to one another in a benzene ring often possess diuretic properties and, even though the prototypes usually have two substituted sulfonamide moieties so disposed, other groups can replace at least one of them. An example of this is piretanide (24), where one such group is a carboxyl

moiety.¹⁰ The published synthesis starts with highly substituted benzoate 21¹⁰ which is reduced with a Raney nickel catalyst and converted to succinimide 23 by reaction with succinic anhydride.




Reduction to the corresponding N-substituted pyrrolidine (23) takes place with sodium borohydride/boron trifluoride. Saponification completes the synthesis of the diuretic agent pirtamide (24).¹¹

Because of resonance stabilization of the anion, a tetrazolyl moiety is often employed successfully as a bioisosteric replacement for a carboxy group. An example in this subclass is provided by azosemide (27). Benzonitrile analogue 25 is prepared by phosphorus oxychloride dehydration of the corresponding benzamide. Next, a nucleophilic aromatic displacement reaction of the fluorine atom leads to 26. The synthesis concludes with the 1,3-dipolar addition of azide to the nitrile function to produce the diuretic azosemide (27).¹²




Reversal of the amide moiety of local anesthetics is

consistent with retention of activity. So too with the derived antiarrhythmic agents. Flecainide (30) is such a substance. It is synthesized from 2,5-dihydroxybenzoic acid by base-mediated etherification with 2,2,2-trifluoroethanol. If done carefully, ester 28 results. Amide ester exchange with the appropriate pyridine amine analogues leads to 29. Catalytic reduction of the more electron-deficient aromatic ring results in the formation of flecainide (30).¹³




A lipid lowering agent of potential value in hypercholesterolemia is cetaben (31). It is synthesized facilely by monoalkylation of ethyl p-aminobenzoate with hexadecyl bromide and then saponification.¹⁴

Benzamide 33, known as bentiromide, is a chymotrypsin substrate of value as a diagnostic acid for assessment of pancreatic function. It is synthesized by amide formation between



ethyl p-aminobenzoate and N-benzoyl-tyrosine using N-methylmorpholine and ethyl chlorocarbonate for activation. The resulting L-amide (32) is selectively hydrolyzed by sequential


use of dimethyl sodium and dilute acid to give bentiromide (33).¹⁵




3. BENZENESULFONIC ACID DERIVATIVES


As has been discussed previously, substituted p-alkylbenzene-sulfonylureas often possess the property of releasing bound insulin, thus sparing the requirement for insulin injections in adult-onset diabetes.¹⁶ A pyrimidine moiety, interestingly, can serve as a surrogate for the urea function.

Gliflumide (37), one such agent, is synthesized from 4-isobutyl-2-chloro-pyrimidine (34) by nucleophilic displacement using p-sulfonamidobenzeneacetic acid (35) to give sulfonamide 36. Reaction, via the corresponding acid chloride, with S-1-amino-1-(2-methoxy-5-fluorophenyl)ethane completes the synthesis of the antidiabetic agent gliflumide (37).¹⁷






A related agent, glicetanile sodium (42), is made by a variant of this process. Methyl phenylacetate is reacted with chlorosulfonic acid to give 38, which itself readily reacts with aminopyrimidine derivative 39 to give sulfonamide 40. Saponification to acid 41 is followed by conversion to the acid chloride and amide formation with 5-chloro-2-methoxyaniline to complete the synthesis of the hypoglycemic agent glicetanile (42).¹⁸



Perhaps surprisingly, the p-methyl benzenesulfonylurea analogue called tosifen (45), which is structurally rather close to the oral hypoglycemic agents, is an antianginal agent

instead. Its synthesis involves ester-amide displacement of carbamate 43 with S-2-aminophenylpropane (44) to give 45.



Several obvious variants exist. Tolbutamide, the prototypic drug, has some antiarrhythmic activity by an unknown mechanism. This side effect has become the principal action with tosifen, which itself does not in turn significantly lower blood sugar.¹⁹

REFERENCES

1. R. N. Boys, B. R. Duce, E. R. Smith, and E. W. Byrnes, German Offen. DE2,235,745 (1973); Chem. Abstr., 78, 140411v (1973).
2. H. C. Ferguson and W. D. Kendrick, J. Med. Chem., 16, 1015 (1973).
3. G. H. Douglas, J. Diamond, W. L. Studt, G. N. Mir, R. L. Alioto, K. Anyang, B. J. Burns, J. Cias, P. R. Darkes, S. A. Dodson, S. O'Connor, N. J. Santora, C. T. Tsuei, J. J. Zulipsky, and H. K. Zimmerman, Arzneim. Forsch., 28, 1435 (1978).
4. C. R. Rasmussen, J. F. Gardocki, J. N. Plampin, J. N. Twardzik, B. E. Reynolds, A. J. Molinari, N. Schwartz, W. W. Bennetts, B. E. Price, and J. Marakowski, J. Med. Chem., 21, 10 (1978).
5. G. D. Diana, French Patent FR2,003,438 (1969); Chem. Abstr., 72, 78, 7352 (1970).

6. H. Koelling, H. Thomas, A. Widdig, and H. Wollwever, German Offen. DE2,423,679 (1975); Chem. Abstr., 84, 73949k (1976).
7. J. W. Baker, G. L. Bachman, I. Schumacher, D. P. Roman, and A. L. Tharp, J. Med. Chem., 10, 93 (1967).
8. C. R. Rasmussen, German Offen. DE2,711,757 (1977); Chem. Abstr., 88, 37603s (1978).
9. J. B. Wright, C. M. Hall, and H. G. Johnson, J. Med. Chem., 21, 930 (1978).
10. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 2, Wiley, New York, 1980, p. 87.
11. W. Merkel, J. Med. Chem., 11, 399 (1976).
12. A. Popelak, A. Lerch, K. Stach, E. Roesch, and K. Hardebeck, German Offen. DE1,815,922 (1970); Chem. Abstr., 73, 45519z (1970).
13. E. H. Banitt, W. R. Bronn, W. E. Coyne, and J. R. Schmid, J. Med. Chem., 20, 821 (1977).
14. J. D. Albright, S. A. Schaffer, and R. G. Shepherd, J. Pharm. Sci., 68, 936 (1979).
15. P. L. DeBenneville and N. H. Greenberger, German Offen. DE2,156,835 (1972); Chem. Abstr., 77, 114888r (1972).
16. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 1, Wiley, New York, 1977, p. 136.
17. C. Rufer, J. Med. Chem., 17, 708 (1974).
18. K. Gutsche, E. Schroeder, C. Rufer, O. Loge, and F. Bahlmann, Arzneim. Forsch., 24, 1028 (1974).
19. L. Zitowitz, L. A. Walter, and A. J. Wohl, German Offen. DE2,042,230 (1971); Chem. Abstr., 75, 5532h (1971).


5 Polycyclic Aromatic Compounds

It will have been noted that important structural moieties are sometimes associated with characteristic biological responses (prostanoids, phenylethanolamines, for example). Just as often, however, such structural features show commonality only in the mind of the organic chemist. As will be readily evident from the very diverse biological activities displayed by drugs built on polycyclic aromatic nuclei, this classification is chemical rather than pharmacological. The nucleus does, however, sometimes contribute to activity by providing a means by which pharmacophoric groups can be located in their required spatial orientation; sometimes too, particularly in the case of the monofunctional compounds, the polycyclic aromatic moiety probably contributes to the partition coefficient so as to lead to efficient transport of the drug to the site of action.

1. INDANONES



A rather simple derivative of 1-indanone itself has been reported to possess analgesic activity. This is particularly noteworthy in that this agent, drindene (3),

departs markedly from the structural pattern of either centrally acting or peripheral analgesics. Condensation of 1-indanone (1) with ethyl chloroformate in the presence of alkoxide gives the corresponding hydroxymethylene derivative 2. Reaction with ammonium acetate leads to the corresponding enamine 3, probably by addition of ammonium ion to the terminus of the enone followed by elimination of hydroxide.¹




The discovery of disodium cromoglycate (4) afforded for the first time an agent that was active against allergies by opposing one of the very first events in the allergic reaction; that is, the release of the various substances (mediators) that cause the characteristic symptomology of an allergic attack. The fact that this agent is active only by the inhalation route led to an extensive search for a compound that would show the same activity when administered orally. The various candidates have as a rule been built around some flat polycyclic nucleus and have contained an acidic proton (carboxylate, tetrazole, etc.). One of the simplest of these is built on an indane nucleus. Base catalyzed condensation of phthalic ester 5 with ethyl acetate affords indanedione 6 (shown in the enol form). Nitration by means of fuming nitric acid leads the mediator

release inhibitor nivimedone (7).² The triply activated proton shows acidity in the range of carboxylic acids.




Further investigation on the chemistry of the very potent diuretic drug ethacrinic acid (8) led to a compound that retained the high potency of the parent with reduced propensity for causing side effects, such as loss of body potassium and retention of uric acid. Friedel-Crafts acylation of dichloroanisole 9 with phenylacetyl chloride gives ketone 10. This is then reacted in a variant of the Mannich reaction which involves the aminal from dimethyl-




amine and formaldehyde. The reaction may be rationalized as leading initially to the adduct 11; loss of dimethylamine leads to the enone 12. Cyclization by means of sulfuric acid affords the indanone (13). This last is in turn alkylated on carbon (14) and O-demethylated under acidic conditions. The phenol (15) thus obtained is then alkylated on oxygen by means of ethyl bromoacetate. Saponification of the ester affords indacrinone (17)³.

2. NAPHTHALENES

As noted earlier, most classical antidepressant agents consist of propylamine derivatives of tricyclic aromatic compounds. The antidepressant molecule tametriline⁴ is thus notable in that it is built on a bicyclic nucleus that directly carries the amine substituent. Reaction of 4-phenyl-1-tetralone (18) (obtainable by Friedel-Crafts cyclization of 4,4-diphenylbutyric acid) with methylamine in the presence of titanium chloride gives the corresponding Schiff base. Reduction by means of sodium borohydride affords the secondary amine as a mixture of cis (21) and trans (20) isomers. The latter is separated to afford the more active antidepressant of the pair, tametriline (20).




Topical fungal infections usually involve the lipid-like dermal and subdermal tissues. Drugs with increased lipophilicity would thus be expected to show enhanced antifungal activity by reason of preferential distribution to the lipid-rich site of action. A modification of the antifungal agent tolnaftate (29), which increases its lipophilicity, affords tolciclate (28). One approach to construction of the required bridged tetrahydronaphthol (25) involves Diels-Alder condensation of a benzyne. Thus reaction of dihalo anisole 22 with magnesium in the presence of cyclopentadiene leads directly to the adduct 24. It is likely that 22 initially forms a Grignard-like reagent at the iodo group; this then collapses to magnesium halide and benzyne 23; 1,4 addition to cyclopentadiene leads to the observed product. Preparation of the requisite phenol 26 is completed by catalytic hydrogenation (25) followed by O-demethylation⁵. Reaction of the sodium salt of the phenol with thiophosgene leads to intermediate 27; condensation of N-methyl-*m*-toluidine gives tolciclate (28).⁶




Research carried on in several laboratories in the mid-1960s indicated that triarylethylenes that carry an ethoxyethylamine substituent on one of the rings show very promising antifertility activity. It was quickly found that such agents owe their activity in the particular test system used to their ability to antagonize the effects of endogenous estrogens. One of the more potent agents synthesized in this period was nafoxidine (30). This agent's antifertility activity turned out to be restricted to rodents due to a peculiarity of the reproductive endocrinology of this species. Further clinical testing of compounds in this class revealed that certain estrogen antagonists were remarkably effective in the treatment of breast tumors, particularly those that can be demonstrated to be estrogen dependent. One such agent, tamoxifen, is currently used clinically for that indication.

More recent work in this series demonstrated that a carbonyl group can be interposed between the side-chain-carrying aromatic ring and the ethylene function with full retention of activity. Claisen condensation of benzoate 31 with 2-tetralone affords the β -diketone 32. Reaction of this with p-anisylmagnesium bromide interestingly proceeds preferentially at the ring carbonyl atom. (The thermodynamically favored enol carries the carbonyl at that position.) Spontaneous dehydration leads to the enone 33. The methoxy group on the ring substituted by the carbonyl group is rendered more reactive by the ketone at the para position thus demethylation with an equivalent of sodium ethylthiol leads to phenol 34. Alkylation with 2-chloroethylpyrrolidine affords the antiestrogenic agent trioxifene (35).⁷




Yet another compound that exhibits antidepressant properties, that does not fit the classical mold, is a rather simple substituted amidine. Reaction of amide 36 with triethyloxonium fluoroborate (Meerwein reagent) affords the corresponding imino ether 37. Exposure of this intermediate to methylamine leads to napactidine (38).⁸




Antifungal activity has been described for an equally straightforward derivative of 1-naphthylmethylamine. This

agent, naftidine (40) is obtained by alkylation of amine 39 with cinnamyl bromide.⁹



Excessive activity of the enzyme aldose reductase sometimes accompanies diabetes. The net result is often accumulation of reduced sugars such as galactose in the lens of the eye and ensuing cataract formation. Alrestatin (43), an aldose reductase inhibitor, is one of the first agents found that holds promise of preventing diabetes-induced cataracts. The compound, actually used as its sodium salt, is prepared in straightforward manner by imide formation between 1,8-naphthalic anhydride (41) and glycine.¹⁰




3. TRICYCLIC COMPOUNDS: ANTHRACENE, PHENANTHRENE AND DIBENZOCYCLOHEPTENE

The discovery of the activity of the phenothiazines such as chlorpromazine (44) against schizophrenia pointed the way to drug therapy of diseases of the mind. The intensive


research on the chemistry and pharmacology of those heterocycles is detailed at some length in the first Volume of this series. Those earlier investigations, it should be noted, centered mainly on modification of the side chain and substitution of the aromatic ring. Subsequent research revealed that considerable latitude exists as to the structural requirements of the central ring. For example, clomacran (45) shows the same antipsychotic activity as its phenothiazine counterpart (44). It is thus interesting to note that the fully carbocyclic analogue fluotracen (54) also exhibits CNS activity. This particular agent in fact shows a combination of antipsychotic and antidepressant activity. (In connection with the last it may be of some relevance that the compound may be viewed as a ring-contracted analogue of the tricyclic antidepressants.)

Reaction of substituted nitrile 46 with phenylmagnesium bromide gives, after hydrolysis, the benzophenone 47. Reaction of the ketone with the ylide from trimethylsulfoxonium iodide leads to the epoxide 48. Reductive ring opening of the oxirane by means of phosphorus and hydriodic acid completes conversion of the carbonyl to the homologous methyl group (49). Replacement of bromine by a nitrile group is accomplished by treatment of 49 with cuprous cyanide. Reaction of the product with the Grignard reagent from 3-methoxybromopropane affords the imine 51, which now contains all the required carbon atoms. Treatment of this intermediate with hydrobromic acid achieves both Friedel-Crafts-like ring closure and conversion of the terminal methoxy group to a bromide (52). The latter transformation


may proceed either by direct $\text{S}_{\text{N}}2$ displacement of the protonated methoxy group by bromide or by prior cleavage of the ether to an alcohol followed by the more conventional transformation. Displacement of the terminal bromine by dimethylamine completes construction of the side chain (53). Catalytic reduction proceeds in the usual fashion to give the 9,10-dihydro derivative, fluotracen.¹¹ (Though not specifically stated, the method of synthesis would suggest that these groups bear a cis relationship.)




There is much evidence to suggest that one of mankind's dreaded afflictions, cancer, is not one but a loosely related series of diseases. This diversity has acted as a significant bar to the elucidation of the mechanisms underlying the uncontrolled cell division that characterizes tumor growth. Though some progress has been made toward rational design of antitumor agents, a significant portion of the drug discovery process still relies on random screening. It is thus that one of the screens sponsored by the National Cancer Institute (US) discovered the antitumor activity of a deep blue compound which had been originally synthesized as a dye for use in ball point pen ink. Preparation of this compound, ametantrone (58) starts by reaction of leucoquinizarin (55) with diamine 56 to give the bisimine 57. Air oxidation of the intermediate restores the anthraquinone oxidation state. There is thus obtained ametantrone (58)¹². A similar sequence starting with the leuco base of tetrahydroxyanthraquinone 59 affords the very potent antitumor agent mitoxantrone (61).¹³



Large-scale treatment of a host of lower organisms with biocides seems to lead almost inevitably to strains of that organism that become resistant to the effects of that agent. We thus have bacteria that no longer succumb to given antibiotics, and insects that seemingly thrive on formerly lethal insecticides. The evolution of strains of plasmodia resistant to standard antimalarial agents, coupled with the US involvement in Vietnam, a hotbed for malaria, led to a renewed search for novel antimalarial agents. Halofantrine (70) is representative of the latest generation of these compounds. The preparation of this phenanthrene starts with the aldol condensation of substituted benzaldehyde 62 with phenylacetic acid derivative 63 to give the cinnamic acid 64. Chemical reduction of the nitro group leads to aniline 65. This is then cyclized to the phenanthrene by the classical Pschorr synthesis (nitrous acid followed by strong acid). Though many methods have been proposed for direct reduction of carboxylic acids to aldehydes, these have usually been found less than satisfactory in practice. A more satisfactory method of achieving the transformation consists in reducing the acid to the carbinol (67) and then oxidizing that back to the aldehyde (68); the present sequence employs lead tetraacetate for the last step. Reformatski condensation of 68 with N,N-di-(n-butyl)bromoacetamide and zinc affords amidoalcohol 69. This is reduced to the amino alcohol by means of diborane to give halofantrine (70).¹⁴



An analogue of amitriptyline which contains an additional double bond in the central seven membered ring shows much the same activity as the prototype. Treatment of dibenzocycloheptanone 71 with N-bromosuccinimide followed by triethylamine serves to introduce the additional double bond by the bromination-dehydrohalogenation sequence. Reaction of the carbonyl group with the Grignard reagent from 3-chloropropyl-N,N-dimethylamine serves to introduce the side chain (73). Acid catalyzed dehydration affords the antidepressant compound cyclobenzaprine (74).¹⁵



REFERENCES

1. P. D. Hammen and G. M. Milne, German Offen., 2,360,096; Chem. Abstr., **81**, 105093 (1974).
2. D. R. Buckle, N. J. Morgan, J. W. Ross, H. Smith, and B. A. Spicer, J. Med. Chem., **16**, 1334 (1973).
3. S. J. DeSolm, D. W. Woltersdorf, E. J. Cragoe, L. S. Waton and G. M. Fanelli, J. Med. Chem. **21**, 437 (1978).
4. S. Reinhard, J. Org. Chem., **40**, 1216 (1975).
5. H. Tanida, R. Muneyuki and T. Tsuji, Bull. Chem. Soc. Jap., **37**, 40 (1964).
6. P. Melloni, M. Rafaela, V. Vecchietti, W. Logemann, S. Castellino, G. Monti, and I. DeCarneri, Eur. J. Med. Chem., **9**, 26 (1964).
7. C. D. Jones, T. Suarez, E. H. Massey, L. J. Black and F. C. Tinsley, J. Med. Chem. **22**, 962 (1979).
8. J. R. McCarthy, U.S. Patent 3,903,163; Chem. Abstr. **83**, 192933 (1975).


9. B. Daniel, German Offen, 2,716,943; Chem. Abstr. 88, 62215 (1978).
10. S. Kazimir, N. Simard-Duquesne and D. M. Dvornik, U.S. Patent 3,821,383; Chem. Abstr. 81, 176158 (1974).
11. P. N. Craig and C. L. Zirkle, French Patent, 1,523,230; Chem. Abstr. 71, 101609 (1969).
12. R. K. Y. Zeecheng and C. C. Cheng, J. Med. Chem., 21, 291 (1978).
13. K. C. Murdock, R. G. Child, P. F. Fabio, R. B. Angier, R. E. Wallace, F. E. Durr, and R. V. Citavella, J. Med. Chem., 22, 1024 (1979).
14. W. T. Colwell, V. Brown, P. Christie, J. Lange, C. Reece, Y. Yamamoto and D. W. Henry, J. Med. Chem., 15, 771 (1972).
15. S. O. Winthrop, M. A. Davis, G. S. Myers, J. G. Gavin, R. Thomas and R. Barber, J. Org. Chem., 27, 230 (1962).

6 Steroids

The steroid nucleus provides the backbone for both the hormones that regulate sexual function and reproduction and those involved in regulation of mineral and carbohydrate balance. The former comprise the estrogens, androgens, and progestins; cortisone, hydrocortisone, and aldosterone are the more important entities in the second category. Synthetic work in the steroid series, accompanied by some inspired endocrinological probes, led to many signal successes. Research on estrogens and progestins thus led to the oral contraceptives, and corresponding efforts on cortisone and its derivatives culminated in a series of clinically important antiinflammatory agents.

Few if any endogenous hormones seldom exert a single action. These compounds typically elicit a series of re-


sponses on different biological end points and organ systems. It should thus not be surprising that both natural and modified steroids also show more than one activity; these ancillary activities, however, often consist of undesirable actions, and are thus considered side effects. Volumes 1 and 2 of this series detail the enormous amount of work devoted to the steroids inspired at least partly by the goal to separate the desired activity from those side effects. When it became apparent that this goal might not be achievable, there was a considerable diminution in the synthetic work in the steroid series; this is well illustrated by comparison of this section with its counterparts in the preceding volumes.



1. ESTRANES

The adventitious discovery of the antitumor action of the nitrogen mustard poison war gases led to intensive investigation of the mode of action of these compounds. In brief, it has been fairly well established that these agents owe their effect to the presence of the highly reactive bis(2-chloroethyl)amine group. The cytotoxic activity of

these drugs is directly related to the ability of this group to form an irreversible covalent bond with the genetic material of cells, that is, with DNA. Since this alkylated material can then no longer perform its function, replication of the cell is disrupted. The slight selectivity shown for malignant cells by the clinically used alkylating agents depends largely on the fact that these divide more rapidly than those of normal tissue. There have been many attempts to achieve better tissue selectivity by any number of other stratagems. One of these involves linking the mustard function to a molecule that itself shows very specific tissue distribution. Steroids are prime candidates as such carriers since they are well known to exhibit highly selective organ distribution and tissue binding. Estramustine (4) and prednimustine (64; see Section 3 below) represent two such site directed cytotoxic agents.




Reaction of bis(2-chloroethyl)amine with phosgene affords the corresponding carbamoyl chloride (2). Acylation of estradiol (3) with this reagent leads to estramustine (4)¹. Though reaction with the more nucleophilic alcohol function at 17 might at first sight lead to a competing reaction, the highly hindered nature of this alcohol greatly reduces its reactivity.

Oral contraceptives almost invariably consist of a mixture of a progestational agent active by the oral route and a small amount of an estrogen. It was discovered quite early that, in contrast to the natural compounds, those lacking the 19 methyl group (19-nor) showed good oral activity; the availability of practical methods for total synthesis led to some emphasis on agents that possess an ethyl group at position 13 rather than the methyl of the natural steroids. Very widespread use of oral contraceptives led to the recognition of some side effects that were associated with the progestin component; there has thus been a trend to design drugs that could be administered in smaller quantity and a corresponding search for ever more potent progestins.


Birch reduction of the norgetrel intermediate 5² followed by hydrolysis of the enol ether gives the enone 6; oxidation of the alcohol at 17 leads to dione 7. Fermentation of that intermediate in the presence of the mold Penicillium raistricky serves to introduce a hydroxyl group at the 15 position (8). Acetal formation with neopentyl glycol affords the protected ketone which consists of a mixture of the Δ^5 and $\Delta^{5,10}$ isomers (9); hindrance at position 17 ensures selective reaction of the 3 ketone. The


hydroxyl is then converted to its mesylate (10). Treatment with sodium acetate leads to elimination of the mesylate and thus formation of the corresponding enone (11). Reaction of the ketone at 17 with ethynylmagnesium bromide introduces the requisite side chain. Removal of the ketal group by means of aqueous oxalic acid completes the synthesis of gestodene (13)³.



A rather more complex scheme is required for preparation of the analogue gestrinone (27) which contains unsaturation in rings A, B, and C. The key intermediate 24 can be obtained by Robinson annulation on dione 14 with enone 15 to give the bicyclic intermediate 16. Successive

reduction of the double bond and the cyclopentyl carbonyl group followed by esterification of the thus obtained alcohol gives ketoester 17. This last can then be converted to the enol lactone 18 by successive partial saponification and treatment with acetic anhydride. Condensation with the Grignard reagent from bromoketal 19 gives after hydrolysis the tricyclic intermediate 20. (This reaction can be rationalized as initial reaction of the organometallic with the lactone carbonyl; the diketone formed by hydrolysis would then cyclize under the reaction conditions.) Treatment of 20 with pyrrolidine serves to close the last ring via its enamine (21). Hydrolysis of the first formed 3-enamine leads to the doubly unsaturated steroid 22. Treatment with dicyanodichloroquinone (DDQ) would serve to introduce the last double bond with consequent formation of the conjugated 5,9,10 triene (23); hydrolysis of the ester at 17 and subsequent oxidation of the alcohol would give the key 3,17 diketone 24. Treatment with ethylene glycol in the presence of acid leads to formation of the 3 ketal. Reaction with ethynylmagnesium bromide (26) followed by removal of the ketal group gives gestrinone (27)^{4,5}.







2. ANDROSTANES


Despite some early hopes, the drugs related to the androgens have found rather limited use. This has in practice been confined to replacement therapy in those cases where endogenous hormone production is deficient; some agents that show reduced hormonal effects have found some application as anabolic agents as a consequence of their ability to rectify conditions that lead to loss of tissue nitrogen. In analogy with the estrogen antagonists, an antiandrogen would seem to offer an attractive therapeutic target for treatment of diseases characterized by excess androgen stimulation (e.g., prostatic hypertrophy) and androgen dependent tumors. Attempts to design specific antagonists to androgens have met with limited success, however.

Halogenation of steroid 3-ketones can lead to complicated mixtures by virtue of the fact that the kinetic enol leads to 3 halo products, whereas the thermodynamic product is that halogenated at the 4 position. Carefully controlled reaction of the 5α -androstanolone 28 with chlorine thus leads to the 2α -chloro derivative (29). Reaction of that intermediate with *O*(*p*-nitrophenyl)-hydroxylamine affords the androgenic agent nistremine acetate (30)⁶.




Replacement of the hydrogen at the 17 position of the prototypical androgen testosterone (31) by a propyl group interestingly affords a compound described as a topical antiandrogen. Reaction of the tetrahydropyranyl ether of dehydroepiandrosterone (32) with propylmagnesium bromide gives after removal of the protecting group the corresponding 17α -propyl derivative 33. Oppenauer oxidation of the 3-hydroxy- Δ^5 function leads to the corresponding conjugated ketone. There is thus obtained topterone (34)⁷.







A highly modified methyltestosterone derivative also exhibits antiandrogenic activity. One synthesis of this compound involves initial alkylation of methyltestosterone (35) by means of strong base and methyl iodide to afford the 4,4-dimethyl derivative 36. Formylation with alkoxide and methyl formate leads to the 2-hydroxymethyl derivative 37. Reaction of this last with hydroxylamine leads to formation of an isoxazole ring. There is then obtained azastene (38)⁸.



3. PREGNANES

With very few exceptions, the biological activities of synthetic steroids tend to parallel those of the naturally occurring hormones on which they are patterned. Compounds with distant pharmacological activity are, as a rule, quite rare. It is thus intriguing that inclusion of a tertiary amine at the 11 position of a pregnane leads to a compound with activity far removed from its close analogues. The agent in question, minaxalone (47), exhibits anesthetic activity. Epoxidation of progesterone derivative 40, obtainable in several steps from 11-ketoprogesterone (39)⁹, gives the corresponding α -epoxide 41. Reaction of that compound with alkoxide leads to diaxial opening of the oxirane and consequent formation of the 2 β -ethoxy 3 α -hydroxy derivative 42. Reaction with ethylene glycol leads cleanly to selective formation of the 17 ketal (43) by reason of the highly hindered environment about the 11 carbonyl. For the same reason, formation of the oxime 44 requires forcing conditions. Chemical reduction of that oxime leads to the thermodynamically favored equatorial α -amine 45. (Catalytic reduction would have given the β -amine.) Methylation of the amine by means of formic acid and formaldehyde leads to the corresponding dimethylamino derivative (46). Removal of the ketal group completes the synthesis of minaxalone (47)¹⁰.








Spironolactone (48) has proved a very useful diuretic and antihypertensive agent. This drug, that owes its effect to antagonism of the endogenous steroid hormone that regulates mineral balance, aldosterone, exhibits in addition some degree of progestational and antiandrogenic activity. Further analogues have thus been prepared in an effort to prepare an agent free of those side effects.


Preparation of the newest of these, spirorenone (61)¹¹, starts by 7-hydroxylation of dehydroepiandrosterone derivative 49. Though this transformation has also been accomplished by chemical means, microbiological oxidation by Botryodiploda malorum apparently proves superior. Acylation with pivalic anhydride proceeds selectively at the 3 hydroxyl group (51). Epoxidation by means of tertiary butylhydroperoxide and vanadium acetylacetonate affords exclusively the β epoxide (52). The remaining hydroxyl is

then displaced by chlorine by means of triphenylphosphine and carbon tetrachloride (53). Sequential reductive elimination (54) followed by saponification gives the allylic alcohol 55. Reaction with the Simmons-Smith reagent affords the corresponding cyclopropane, 56, the stereochemistry being determined by the adjacent hydroxyl group. Addition of the dianion from propargyl alcohol to the carbonyl group at position 17 adds the required carbon atoms for the future lactone (57). The side chain is then reduced by catalytic hydrogenation (58). Oxidation of this last intermediate by means of pyridinium chlorochromate simultaneously oxidizes the primary alcohol to an acid and the secondary alcohol at position 3 to a hydroxyketone; under the reaction conditions, the latter eliminates to give an enone while the hydroxy acid lactonizes. There is thus obtained directly the intermediate 60. Dehydrogenation by means of DDQ introduces the remaining double bond to afford spirorenone (61)¹¹.






As noted above, the steroid nucleus has been a favorite for the design for site directed alkylating antitumor drugs. Thus reaction of prednisolone (62) with anhydride 63 affords the 21 acylated derivative, prednimustine (64)¹².




A preponderance of the work devoted to steroids, as judged from the number of compounds bearing generic names, has clearly been that in the area of corticosteroids related

to cortisone. Much of the effort, particularly that detailed in the earlier volumes was no doubt prompted by the very large market for these drugs as antiinflammatory agents. Heavy usage led to the realization that parenteral use of these agents carried the potential for very serious mechanism related side effects. There has thus been a considerable recent effort to develop topical forms of these drugs for local application to rashes, irritation and other surface inflammations. A good bit of the work detailed below is aimed at improving dermal drug penetration.


Because skin exhibits many of the properties of a lipid membrane, dermal penetration can often be enhanced by increasing a molecule's lipophilicity. Preparation of an ester of an alcohol is often used for this purpose since this stratagem simultaneously time covers a hydrophilic group and provides a hydrophobic moiety; the ready cleavage of this function by the ubiquitous esterase enzymes assures availability of the parent drug molecule. Thus acylation of the primary alcohol in flucinolone (65)¹³ with propionyl chloride affords procinonide (66)¹⁴; the same transform employing cyclopropyl carbonyl chloride gives ciprocinonide (67)¹⁴.






Further oxidation of the carbon at position 21 is interestingly consistent with antiinflammatory activity. Thus oxidation of flucinolone with copper (II) acetate affords initially the hydrated ketoaldehyde 68; exchange with methanol gives the 21 dimethyl acetal (69), flumoxonide.¹⁵

Omission of the fluoro substituents at the 6 and 9 positions leads to the antiinflammatory compound agent budesonide (71). This compound is obtained by formation of the acetal of the 16,17 diol 70¹⁶ with butyraldehyde.¹⁷




It is a hallmark of the structure activity relationships of the corticoids that the effects of structural modifications that lead to increased potency are usually additive. The fact that more than half a dozen such modifications each lead to increased potency opens ever new possibilities for combinations and permutations. Meclorison dibutyrate (74) thus combines the known potentiating effects of the replacement of the 11-hydroxyl by chlorine as well as incorporation of a 16 α -methyl

group. Addition of chlorine to the 9,11 double bond of 72 would afford the corresponding dichloro derivative, 73. (Addition of halogen is initiated by formation of the chloronium ion on the less hindered alpha face; diaxial opening of the intermediate leads to the observed regio- and stereochemistry.) Acetylation of that intermediate with butyric acid in the presence of trifluoroacetic acid leads to meclorisone dibutyrate (74).




Halogenation of the 7 position also proves compatible with good antiinflammatory activity. Construction of this compound, aclomethasone dipropionate (80), starts by introduction of the required unsaturation at the 6,7 position by dehydrogenation with DDQ (76). The highly hindered nature of the hydroxyl at position 17 requires that a roundabout scheme be used for formation of the corresponding ester. Thus treatment of 76 with ethyl orthoformate affords first the cyclic orthoformate 77. This then rearranges to the 17 ester 78 on exposure to acetic acid. Acylation of the 21 alcohol is accomplished in straightforward fashion with

propionic anhydride (79). Addition of hydrogen chloride completes the synthesis of aclomethasone dipropionate (80).²⁰ (This last reaction may in fact involve 1,6 conjugate addition of the reagent; this would then ketonize to the observed product under the acidic reaction conditions).



Incorporation of a vinylic bromide at the 2 position also gives a compound with good activity. Bromination of fluorohydrin 81²¹ under carefully controlled conditions gives the 2 α bromo derivative 82. The hydroxyl at the 11 position is then converted to its mesylate with methanesulfonyl chloride (83). Reaction of that intermediate with acetic anhydride under forcing conditions (perchloric acid) gives the 5,17,21 triacetate 84. Treatment with sodium acetate leads to elimination of the acetate at 5 and formation of the enone 85. The presence of that function



eliminates possible future complication due to the known facile rearrangement of halogen from the 2 to the 4 position. Thus exposure of the intermediate 85 to bromine gives the 2,2 dibromo derivative 86. Elimination of one of those halogens by means of lithium carbonate and lithium bromide leads to the vinylic bromide 87; these basic conditions achieve simultaneous elimination of the 11 mesylate and formation of the 9,11 olefin. That last


function is then used for introduction of the 9α -fluoro-11- β -alcohol by the standard scheme. Thus exposure of the olefin to HOBr (aqueous NBS) gives bromohydrin 88 (diaxial opening of the initially formed α -bromonium intermediate). Treatment with base gives the β -epoxide 89. Opening of the oxirane with hydrogen fluoride leads finally to the antiinflammatory agent haloprednone (90).²²

4.MISCELLANEOUS STEROIDS


The cardiostimulant action of extracts of the leaves of the foxglove plant (Digitalis purpurea) were recognized as early as the eighteenth century. Careful examination of these extracts led to the isolation of a series of so-called cardiac glycosides which consist of hydroxylated steroids generally substituted by an unsaturated five-membered lactone at the 17 position and glycosidated by a series of sugars at the 3 position. Though these drugs have proved extremely useful in the treatment of diseases marked by failure of the heart muscle, they are at the same time extremely toxic. Only a very narrow margin exists between effective and toxic doses. (It is of interest that the need to carefully adjust blood levels of digitalis contributed to the birth of the science of pharmacokinetics.) Though there is a great need for a digitalis-like drug with a greater margin of safety, there has been surprisingly little synthetic effort devoted to this area.

It is known that the presence of the 14β -hydroxyl group and a sugar at the 3 position are absolute requirements for activity. A modified drug actodigin (100) demonstrates that reversal of the lactone 17 is consistent with activity. Reduction of digitoxigenin 91, the aglycone of digitoxin



(92) with diisobutylaluminum hydride leads to the diol 93. Oxidation of that intermediate with triethylamine-sulfur trioxide complex leads to the furan 95. (This transformation can be rationalized by invoking the intermediacy of an unsaturated hydroxyaldehyde (94), followed by formation of internal acetal 94.) The hydroxyl at the 3 position is then protected as its chloroacetate (96). Reaction of the furan ring with NBS followed by hydrolysis of the halide leads to the furanone ring at 17 which is in effect the reversed lactone from 91. The protecting group is then removed (98), and the alcohol glycosidated with the acetylated halo sugar 99. Removal of the acetate groups by saponification affords actodigin (101).²³



One of the triumphs of the science of nutrition is the careful investigation that linked childhood rickets with vitamin D deficiency. This work, which led to methods for treating the disease, is too familiar to need repetition. A direct consequence of these efforts was the elucidation of the pivotal role played by vitamin D in calcium metabolism, as well as the structural studies that revealed that this compound (102) is in fact a steroid derivative. The past several decades have seen the development of physical and spectroscopic methods which allow the study of ever smaller quantities of organic compounds. As a direct outgrowth of this, it has become possible to carry out very detailed studies on the metabolism of endogenous and exogenous organic compounds. Applications of such methods to vitamin D revealed that this agent in fact undergoes further hydroxylation in the body. The very high biological activity of the resulting compounds soon cast doubt on the question whether the vitamin (102) was in fact the ultimate biologically active agent. Detailed work revealed that this metabolism is in fact required for calcium regulatory action. It has been demonstrated that the mono and dihydroxy metabolites act more quickly than vitamin D and that they show much higher potency as antirachitic agents. This finding has some very practical significance since a number of disease states such as kidney failure, which are marked by calcium loss from bone, are associated with deficient vitamin D hydroxylation. The two hydroxylated metabolites, calcifediol (113) and calcitriol (145), have been introduced for clinical treatment of diseases associated with impaired vitamin D metabolism.




The reported synthesis of the monohydroxy metabolite (113) starts with acid 105 obtained as a by-product from oxidative cleavage of the side chain of cholesterol. This is transformed to the corresponding diazomethyl ketone 107 by reaction of the acetylated acid chloride 106 with diazomethane. Arndt-Eistert rearrangement of that intermediate affords the homologated ester 108. Allylic bromination (109) by means of dibromodimethylhydantoin followed by dehydrohalogenation with trimethyl phosphite establishes the cis diene functionality (110) required for opening of ring B. Reaction of the ester with methyl Grignard reagent completes construction of the side chain (111). Photolysis of that diene affords the product of electrocyclic ring opening (112). It now remains to isomerize the triene function. This is accomplished by thermal equilibration. There is thus obtained calcifidiol (113).²⁴ The low yields reported leave it open to question that this is the commercial route.




A rather complex stereospecific convergent total synthesis has been reported for calcitriol (145). Construction of the A ring fragment starts with epoxidation of chiral d-carvone (114) to afford epoxide 115. Condensation with the ylide from diethyl (carboxyethyl)-phosphonate gives the corresponding ester largely as the E isomer. The epoxide is then opened with acetate (117) and acetylated to give diacetate 118. The methylene group on the side chain is then cleaved to the ketone by means of osmium tetroxide periodate reagent. Bayer-Villiger cleavage of the resulting methyl ketone (trifluoroperacetic acid) affords finally triacetate 120. This is then hydrolyzed (121) and converted to the bis trisilyl derivative 122. Dehydration under very specialized conditions gives the exo-

methylene derivative 123. The conjugated double bond is then isomerized by irradiation in the presence of a sensitizer (124). The carbethoxy group is then reduced to an alcohol (125) and this converted to the corresponding allylic chloride (126). This reactive function is displaced with lithium diphenylphosphide (127). Oxidation of phosphorus affords the A ring intermediate 128 functionalized so as to provide an ylide.



The CD fragment is synthesized starting with resolved bicyclic acid 129. Sequential catalytic hydrogenation and reduction with sodium borohydride leads to the reduced hydroxy acid 130. The carboxylic acid function is then converted to the methyl ketone by treatment with methyl-lithium and the alcohol is converted to the mesylate. Elimination of the latter group with base leads to the conjugated olefin 133. Catalytic reduction followed by equilibration of the ketone in base leads to the saturated methyl ketone 134. Treatment of that intermediate with peracid leads to scission of the ketone by Bayer Villiger reaction to afford acetate 135. The t-butyl protecting group on the alcohol on the five membered ring is then removed selectively by means of trimethylsilyl iodide (136). Oxidation by means of pyridinium chlorochromate gives the ketone 137. That function is then reacted with ethylidene phosphorane to afford the olefin 138. Ene reaction with ethyl propiolate proceeds stereo- and regioselectively to the extended side chain of 139; note particularly that the chiral center at C₂₀ has been introduced in the correct absolute configuration. Catalytic reduction leads to the saturated intermediate 140. The ester function is then reduced to the aldehyde by means of diisobutylaluminum hydride and the acetate saponified to afford 141. Condensation of the aldehyde on 141 with isopropyl phosphorane adds the last required carbon atoms (142). The tertiary hydroxyl group at the future 25 position is then introduced by means of an oxymercuration reaction (143). Oxidation of the secondary hydroxyl group completes the synthesis of the CD moiety, 144.

Condensation of the ylide from 144 with the A ring fragment (as its TMS derivative) gives, after removal of the protecting groups, the vitamin D metabolite cacitriol (145).²⁵



REFERENCES


1. K. B. Hogberg, H. J. Fex, I. Konyves, and H. O. J. Kneip., German Patent 1,249,862 Chem. Abstr. 68, 3118j (1968).
2. See D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol.1, Wiley, New York, 1977, p. 168.
3. H. Hofmeister, R. Wiechert, K. Annen, H. Laurent and H. Steinbeck, German Offen. 2,546,062; Chem. Abstr. 87, 168265k (1977).
4. L. Velluz, G. Nomine, R. Bucort, and J. Mathieu, Compt. Rend., 257, 569 (1963).
5. D. Bertin and A. Pierdet, French Patent 1,503,984; Chem. Abstr. 70, 4391w (1969).
6. A. Hirsch, German Offen. 2,327,509; Chem. Abstr. 80, 83401g (1974).
7. A. L. Beyler and R. A. Ferrari, Ger. Offen., 2,633,925; Chem. Abstr., 86 161310s (1977).
8. G. O. Potts, U.S. Patent 3,966,926; Chem. Abstr. 85, 83244 (1976); note this is a use patent.
9. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 1, Wiley, New York, 1977, p. 191.
10. G. H. Phillips and G. Ewan, German Offen. 2,715,078; Chem. Abstr. 88, 38078m (1978).
11. D. Bittler, H. Hofmeister, H. Laurent, K. Nickiseh, R. Nickolson, K. Petzold and R. Wiechert, Angew. Chem. Int. Ed. Engl. 21, 696 (1982).
12. H. J. Fex, K. B. Hogberg and I. Konyves, German Offen. 2,001,305; Chem. Abstr. 73, 99119n (1970).
13. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 1, Wiley, New York, 1977, p. 202.

14. B. J. Poulson, U.S. Patent 3,934,013; Chem. Abstr. 84, 111685f (1976).
15. M. Marx and D. J. Kertesz, German Offen. 2,630,270; Chem. Abstr., 86, 140,345s (1977).
16. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 2, Wiley, New York, 1980, p. 180.
17. A. Thalen and R. Brattsand, Arzneim Forsch. 29, 1607 (1979).
18. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 1, Wiley, New York, 1977, p. 198.
19. E. Shapiro, et.al. Steroids 9, 143 (1967).
20. M. J. Green, H. J. Shue, E. L. Shapiro and M. A. Gentles, U.S. Patent 4,076,708; Chem. Abstr. 89, 110119r (1978).
21. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Vol. 1, Wiley, New York, 1977 p. 195.
22. M. Riva and L. Toscano; German Offen. 2,508,136; Chem. Abstr. 84, 31311 (1976).
23. J. M. Ferland, Can. J. Chem. 52, 1652 (1974).
24. J. A. Campbell, D. M. Squires, and J. C. Babcock, Steroids 13, 567 (1969).
25. E. G. Baggiolini, J. A. Iacobelli, B. M. Hennessy, and M. V. Uskokovic, J. Am. Chem. Soc., 104, 2945 (1982).

7 Compounds Related to Morphine

Pain is probably the immediate stimulus for more visits to the physician's office than all other complaints combined. Since pain serves as an alert to injury, it is often the first harbinger of disease; pain is thus associated with a multitude of physical ills. The fact that this sensation often persists well beyond the point where it has served its alerting function makes its alleviation a prime therapeutic target. In a very general way, the sensation of pain can be divided into two segments. The first is the immediate stimulus that sets off the chain of events; this could be a surface injury such as a burn or a cut, an inflamed internal organ, or any other disorder that causes pain receptors to be triggered. Following a rather complex series of neurochemical transmissions, the signal reaches the brain, where it is processed and finally

presented as the sensation of pain. Treatment of pain follows a roughly similar duality. The pain receptors, for example, can be blocked by local anesthetics in those few cases where the insult is localized in the periphery of the body. (In practice, this is restricted to minor surgery and dentistry.) It has recently become recognized that the pain that accompanies inflammation and related conditions is actually triggered by the local synthesis of high levels of prostaglandins; compounds that inhibit this reaction will, in fact, attenuate the pain attendant to the causative inflammation. Cyclooxygenase inhibitors such as aspirin and other nonsteroid antiinflammatory agents have thus found a secure place in the treatment of the mild to moderate pain associated with elevated prostaglandin synthesis. There remains, however, a large category of pain that is not affected by interference at the receptor stage; intervention is achieved in this case at the level of the central nervous system; hence the sobriquet, central analgetics. Rather than blocking or in some way interfering with the pain signal, agents in this class change the perception of the signal. The opium alkaloid morphine (1) is the prototype central analgetic. The fact that its analgetic properties were discovered centuries ago make it clear that in this case theory came a good bit later than practice.




Though morphine is an extremely effective analgesic, it has an associated series of side effects that limit its legitimate use. The most prominent among these is, of course, its propensity to cause physical addiction. A significant amount of work has thus been devoted to the synthesis of analogues with a view to modifying the pharmacological spectrum and, in particular, avoiding addiction potential. As will be noted from the following discussion (and that in the earlier volumes), this work has led to structures that have little in common with the prototype molecule.

1. BRIDGED POLYCYCLIC COMPOUNDS

It has been found empirically that central analgesics that possess some degree of activity as antagonists of the effects of morphine tend to show a reduced propensity for causing physical addiction. Again empirically, it was noted that this could often be achieved by replacement of the N-methyl group by allyl, cyclopropylmethyl, or cyclobutylmethyl; additional nuclear modifications often contributed to this activity.


Exposure of the opium alkaloid thebaine (2) to mild acid leads to hydrolysis of the enol ether function followed by migration of the double bond to yield the conjugated enone 3. Addition of lithium diethylcuprate proceeds by 1,4 addition from the less hindered side to give the intermediate 4. Treatment of that with cyanogen bromide under von Braun conditions leads to the isolable aminocyanide (5). This is then converted to the secondary amine (6) by treatment with aqueous base. Alkylation of

that intermediate with cyclopropylmethyl chloride affords the analgesic codorphone (7).¹




The development of schemes for the total synthesis of the carbon skeleton of morphine revealed that the fused furan ring was not necessary for biological activity. More recently it has been found that substitution of a pyran ring for the terminal alicyclic is also consistent with biological activity. Starting material for this preparation is ketoester 8, available by one of the classical benzomorphan syntheses.² Condensation with the ylide from diethyl(carbethoxyethyl)phosphonate affords diester 9. (The course of the reaction is probably helped by the fact that the β -ketoester can not undergo tautomerism to its enol form.) Catalytic reduction proceeds from the less hindered face to afford the corresponding saturated diester (10). Reduction of the carbonyl function by means of lithium aluminum hydride gives the glycol 11; this undergoes internal ether formation on treatment with acid to form the pyran ring of 12. Treatment with cyanogen bromide (or ethyl chloroformate) followed by saponification of the

intermediate leads to the secondary amine (14). This is converted to the cyclopropylmethyl derivative 16 by acylation with cyclopropylcarbonyl chloride followed by reduction of the thus formed amide (15) with lithium aluminum hydride. Cleavage of the O-methyl ether with sodium ethanethiol affords proxorphan (17).




Replacement of the alicyclic ring of morphine in addition to omission of the furan ring leads to a thoroughly investigated series of analgesic compounds known as the

benzomorphans. Depending on the substitution pattern, these agents range in activity from potent agonists to antagonists. Reduction of the carbonyl group in oxygenated benzomorphan 18 affords the corresponding alcohol (19). This intermediate is then N-demethylated by means of cyanogen bromide (20). Acylation with cyclopropylcarbonyl chloride gives the amide 21. The alcohol is then converted to the ether 22 by treatment with methyl iodide and base. Treatment with lithium aluminum hydride serves to reduce the amide function (23). Cleavage of the phenolic ether by one of the standard schemes affords moxazocine (24).⁴





A rather unusual reaction forms the key step in the preparation of a benzomorphan bearing a fatty side chain. The scheme used to form the benzomorphan nucleus, which is patterned after the Grewe synthesis originally developed for

preparing morphinans, is fairly general for this class of compounds. Preparation of tonazocine (33) starts with the condensation of the Grignard reagent from 25 with the pyridinium salt 26. (Note that reaction actually occurs at the more sterically hindered center.) In the usual synthetic route, the enamine function would next be reduced and the olefin cyclized. In the case at hand, however, the diene function of 27 is condensed with ethyl acrylate in a Diels-Alder reaction (28). Treatment with strong acid leads to cyclization of the olefin into the aromatic ring and formation of the benzomorphan nucleus (29). Acylation of the anion obtained from that intermediate by means of lithium diisopropyl amide with hexanoyl chloride gives the β -ketoester 30. Heating of that compound with formic acid leads directly to the ring-opened benzomorphan 32. This transform can be rationalized as involving proton mediated cleavage of the blocked β -ketoester to afford 31; decarboxylation under the strongly acidic conditions then leads to the observed product. Cleavage of the phenolic ether affords the analgesic agent tonazocine (33).⁵



2. PIPERIDINES


Still further simplification of the structural requirements for central analgesic activity came from the serendipitous observation that the simple phenylpiperidine, meperidine (34), shows biological activity almost indistinguishable from that of morphine. Further elaboration of that molecule led to one of the most potent opioid analgesics, fentanyl (35).




In-depth investigation of the structure activity relationships in the fentanyl meperidine series in the Janssen laboratories revealed that additional substitution of the amide nitrogen-bearing center resulted in still further enhancement of analgesic potency. Several compounds were obtained as a result of this work, which showed analgesic activity in animal models at doses some five decade orders of magnitude lower than morphine; the biological profile of these agents is, however, almost identical to that of the classical opioids.

Reaction of the carbonyl group of piperidone 36 with cyanide and aniline leads to formation of a cyanohydrin-like function known as an α -aminonitrile (37); hydrolysis under


strongly acidic conditions gives the corresponding amide 38. (Although aminonitriles are somewhat labile, particularly under basic conditions, the corresponding aminoamide is a quite stable function.) The benzyl amine protecting group is then removed by catalytic hydrogenation (39). Alkylation with 2-phenethyl bromide proceeds on the more nucleophilic aliphatic amine to afford 40. Ethanolysis of the amide function leads to the corresponding ester, 41. Acylation of the remaining secondary amine function with propionyl chloride affords carfentanyl (42).⁶ The same sequence starting with the corresponding 3-methylpiperidone (43)⁷ affords lofentanyl (44).




Alkylation of intermediate 39 with 2-(2-bromoethyl)thiophene affords the corresponding thiophene-containing compound 45. Ethanolysis then leads to ethyl ester (46). Reduction of the carbonyl function affords the carbinol 47. Alkylation of the alkoxide obtained from the alcohol with methyl iodide gives the methyl ether 48. Acylation with propionyl chloride leads to the very potent opioid analgesic sulfentanyl (49).⁸ In the absence of a specific reference, one may speculate that alkylation of heterocycle 50 with 1-bromo-2chloroethane would give the chloroethyl derivative 51. Use of this for alkylation of 39 would give the heterocycle substituted intermediate 52. A similar scheme via structures 53, 53a, and 54 would then afford alfentanil (55).






Fusion of an alicyclic ring onto the piperidine so as to form a perhydroisoquinoline is apparently consistent with analgesic activity. Synthesis of this agent, ciprefadol (68), starts with the Michael addition of the anion from cyclohexanone 56 onto acrylonitrile (57). Saponification of the nitrile to the corresponding acid (58) followed by Curtius rearrangement leads to isocyanate 59. Acid hydrolysis of the isocyanate leads directly to the indoline 61, no doubt by way of internal Schiff base formation from the intermediate amine 60. Alkylation by means of trimethyloxonium fluoroborate affords ternary iminium salt 62. Treatment of that reactive carbonyl-like functionality with diazomethane gives the so-called azonia salt 63 (note the analogy to the hypothetical oxirane involved in ring expansion of ketones with diazomethane). Exposure of the aziridinium intermediate to base leads to ring opening and consequent formation of the octahydroisoquinoline (64). Reduction of the enamine (catalytic or borohydride) affords the perhydroisoquinoline 65. This compound is then subjected to one of the N-demethylation sequences and the resulting secondary amine (66) alkylated with cyclopropylmethyl bromide (67). O-Demethylation of the phenol ether completes the preparation of ciprefadol (68).⁹





An internally bridged arylpiperidine in which the aryl group has been moved to the 3 position interestingly retains analgesic activity. It should be noted, however, that this compound has been described as nonnarcotic on the basis of its animal pharmacology. Reaction of the carbene obtained from treatment of bromoaryl acetate 69 with ethyl acrylate affords the cyclopropane 70. The *cis* stereochemistry of the product probably represents the fact that this isomer shows fewer nonbonding interactions than does its *trans* counterpart. Saponification of the ester followed by reaction of the resulting diacid (71) with urea leads to the cyclic imide 72. Reduction of the carbonyl groups is achieved by treatment with sodium aluminumbis[2(methoxy)ethoxy]hydride. There is thus obtained bicifadine (73).¹⁰






3. MISCELLANEOUS COMPOUNDS


The large amount of synthetic work devoted to central analgesic agents led to the elaboration of fairly sound structure activity relationships. Until fairly recently, structural requirements for analgesic activity could be reliably covered by the so-called Beckett and Casey rule. In its most general form, this requires an aromatic ring attached to a quaternary center with basic nitrogen at a distance of about two carbon atoms from that center. (Fentanyl and its analogues were incorporated by assuming that the fully substituted anilide nitrogen is equivalent to the quaternary center.) Some quite potent analgesics that have recently been prepared do not fit this generalization very well, suggesting that it perhaps needs to be revised.

The benzazepines, verilopam (79) and anilopam (81), for example, represent significant departures from the above generalization. Construction of the former starts with the alkylation of veratrylamine (74) with the dimethyl acetal of bromoacetaldehyde to give the secondary amine 75. Cyclization under acidic conditions leads to the benzazepine


76. The benzylic methoxy group is then removed by metal-ammonia reduction.¹¹ Alkylation with p-nitrophenethyl bromide would then give the intermediate 78. Reduction of the nitro group would thus afford verilopam (79). The same sequence starting with amine 80 affords anilopam (81).




The good analgesic efficacy observed with ciramadol (87) and doxpicomine (92) show that location of the basic center directly on the quaternary benzylic center is quite consistent with activity. It is interesting to note in this connection that compound 82, in which nitrogen is similarly located, shows analgesic potency in the range of sulfentanyl that is, some five decade orders of magnitude greater than morphine.¹²



Aldol condensation of the methoxymethyl ether of *m*-methoxybenzaldehyde (83) with cyclohexanone affords the conjugated ketone 84. Michael addition of dimethylamine leads to the aminoketone 85. Reduction of the ketone proceeds stereospecifically to afford the *cis* aminoalcohol 86. Mild hydrolysis of the product gives the free phenol, ciramadol (87).¹³



In a similar vein, Knoevenagel condensation of nicotinaldehyde (88) with diethyl malonate gives the unsaturated ester 89. Michael addition of dimethylamine gives the corresponding aminoester (90). Reduction of the carbonyl groups with lithium aluminum hydride affords the glycol 91. Formation of the acetal between the diol and acetone gives the analgesic agent doxipicomine (92).¹⁴



REFERENCES

1. M. P. Kotick, D. L. Leland, J. O. Polazzi, and R. N. Schut, J. Med. Chem., **23**, 166 (1980).
2. See, for example, J. A. Baltrop, J. Chem. Soc., **1947** 349.
3. T. A. Montzka, J. D. Matiskella and R. A. Partyka, U.S. Patent 4,246,413; Chem. Abstr. **95**, 43442z (1981).
4. T. A. Montzka and J. D. Matiskella, German Offen. 2,517,220, Chem. Abstr. **84**, 59832k (1976).
5. W. F. Michne, T. R. Lewis, S. J. Michalec, A. K. Pierson, and F. J. Rosenberg, J. Med. Chem., **22**, 1158 (1979).
6. P. G. H. VanDaele, M. L. F. DeBruyn, J. M. Boey, S. Sanczuk, J. T. M. Agten, and P. A. J. Janssen, Arzneim. Forsch., **26** 1521 (1971).
7. W. F. M. VanBever, C. J. E. Niemegeers, and P. A. J. Janssen, J. Med. Chem., **17**, 1047 (1974).

8. W. F. M. VanBever, C. J. E. Niemegeers, K. H. Schellekens, and P. A. J. Janssen, Arzneim. Forsch., 26, 1548 (1976).
9. D. M. Zimmerman and W. S. Marshall, U.S. Patent 4,029,796; Chem. Abstr. 87, 102192 (1977).
10. J. W. Epstein, H. J. Brabander, W. J. Fanshawe, C. M. Hofmann, T. C. McKenzie, S. R. Safir, A. C. Usterberg, D. B. Cosulich, and F. M. Lovell, J. Med. Chem., 24, 481 (1981).
11. T. A. Davison and R. D. Griffith, German Offen 2,946,794; Chem. Abstr. 93, 186198 (198).
12. D. Lednicer and P. F. VonVoigtlander, J. Med. Chem., 22, 1157 (1979).
13. J. P. Yardley, H. Fletcher, III, and P. B. Russell, Experientia 34, 1124 (1978).
14. R. N. Booher, S. E. Smits, W. W. Turner, Jr., and A. Pohland, J. Med. Chem., 20, 885 (1977).


8 Five-Membered Heterocycles

A five-membered heterocyclic ring packs a relatively large number of polarized bonds into a relatively small molecular space. This provides a convenient framework to which to attach necessary side chains. In some cases, the framework itself is believed to be part of the pharmacophore.


1. PYRROLES AND PYRROLIDINES

In recent years increasing attention has been paid to the possibility of delaying or even reversing the memory loss that accompanies old age or the more tragic loss of human capabilities associated with premature senility - Alzheimer's disease. Progress is hampered by the difficulty of identifying suitable animal tests, and there is presently no reliable therapy.


A series of pyrrolidones shows promise of being cognition-enhancing agents. One of these, amacetam (3), is synthesized readily by ester-amide exchange between ethyl 2-oxo-1-pyrrolidineacetate (1) and N,N-diisopropylethylenediamine (2).¹



A tragic amount of morbidity and mortality is associated with high blood pressure. Many drugs operating by a wide variety of mechanisms have been employed to control this condition. Of the biochemical mechanisms employed by the body to maintain blood pressure an important one involves conversion of a kidney protein, angiotensinogen, to the pressor hormone angiotensin by a series of enzymes. The last step in the activation involves cleavage of angiotensin I to the much more active angiotensin II by the so called angiotensin-converting enzyme. It was believed that inhibitors of converting enzyme would have anti-hypertensive activity. Captopril (6), designed expressly for this purpose, has found exceptional clinical acceptance. One of the several syntheses of this fairly simple molecule involves amidation of t-butyl prolineate by 3-thio-2-methylpropionic acid (4) followed by acid treatment of the protected intermediate (5) to give captopril (6).²



A nonsteroidal antiinflammatory agent in which the benzene ring carrying the acetic acid moiety has been replaced by a pyrrole grouping is zomepirac (10). It is synthesized from




diethyl acetonedicarboxylate, chloroacetone, and aqueous methylamine via modification of the Hantzsch pyrrole synthesis to give key intermediate 7. Saponification, monoesterification and thermal decarboxylation give ester 8. This is acylated with *N,N*-dimethyl-*p*-chlorobenzamide (to give 9) and, finally, saponification gives zomepirac (10).³

Treatment of *p*-benzoquinone with 1-pyrrolidinylamine provides a convenient synthesis of the immunoregulator and antibacterial agent, azarole (11).⁴


2. FURANS

A biarylpropionic acid derivative containing a furan ring as a prominent feature has antiinflammatory activity. The patented synthesis involves a straightforward organometallic addition of ethyl lithioacetate to aldehyde 12 followed by saponification




to give orpanoxin (13).⁵

Installation of a different side chain completely alters the pharmacological profile leading to a new class of muscle relaxants. The synthesis begins with copper(II)-promoted diazonium coupling between furfural (14) and 3,4-dichlorobenzene-diazonium chloride (15) to give biarylaldehyde 16. Next, condensation with 1-aminohydantoin produces the muscle relaxant clodanolene (17).⁶




The pharmacological versatility of this general substitution strategy is further illustrated by diazonium coupling of 14 with 2-nitrobenzenediazonium chloride to produce biarylaldehyde 18. Formation of the oxime with hydroxylamine is followed by dehydration to the nitrile. Reaction with anhydrous methanolic hydrogen chloride leads to imino ether 19 and addition-elimination of ammonia leads to the antidepressant amidine, nitrafudam (20).⁷



The presence of a furan ring is also compatible with


cimetidine-like antiulcer activity, despite the prominent possession of an imidazole ring by histamine, the natural agonist, which served as a structural point of departure for the histamine H_2 antagonists. The synthesis of ranitidine (25) begins with an acid-catalyzed displacement of the primary alcoholic linkage of 21 with cysteamine to give 22. An addition-elimination reaction with 23 (itself made from 24 by an addition-elimination reaction) completes the synthesis of ranitidine (25).⁸



3. IMIDAZOLES


Amoebal infections, particularly of farm animals and the female human genitalia, are at best only annoying. All too often the problem encountered leads to difficult diarrheas. A group of nitroimidazoles have activity against the causative organisms and consequently have been widely synthesized.

One such agent is synthesized from 2-methylimidazole by reaction with epichlorohydrin under acidic conditions. This produces the antiprotozoal agent ornidazole (26).⁹





Similarly, reaction of 2-nitroimidazole with 1,2-epoxy-3-methoxypropane in the presence of potassium carbonate gives misonidazole (27).¹⁰ This agent also has the interesting and potentially useful additional property of sensitizing hypoxic tumor cells to ionizing radiation.

Many nitroimidazoles possess antiprotozoal activity. One of these is bamnidazole (29). Synthesis involves reaction of imidazole carbonate 28 with ammonia.¹¹




Removal of the nitro group results in an alteration of antimicrobial spectrum leading to a series of antifungal agents. For example, reaction of 2,4-dichloroacetophenone with glycerol and tosic acid leads to dioxolane 30. Under brominating conditions, sufficient carbonyl-like character exists to allow transformation of 30 to 31 and this product, after esterification, undergoes displacement to 32 with imidazole. Saponification and reaction with mesyl chloride then give 33. The synthesis of antifungal ketoconazole (34) then concludes by displacement with the phenolate derived from 4-acetylpiperazinylphenol.¹²









Displacement of bromine on phenacyl halide 35 with imidazole gives 36. Reduction with sodium borohydride followed by displacement with 2,6-dichloro-benzyl alcohol in HMPA then produces antifungal orconazole (37).¹³




If the displacement reaction is carried out between imidazole derivative 38 and thiophene analogue 39, the antifungal agent tiaconazole (40) results.¹⁴ A rather slight variant of this sequence produces antifungal sulconazole (41).¹⁵ Obvious variants of the route explicated above for ketoconazole (34) lead to parconazole (42)¹⁶ and doconazole (43),¹⁷ instead.

Insertion of a longer spacer is compatible with antifungal activity. Reaction of epichlorohydrin with 4-chlorobenzylmagnesium chloride leads to substituted phenylbutane 44. Dis-







placement with sodium imidazole, conversion of the secondary alcohol group to the chloride (thionyl chloride), and displacement with 2,6-dichlorothiophenolate concludes the synthesis of antifungal butoconazole (45).¹⁸



Progressive departure from the fundamental structure of the lead agent cimetidine led to the antiulcer agent oxmetidine (47). The synthesis involves S-methylation (CH_3I) of the 2-thiouracil intermediate 46 and is followed by an addition-elimination reaction with 2-(5-methyl-4-imidazolylmethylthio) ethylamine to give oxmetidine (47).¹⁹




Another entry into the antiulcer sweepstakes is etinfidine (50). It is synthesized by displacement of halide from 4-chloromethyl-5-methylimidazole (48) with substituted thiol 49. The latter is itself made from thiourea analogue 51 by an addition-elimination reaction with cysteamine 52.²⁰




The imidazole-containing hypnotic/injectable anesthetic agent etomidate (58) is synthesized from 1-amino-1-phenylethane starting with triethylamine mediated displacement with chloroacetonitrile leading to secondary amine 53. The d-enantiomer is preferred as starting material. This is converted to the formamide (54) on heating with formic acid. Next, the active methylene group is formylated by reaction of 54 with ethyl formate and sodium methoxide in order to give 55. The now superfluous N-formyl group is removed and the imidazole ring is established upon reaction of 55 with potassium thiocyanate. The key intermediate in this transformation is probably thio-urea 55a. Oxidative desulfurization occurs on treatment of 56 with a mixture of sulfuric and nitric acids and the resulting 57 is subjected to amide-ester interchange with anhydrous ethanolic hydrogen chloride to complete the synthesis of

etomidate (58).²¹

It is possible to form 2-imino-4-imidazolines, such as 59, in situ from creatinine. Treatment of this heterocycle with 3-chlorophenylisocyanate leads to a sedative agent, fenobam (60).²²




A substituted thiazole ring attached to a reduced imidazole moiety is present in a compound that displays antihypertensive activity. Reaction of thiourea 61 with methyl iodide to



give the corresponding S-methyl analogue, followed by heating with ethylenediamine, completes the synthesis of tiamenidine (62).²³


4. TRIAZOLES

Insertion of a triazole ring in place of an imidazole ring is consistent in some cases with retention of antifungal activity. The synthesis of one such agent, azoconazole (64), proceeds simply by displacement of halide 63 with 1,2,4-triazole.²⁴ The route to terconazole (65) is rather like that to ketoconazole (34).²⁵




5. PYRAZOLINES

Reaction of substituted hydrazine analogue 66 with protected β -dicarbonyl compound 67 leads to a ring-forming two-site reaction and formation of the pyrazoline diuretic agent, muzolimine (68).²⁶




As a bioisosteric replacement for a substituted pyrrole ring, a pyrazole ring is a key feature of the nonsteroidal

concludes by acylation with isopropyl chloride.²⁸




7. TETRAZOLES

Conversion of *m*-bromobenzonitrile to the tetrazole and addition of the elements of acrylic acid gives 75, starting material for the patented synthesis of the antiinflammatory agent, broperamole (76). The synthesis concludes by activation with thionyl chloride and a Schotten-Baumann condensation with piperidine.²⁹




8. MISCELLANEOUS

Ropitoin (79) is an antiarrhythmic compound containing a hydantoin ring. Its synthesis is accomplished by alkylating 77 with chloride 78 with the aid of sodium methoxide.³⁰




Reaction of ethyl cyanoacetate with ethyl thiolacetate produces a Z and E mixture of the dihydrothiazole derivative 80. This is N-alkylated with methyl iodide and base (81), the active methylene group is brominated (82), and then a displacement with piperidine (83) is performed. Hydrolysis completes the synthesis of the diuretic agent, ozolinone (84).³¹

Finally, a mesoionic syndnone, molsidomine (88), is active as an antianginal agent. Its synthesis starts by reacting 1-aminomorpholine with formaldehyde and hydrogen cyanide to give 85. Nitrosation gives the N-nitroso analogue (86) which



cyclizes to the syndnone (87) on treatment with anhydrous acid. Formation of the ethyl carbonate with ethyl chlorocarbonate completes the synthesis of molsidomine (88).³²



REFERENCES

1. Y. L. L'Italien and I. C. Nordin, U.S. Patent 4,145,347 (1979); Chem. Abstr., 91, 39332p (1979).
2. M. A. Ondetti, B. Rubin, and D. W. Cushman, Science, 196, 441 (1977); Anon., Belgian Patent 851,361.
3. J. R. Carson and S. Wong, J. Med. Chem., 16, 172 (1973).
4. R. E. Johnson, Belgian Patent 16,542 (1974); Chem. Abstr., 83, 97007g (1975).
5. S. S. Pelosi, Jr., U.S. Patent 3,962,284 (1976); Chem. Abstr., 85, 159860g (1976).
6. H. R. Snyder, Jr., C. S. Davis, R. K. Bickerton, and R. P. Halliday, J. Med. Chem., 10, 807 (1967).
7. S. S. Petosi, Jr., R. E. White, R. L. White, G. C. Wright, and C.-N. You, U.S. Patent 3,919,231 (1975); Chem. Abstr., 84, 59168y (1976).
8. Anon., French Patent 2,384,765 (1980); Chem. Abstr., 92, 58595p (1980).
9. M. Hoffer and E. Grunberg, J. Med. Chem., 17, 1019 (1974).
10. A. G. Beaman and W. P. Tantz, U.S. Patent 3,865,823 (1975); Chem. Abstr., 82, 170944w (1975).
11. C. Jeanmart and M. N. Messer, German Offen. 2,035,573 (1969); Chem. Abstr., 74, 100044p (1971).
12. J. Heeres, L. J. J. Backx, J. H. Mostmans, and J. van Cutsem, J. Med. Chem., 22, 1003 (1979).
13. E. F. Godefroi, J. Heeres, J. Van Cutsem, and P. A. J. Janssen, J. Med. Chem., 12, 784 (1969).
14. G. E. Gymer, U.S. Patent 4,062,966.
15. K. A. M. Walker and M. Marx, U.S. Patent 4,038,409 (1976); Chem. Abstr., 87, 152210c (1977).
16. J. Heeres, U.S. Patent 3,936,470.

17. J. Heeres, German Offen. 2,602,770 (1976); Chem. Abstr., 86, 29811b (1977).
18. K. A. M. Walker, A. C. Braemer, S. Hitt, R. E. Jones, and T. R. Matthews, J. Med. Chem., 21, 840 (1978).
19. T. H. Brown, G. J. Durant, J. C. Emmett, and C. R. Ganellin, U.S. Patent 4,145,546 (1979); Chem. Abstr., 90, 204137t (1979).
20. R. R. Crenshaw, G. Kavadias, and R. F. Santonge, U.S. Patent 4,157,340 (1979); Chem. Abstr., 91, 157312e (1979).
21. E. F. Godefroi, P. A. J. Janssen, C. A. M. Van der Eycken, A. H. M. T. Van Heertum, and C. J. E. Niemegeers, J. Med. Chem., 8, 220 (1965).
22. C. R. Rasmussen, U.S. Patent 3,983,135 (1976); Chem. Abstr., 86, 55441a (1977).
23. H. Rippel, H. Ruschig, E. Lindner, and M. Schorr, German Offen. 1,941,761 (1971); Chem. Abstr., 74, 100054s (1971).
24. G. Van Reet, J. Heeres, and L. Wals, German Offen. 2,551,560 (1976); Chem. Abstr., 85, 94368f (1976).
25. J. Heeres, L. J. J. Backx, and J. H. Mostmans, German Offen. 2,804,096 (1978); Chem. Abstr., 89, 180014b (1978).
26. E. Moeller, K. Meng, E. Wehinger, and H. Horstmann, German Offen. DE 2,319,278 (1974); Chem. Abstr., 82, 57690x (1975).
27. H. Biere, E. Schroder, H. Ahrens, J.-F. Kapp, and I. Bottcher, Eur. J. Med. Chem., 17, 27 (1982).
28. W. J. Ross, R. G. Harrison, M. J. R. Jolley, M. C. Neville, A. Todd, J. P. Berge, W. Dawson, and W. J. F. Sweatman, J. Med. Chem., 22, 412 (1979).
29. Anon., British Patent 1,319,357 (1973); Chem. Abstr., 79, 92231h (1973).

30. S. Hayao, H. J. Havera, and W. G. Stryker, U.S. Patent 4,006,232 (1977); Chem. Abstr., 87, 5968c (1977).
32. K. Masuda, T. Kamiya, Y. Imashiro, and T. Kaneko, Chem. Pharm. Bull., 19, 72 (1971).

9 Six-Membered Heterocycles

The five-membered heterocycles discussed in the preceding chapter more often than not constituted the pharmacophoric moieties of the drugs in question. Drugs based on six-membered rings, on the other hand, constitute a somewhat more diverse group. In many cases such as the dihydropyridines and the antibacterial pyrimidines, the ring system again provides the pharmacophore; this section, however is replete with agents in which the heterocyclic ring simply serves as surrogate for an aromatic ring.


1. PYRIDINES

Derivatives of anthranilic acid have a venerable history as nonsteroid antiinflammatory agents. It is thus not surprising that the corresponding derivatives in which phenyl is replaced by pyridinyl show much the same activity.

Drugs that are too highly hydrophilic are often absorbed rather poorly from the gastrointestinal tract. It is sometimes possible to circumvent this difficulty by preparing esters of such compounds so as to change their water lipid partition characteristics in order to enhance absorption. Once absorbed, the esters are cleaved by the numerous esterase enzymes in the bloodstream, releasing free drug.

Preparation of the first of these antiinflammatory prodrugs starts with the displacement of halogen on bromophthalide 2 by the anion of the nicotinic acid derivative 1. Reaction of the intermediate 3 with aniline 4 leads to formation of talniflumate (5).¹


In much the same vein, the basic ester 7 can be obtained by reaction of the same chloroacid with morpholine derivative 6. Reaction with aniline 4 affords morniflumate (8).




Congestive heart failure represents the end result of a


complex process which leads eventually to death when the heart muscle is no longer able to perform its function as a pump. Cardiotonic agents such as digitalis (see Steroids) have proved of value in treatment of this disease by stimulating cardiac muscle. The toxicity of these agents has led to an extensive search for alternate drugs. The bipyridyl derivative amrinone has shown promise in the clinic as a cardiotonic agent.

Starting material for the synthesis is the enaminaldehyde 10, obtainable by some version of the Villsmeyer reaction on picoline derivative 9. Condensation of that with cyanoacetamide in the presence of methoxide leads to pyridone 12. The reaction can be rationalized by assuming that the initial step consists in Michael addition of the anion from acetamide to the acrolein; elimination of dimethylamine would afford the intermediate 11. Condensation of amide nitrogen with the aldehyde leads to the observed product. Saponification of the nitrile then gives acid 13. Treatment under nitrating conditions leads to an interesting reaction that results ultimately in replacement of the carboxyl by a nitro group (14). Reduction of that last function affords the amine; there is thus obtained amrinone (15).²






Organic nitrite esters such as nitroglycerin have constituted standard treatment for anginal attacks for the better part of the century. These agents, which are thought to owe their efficacy to reducing the oxygen demand of the heart, as a class suffer from poor absorption and very short duration of action. Nicordanil (18) represents an attempt to overcome these shortcomings by combining a nitrite with nicotine, the latter moiety itself having some vasodilating action. The drug is obtained in straightforward manner by reaction of nicotinoyl chloride (16) with the nitrite ester (17) of ethanolamine.³




The excessive production of sebum associated with acne has made life miserable for many an adolescent. Research on acne has, as a rule, concentrated on therapy rather than prophylaxis. A pyrimidone forms an interesting exception, being described as an antiseborrheic agent. Starting keto-ester 21 can, at least in principle, be obtained by γ -

acylation of the anion from acrylate 19 with a derivative of acid 20. Reaction with hydroxylamine under basic conditions would afford initially the oxime 22. This cyclizes to the N-hydroxypyridine, piroctone (23) under the reaction conditions.⁴





The so-called calcium channel blockers constitute a class of cardiovascular agents that have gained prominence in the past few years. These drugs, which obtund contraction of arterial vessels by preventing the movement of calcium ions needed for those contractions, have proved especially useful in the treatment of angina and hypertension. Dihydropyridines such as nifedipine (30) are particularly effective for these indications. A variation of the Hantzsch pyridine synthesis used to prepare the parent molecule provides access to unsymmetrically substituted dihydropyridines. Though preparation of such compounds by ester interchange of but one ester has been described, these schemes are marked by lack of selectivity and low yields. Thus condensation of enaminoester 23 (obtained from the corresponding acetoacetate) with acetoacetate 24 and benzaldehyde 25 affords nimodipine (26).⁵ In a similar

sequence, condensation of the enaminoester from methyl acetoacetate (28) with acetoacetate 27 and benzaldehyde gives the calcium channel blocker nicaripidine (29).⁶




p-Fluorobutyrophenone derivatives of phenylpiperidines constitute a class of very effective antipsychotic agents. It is thus interesting to note that activity is retained when a carbonyl group is inserted between phenyl and the piperidine ring. The starting benzoylpiperidine 32 can be obtained by any of several schemes starting with the reduced derivative of isonicotinic acid 31. Alkylation with bromoacetal 33 leads to the tertiary amine 34. Hydrolysis of the acetal group leads to cloperone (35).⁷






2. PYRIDAZINES

As noted earlier (Chapter 2) β -adrenergic blocking agents have found extensive use in the treatment of hypertension. Drawbacks of these drugs include relatively slow onset of action and efficacy in only about half the patients treated. In an effort to overcome these shortcomings, considerable research has been devoted to β -blockers which incorporate moieties associated with direct-acting vasodilating agents. (On the other side of the coin, the β -blocking moiety should control some of the side effects characteristic of the vasodilators, such as increase in heart rate.) Condensation of ketoester 36 with hydrazine affords the corresponding pyridazinone 37. Reaction of that with phosphorus oxychloride leads to the chloro derivative 38. The target compound could then be obtained by first reacting the phenol with epichlorohydrin to give epoxide 39. Opening of the oxirane with tertiary butylamine would then complete construction of the β -blocking side chain (40). Displacement of chlorine by hydrazine then affords prizidilol (41).⁸




3. PYRIMIDINES


Though the great majority of antiinflammatory agents contain some form of acidic proton, occasional compounds devoid of such a function do show that activity. Thus the nonacidic pyrazolylpyrimidine epirazole (47) is described as a nonsteroid antiinflammatory agent. Reaction of pyrimidinone 42 with phosphorus oxychloride leads to the chloro derivative 43. Replacement of halogen with hydrazine gives the intermediate 44. Reaction of that with the methyl acetoacetate derivative 45 (obtained, for example, by pyrolysis of the orthoester) leads to formation of the pyrazole ring. The reaction may be rationalized by assuming initial formation of hydrazone 46; addition of the more basic hydrazine nitrogen to the masked carbonyl group followed by elimination of methoxide gives the observed product: there is thus obtained epirazole (47).⁹




Two closely related diaminopyrimidines have been described as antineoplastic agents. In the absence of specific references, one may speculate that these can be prepared by a general method for the synthesis of aryl diaminopyrimidines.¹⁰ Thus acylation of arylacetone (48) with ethyl acetate affords the corresponding cyanoketone (49). Reaction of that intermediate with guanidine can be visualized as first involving formation of the imine derivative (50); addition of a second amino group from guanidine to the nitrile gives the cyclized derivative (51); tautomerization then gives the observed product, metoprine (52). The same sequence starting with ethyl propionate instead of ethyl acetate will lead to etoprine (53).



Interposition of a methylene group between the phenyl ring and the heterocycle leads to the benzyldiaminopyrimidines, a class of compounds notable for their anti-bacterial activity. Condensation of hydrocinnamate 54 with ethyl formate leads to the hydroxymethylene derivative 55. In this case, too, the heterocyclic ring is formed by reaction with guanidine. This sequence probably involves initial addition-elimination to the formyl carbon to form 56; cyclization in this case involves simple amide formation. Tautomerization then affords the hydroxy derivative 57. This is converted to tetroxoprim (58) by first replacing the hydroxyl by chlorine and then displacement of halogen with ammonia.¹¹




Preparation of the analogue metioprim involves an alternate approach. Aldol condensation of aldehyde 59 with propionitrile 60 gives the cinnamitrile 61. Reaction of this intermediate with guanidine probably involves displacement of the allylic aniline group as the first step (62). Cyclization followed by tautomerization affords metioprim (63).¹²




The pyrimidine 5-fluorouracil (**64**) is used extensively in the clinic as an antimetabolite antitumor agent. As a consequence of poor absorption by the oral route, the drug is usually administered by the intravenous route. A rather simple latentiated derivative, tegafur (**66**), has overcome this limitation by providing good oral absorption. Reaction of 5-fluorouracil with trimethylsilyl chloride in the presence of base gives the disilylated derivative (**65**). Reaction of this with dihydrofuran (obtained by dehydro-

halogenation of 2-chlorofuran) in the presence of stannic chloride affords directly tegafur (66).¹³




An aryloxypyrimidone has been described as an antiulcer agent; this activity is of note since the agent does not bear any structural relation to better known antiulcer drugs. Displacement of halogen on the acetal of chloroacetaldehyde by alkoxide from *m*-cresol gives the intermediate 67. This affords enaminoaldehyde 68 when subjected to the conditions of the Vilsmeier reaction and subsequent hydrolysis. Condensation with urea may be rationalized by assuming the first step to involve displacement of the dimethylamino group by an addition-elimination sequence (69). Ring closure then leads to the pyrimidone and thus tolimidone (70).¹⁴




4. MISCELLANEOUS HETEROCYCLES


A cinnamoylpiperazine is described as an antianginal agent. The key intermediate 73 can, in principle, be obtained by alkylation of the monobenzyl derivative of piperazine 71 with ethyl bromoacetate (72). Removal of the protecting group then affords the substituted piperazine (73). Acylation of this with 3,4,5-trimethoxycinnamoyl chloride gives cinepazet (74).¹⁵



Though dental afflictions constitute a very significant disease entity, these have received relatively little attention from medicinal chemists. (The fluoride tooth-pastes may form an important exception.) This therapeutic target is, however, sufficiently important to be the focus of at least some research. A highly functionalized piperazine derivative that has come out of such work shows prophylactic activity against dental caries. Condensation of the enol ether 75 of thiourea with n-pentylisocyanate gives the addition product 77. Reaction of this with diamine 78, derived from piperazine, leads to substitution of the methylthio moiety by the primary amine, in all likelihood by an addition-elimination sequence. There is thus obtained ipexidine (79).¹⁶




Heterocycles that carry p-anisyl groups on adjacent positions such as indoxole¹⁷ and flumizole¹⁷ constitute an important subclass among the nonsteroid antiinflammatory agents that do not possess an acidic proton. It is thus not very surprising to note that a similarly substituted 1,2,4-triazine also shows antiinflammatory activity. Condensation of the dibenzyl derivative 80 with semicarbazine affords the heterocyclic ring directly (82). Reaction with phosphorus oxychloride serves to convert the hydroxyl to chloro (83). Taking advantage of a reaction pioneered by Taylor, this intermediate is then reacted with an excess of the ylide from methyltriphenylphosphonium bromide. The first equivalent in all probability displaces halogen to form the substituted phosphonium salt 84. This is then converted to its ylide by excess phosphorane. Hydrolysis leads to loss of triphenylphosphine oxide. There is thus obtained anitrazafen (85).¹⁸






Acetylcholine is one of the fundamental neurotransmitters involved in a wide variety of normal regulatory functions. A number of disease states that may be associated with local excesses of this compound can at least in theory be treated by suppressing its action. Anticholinergic drugs have in practice proved of limited utility because it has been difficult to devise molecules that show much selectivity. The very widespread distribution of necessary cholinergic responses leads to the manifestation of a multitude of side effects when anticholinergic drugs are used in therapy. However, a number of syndromes could in principle, be treated with these drugs if they are applied by the topical route; lack of systemic absorption should avoid the side effects. It should, for example, be possible to treat the stomach lining to suppress the cholinergically mediated acid secretion associated with gastric ulcers. By the same token, direct administration to

the lung should prevent the bronchoconstriction associated with asthma. Considerable work based on this concept has been occasioned by the observation that quaternary salts of atropine (93), which should not be absorbed systemically, do retain the anticholinergic activity of the parent base. One such salt, ipratropium bromide (92), has undergone considerable clinical investigation as an antiasthmatic agent administered by insufflation (*i.e.*, topical application to the bronchioles). The fact that the stereochemistry of this agent is the opposite from that which would be obtained by direct alkylation with isopropyl bromide requires that a somewhat longer sequence be employed for its synthesis.

Preparation of the key tropine 86 is achieved by any one of several variations on the method first developed by Robinson, which involves reaction of a primary amine with dihydroxyacetone and glyoxal. Reduction of the carbonyl group in the product (86) followed by acylation affords the aminoester (88). Transesterification with ester aldehyde 89



leads to 90. The ester is then reduced to the atropic ester 91 by means of borohydride.¹⁹ Attack of methyl bromide occurs from the more open face of the molecule to give ipratropium bromide (92).



Until the advent of the antitumor antibiotics, alkylating agents were the mainstay of cancer chemotherapy. The alkylating drug cyclophosphamide (100) found probably more widespread use than any other agent of this class. Two closely related agents, ifosfamide (96) and trofosfamide (97), show very similar activity; clinical development of these drugs hinges on the observation that the newer agents may show efficacy on some tumors that do not respond to the prototype. The common intermediate (95) to both drugs can be obtained from reaction of phosphorus oxychloride with amino alcohol 94. Reaction of the oxazaphosphorane oxide with 2-chloroethylamine gives ifosfamide (96); displacement on bis(2-chloroethyl)amine gives trofosfamide (97).²⁰ In an alternate synthesis, the phosphorane is first condensed with the appropriate amino alcohols to give respectively 98 and 99. These are then converted to the nitrogen mustards by reaction with mesyl chloride, or thionyl chloride.

REFERENCES

1. M. A. Los, U.S. Patent 4,255,581; Chem. Abstr., 95, 62002x (1981).
2. G. Y. Leshner and C. J. Opalka, Jr., U.S. Patent 4,107,315; Chem. Abstr. 90, 103844r (1979).
3. H. Nagano, T. Mori, S. Takaku, I. Matsunaga, T. Kujirai, T. Ogasawara, S. Sugano, and M. Shindo, German Offen, 4,714,713; Chem. Abstr., 88, 22652h (1978).
4. G. Lohaus and W. Dittmar, German Patent 1,795,831; Chem. Abstr., 89, 197347k (1978).
5. E. Wehinger, H. Meyer, F. Bossert, W. Vater, R. Towart, K. Stoepe1, and S. Kazada, German Offen, 2,935,451; Chem. Abstr. 95, 42922u (1981).
6. Anon. Japanese Patent, 74,109,384; Chem. Abstr. 82, 170642c (1975).
7. J. W. Ward and C. A. Leonard, French Demande 2,227,868; Chem. Abstr. 82, 170720v (1975).
8. B. L. Lam, Eur. Pat. Appl. EP 47,164; Chem. Abstr., 96, 217866d (1982).
9. Y. Morita, Y. Samejima, and S. Shimada, Japanese Patent 73 72,176; Chem. Abstr., 79, 137187s (1973).
10. B. Roth and J. Z. Sterlitz, J. Org. Chem., 34, 821 (1969).
11. W. Liebenow and J. Prikryl, French Demande 2,221,147; Chem. Abstr., 82, 156363z (1975).
12. Anon. Belgian Patent 865,834; Chem. Abstr., 90, 54971u (1979).
13. S. Hillers, R. A. Zhuk, A. Berzina, L. Serina, and A. Lazdins; U.S. Patent 3,912,734; Chem. Abstr., 84, 59538u (1976).

14. C. A. Lipinski, J. G. Stam, G. D. DeAngelis, and H. J. E. Hess; U.S. Patent 3,922,345; Chem. Abstr., 84, 59552u (1976).
15. F. Faurau, G. Huguet, G. Raynaud, B. Pourfas, and M. Turin; British Patent 1,168,108; Chem. Abstr., 72, 12768 (1970).
16. R. A. Wohl; South African Patent 7,706,373; Chem. Abstr., 90, 54980 (1979).
17. D. Lednicer and L. A. Mitscher, "The Organic Chemistry of Drug Synthesis", Vol. 2, Wiley, New York, 1980, p. 255
18. W. B. Lacefield and P. P. K. Ho, Belgian Patent 839,469; Chem. Abstr., 87, 68431 (1977).
19. W. Schultz, R. Banholzer, and K. H. Pook, Arzneim Forsch. 26, 960 (1976).
20. H. Arnold, N. Brock, F. Bourreaux, and H. Bekel, U.S. Patent 3,732,340; Chem. Abstr., 79, 18772 (1973).


10 Five-Membered Heterocycles Fused to Benzene

1. INDOLES


Though the therapeutic utility of aspirin has been recognized for well over a century, this venerable drug was not classified as a nonsteroid antiinflammatory until recently. The first drug to be so classified was in fact, indomethacin (1), a very useful agent introduced into clinical practice within the past two-score years. Much of the work that led to the elucidation of the mechanism of action of this class of therapeutic agents was in fact carried out using indomethacin. This drug is often considered the prototype of cyclooxygenase (prostaglandin synthetase) inhibitors; it is still probably the most widely used inhibitor in various pharmacological researches. The undoubted good efficacy of the drug in the treatment of arthritis and inflammation at the same

time has led to very widespread use in medical practice.


The relatively short duration of action of indomethacin resulted in various attempts to develop prodrugs so as to overcome this drawback. One of these consists of an amino acid derivative. Thus, reaction of the drug with the chlorocarbonate derivative of dimethylethanol(2) affords the mixed anhydride 3. Reaction of that reactive intermediate with serine (4) leads directly to sermatacin (5).¹




Replacement of chlorine on the pendant benzoyl group by azide is apparently consistent with antiinflammatory activity. Acylation of indomethacin intermediate 6 with p-nitrobenzoyl chloride leads to the corresponding amide (7). Saponification (8) followed by reduction of the nitro group gives the amine 9. The diazonium salt (10) obtained on treatment with nitrous acid is then reacted with sodium azide; there is thus obtained zidomethacin (11).¹²





Serotonin (12) is a ubiquitous endogenous compound that has a multitude of biological activities. For example, the compound lowers in certain biological tests. A compound that would lead to a serotonin derivative after decarboxylation has been described as an antihypertensive agent. (Note, however, that decarboxylation would have to occur by a mechanism different from the well-known biosynthetic loss of carbon dioxide from α -amino acids.) Mannich reaction on indole 13 with formaldehyde and dimethylamine gives the gramine derivative 14. Reaction with cyanide leads to replacement of the dimethylamino group to give the nitrile 15. Condensation of that intermediate with dimethyl carbonate and base gives the corresponding ester (16). Catalytic reduction of the nitrile group (17) followed by saponification affords indorenate (18).³






Two closely related indoles fused to an additional saturated ring have been described as CNS agents. The first of these is obtained in straightforward manner by Fischer indole condensation of functionalized cyclohexanone 20 with phenylhydrazine (19). The product, cyclindole (21) shows antidepressant activity.⁴ The fluorinated analogue flucindole (26) can be prepared by the same scheme. An alternate route starting from a somewhat more readily available intermediate involves as the first step Fischer condensation of substituted phenylhydrazine 22 with 4-hydroxycyclohexanone (23). The resulting alcohol (24) is then converted to its tosylate (25). Displacement by means of dimethylamine leads to the antipsychotic agent flucindole (26).⁵







Changing the functionality on the alicyclic ring from an amine to a carboxylic acid leads to a compound that shows antiallergic activity, acting possibly by means of inhibition of the release of allergic mediators. Thus, condensation of acylated indole 27 with cyclohexanone carboxylic acid 28 affords directly oxarbazole (29).⁶




A fully unsaturated tricyclic indole derivative serves as the aromatic moiety for a nonsteroid anti-inflammatory agent. Preparation of this compound starts with the Michael addition of the anion from methyl diethylmalonate to cyclohexanone. The product (32) is then hydrolyzed and decarboxylated to give ketoester 33. Fischer condensation with p-chlorophenylhydrazine leads to the indole 34. This is then esterified (35) and dehydrogenated to the carbazole 36. Saponification leads to the acid and thus carprofen (37).⁷




The salicylic acid functionality incorporated in a rather complex molecule interestingly leads to a compound that exhibits much the same activity as the parent. The 1,4 diketone required for formation of the pyrrole ring can be obtained by alkylation of the enamine from 2-tetralone (38) with phenacyl bromide. Condensation of the product, 39, with salicylic acid derivative 40 leads to the requisite heterocyclic system (41). The acid is then esterified (42) and the compound dehydrogenated to the fully aromatic system (43). Saponification affords fendosal (44).⁸







An isoindolinone moiety forms part of the aromatic moiety of yet another antiinflammatory propionic acid derivative. Carboxylation of the anion from *p*-nitroethylbenzene (45) leads directly to the propionic acid (46). Reduction of the nitro group followed by condensation of the resulting aniline (47) with phthalic anhydride affords the corresponding phthalimide (48). Treatment of that intermediate with zinc in acetic acid interestingly results in reduction of only one of the carbonyl groups to afford the isoindolone. There is thus obtained indoprofen (49).⁹




2. BENZIMIDAZOLES

A series of benzimidazole and benzimidazolone derivatives from the Janssen laboratories has provided an unusually large number of biologically active compounds, particularly in the area of the central nervous system. Reaction of imidazolone itself with isopropenyl acetate leads to the singly protected imidazolone derivative 51. Alkylation of this with 3-chloro-1-bromopropane affords the functionalized derivative 52. Use of this interintermediate to alkylate piperidine 53 (see cloperone, Chapter 6) affords the derivative 54. Hydrolytic removal of the isopropenyl group then gives the veterinary sedative milenperone (55).¹⁰ The same sequence using p-fluorobenzoylpiperidine (56) gives the antipsychotic agent declenperone (57).¹⁰







Alkylation of the monobenzhydryl derivative of piperazine (58) with the same alkylating agent gives oxatimide (59), after removal of the protecting group.¹¹ This agent shows antihistaminic activity as well as some mediator release inhibiting activity, a combination of properties particularly useful for the treatment of asthma.




A somewhat more complex scheme is required for the preparation of benzimidazolones in which one of the nitrogen atoms is substituted by a 4-piperidyl group. The sequence starts with aromatic nucleophilic substitution on dichlorobenzene 60 by protected amino-piperidine derivative 61 to give 62. Reduction of the nitro group gives the diamine 63, which on treatment with urea affords the desired benzimidazolone 64.¹² The carbamate protecting group is then removed under basic conditions to give the secondary amine 65. Alkylation of this with the halide obtained by prior hydrolysis of

promising antiemetic agent.¹³




Piperidinobenzimidazole 65 also serves as starting material for the antipsychotic agent halopemide (69). In the absence of a specific reference, one may speculate that the first step involves alkylation with bromochloroethane to give halide 67. The chlorine may then be converted to the primary amine 68 by any of several methods such as reaction with phthalimide anion followed by hydrazinolysis. Acylation with p-fluorobenzoyl chloride then gives the desired product.

A still different scheme is used for the preparation of the benzimidazole buterizine (74). Alkylation of benzhydrylpiperazine 58 with substituted benzyl chloride 70 gives the intermediate 71. Nucleophilic aromatic displacement on this compound by means of ethylamine leads to 72; reduction of the nitro group then gives the diamine 73. Treatment of that with the orthoformate ester of pentanoic acid serves to form the imidazole ring. There is thus obtained the peripheral vasodilating agent buterizine (74).¹⁴




Amides and carbamates of 2-aminobenzimidazole have proved of considerable value as anthelmintic agents, particularly in veterinary practice. A very considerable number of these agents have been taken to the clinic in the search for commercially exploitable agents. (See the section on Benzimidazoles in Chapter 11 of Volume 2 of this series.) A small number of additional compounds have been prepared in attempts to uncover superior agents.

In a typical synthesis, reduction of the nitro group in starting material 75 leads to the corresponding diamine 76. Reaction with intermediate 77 obtained by acylation of the methyl ether of thiourea with methyl chloroformate, leads directly to fenbendazole (78).¹⁵




Friedel-Crafts acylation of fluorobenzene with thiophene-1-carboxylic acid gives the ketone 79. Nitration proceeds ortho to the fluoro group to give the intermediate 80. Nucleophilic displacement by means of ammonia (81) followed by reduction of the nitro group leads to the corresponding amine 81. Treatment of that with reagent 77 gives the anthelmintic agent nocodazole (83).¹⁶




It is of particular note that slight changes in the functionality of this last-named compound lead to a pro-

found change in biological activity. The agent in question, enviroxime, shows pronounced antiviral activity. The synthesis of this compound begins with the reaction of diamine 84 with cyanogen bromide. The reaction may be rationalized by assuming that cyanamide 85 is the initially formed product; addition of the remaining amine to the nitrile will give the observed product. Reaction of the anion obtained on treatment of 86 with sodium hydride with isopropylsulfonyl chloride apparently affords 87 as the sole product. (Note that minor tautomeric shifts could provide at least two alternate products.) Reaction with hydroxylamine affords the E oxime as the predominant product. There is thus obtained enviroxime (88).¹⁷ Examination of the isomeric oximes show the E isomer to be a good deal more active than the Z counterpart.



Incorporation of a 4-aminopiperidine moiety leads to a major change in biological activity. The agent obtained by this modification, astemizole (96) is a rather potent antihistaminic compound. Reaction of the isothiocyanate 89 with phenylenediamine under carefully

controlled conditions would lead to the thiourea 90. Alkylation with p-fluorobenzyl bromide then leads to the alkylated derivative 92. Cyclization of that intermediate gives the benzimidazole 93. The carbamate protecting group is then removed under basic conditions. Alkylation of the resulting secondary amine with substituted phenethyl bromide 95 proceeds to give astemizole (96).¹⁸





3. BENZOTHAZOLES

Bioisosteric relations constitute one of the more familiar tools in medicinal chemistry. There are thus sets of atoms that can often be interchanged without much

influence on the biological activity of the resulting molecules. In many series, for example, it may be quite useful to replace oxygen by sulfur; a sulfoxide sometimes serves in lieu of a ketone. Sulfur and nitrogen, on the other hand, are seldom considered to be a bioisosteric pair. It is thus of note that activity is retained in the antihelminthic compounds in the face of exactly such a substitution.

Reaction of aminothiophenol 97 with reagent 98 obtainable from phenylurea and thiophosgene leads directly to the anthelmintic agent frentizole (99).¹⁹ In much the same vein, condensation of 100 with reagent 101 affords tioxidazole (102).



REFERENCES

1. H. Biere, H. Arens, C. Rufer, E. Schroeder, and H. Koch, German Offen. 2,413,125; Chem. Abstr., **84**, 17731 (1976)
2. S. Tricerri, E. Panto, A. Bianchetti, G. Bossoni, and R. Venturini, Eur. J. Med. Chem., **14**, 181 (1979)

3. R. N. Schut, M. E. Safdy, and E. Hong, German Offen. 2,921,978; Chem. Abstr., 92, 128724f (1980).
4. A. Mooradian, U.S. Patent 3,959,309; Chem. Abstr., 85, 123759s (1976).
5. A. Mooradian, German Offen. 2,240,211 (1973); Chem. Abstr., 78, 136069x (1973).
6. J. E. Alexander and A. Mooradian, U.S. Patent 3,905,998; Chem. Abstr., 84, 74101 (1976).
7. L. Berger and A. J. Corraz, German Offen. 2,337,040; Chem. Abstr., 80, 108366g (1974).
8. R. C. Allen and V. B. Anderson, U.S. Patent 3,931,457; Chem. Abstr., 84, 105391r (1976).
9. G. Nannini, P. N. Giralidi, G. Malgara, G. Biasoli, F. Spinella, W. Logemann, E. Dradi, G. Zanni, A. Buttinoni, and A. Tommassini, Arzneim. Forsch. 23, 1090 (1973).
10. J. Vandenberg, L. E. J. Kennis, M. J. M. C. Van der Aa, and A. H. M. T. Van Heertum, German Offen. 2,645,125; Chem. Abstr., 87, 102326 (1977).
11. J. Vandenberg, L. E. J. Kennis, M. J. M. C. Van der Aa, and A. H. M. T. Van Heertum, German Offen. 2,714,437; Chem. Abstr., 88, 50920n (1978).
12. P. A. J. Janssen, A. H. M. T. Van Heertum, J. Vandenberg, and M. J. M. C. Van der Aa, German Offen. 2,257,261; Chem. Abstr., 84, 135657 (1976).
13. J. Vandenberg, L. E. J. Kennis, M. J. M. C. Van der Aa, and A. H. M. T. Van Heertum, German Offen. 2,632,870; Chem. Abstr., 87, 23274c (1977).
14. A. H. M. Raeymaekers, J. L. H. Van Gelder, G. M. Boeckx, and L. L. Van Hemeldonck, German Offen. 2,813,523; Chem. Abstr., 90, 54975y (1979).
15. H. Loewe, J. Urbanietz, R. Kirsch, and D. Duewel, German Offen. 2,164,690; Chem. Abstr., 79, 92217h (1973).

16. J. L. H. Van Gelder, A. H. M. Raeymaekers, and L. F. C. Roevens, German Offen. 2,029,637; Chem. Abstr., 74, 100047s (1971).
17. J. H. Wikel, C. J. Paget, D. C. DeLong, J. D. Nelson, C. Y. E. Wu, J. W. Paschal, A. Dinner, R. J. Templeton, M. O. Chaney, N. D. Jones, and J. W. Chamberlin, J. Med. Chem., 23, 368 (1980).
18. F. Janssens, M. Luyckx, R. Stokbroekx and J. Torremans, U.S. Patent 4,219,559; Chem. Abstr., 94, 30579d (1981).
19. C. J. Paget and J. L. Sands, German Offen. 2,003,841; Chem. Abstr., 73, 87920c (1970).


11 Benzofused Six-Membered Heterocycles

A diversity of biological effects are possessed by benzofused six-membered heterocycles. These range from antimicrobial activity to cardiovascular, CNS, and inflammation-influencing agents. It can be inferred that the ring system itself is primarily a molecular scaffold upon which to assemble the characteristic pharmacophore for the various receptors involved. It is interesting also to note that the range of bioactivities involved differ substantially from those seen with the benzofused five-membered heterocycles described in Chapter 10.


1. QUINOLINE DERIVATIVES

Various bioisosteric replacements for a phenolic hydroxyl have been explored. One such, a lactam NH, is incorporated into the design of the β -adrenergic blocker, carteolol (3). The fundamental synthon is carbostyryl derivative 1. This is reacted in the usual manner with epichlorohydrin to give 2, which is in turn reacted with t-butylamine to complete the synthesis of carteolol (3), a drug that appears to have relatively reduced nonspecific myocardial depressant action.¹ Carrying this de-

vice farther results in the pseudocatechol, procaterol (6).




Friedel-Crafts alkylation of 8-hydroxycarbofentyls, such as 4, leads to substitution at the C-5 position, namely, 5. In this case an α -haloacyl reagent is employed. Displacement with isopropylamine and careful sodium borohydride reduction (care is




needed to avoid reduction of the carbofentyl double bond) leads to procaterol (6). Procaterol is an adren-ergic agonist selective for β_2 -receptors. Thus it dilates bronchioles without significant cardiac stimulation.²

N-aryl anthranilic acid derivatives ("fenamic acids") often inhibit cyclooxygenase and thereby possess antiinflammatory and analgesic potency. One such agent, floctafenine (9), can also be regarded as a 4-aminoquinoline. The synthesis begins with a Gould-Jacobs reaction of m-trifluoromethylaniline with diethyl methoxymethylene malonate to give (after addition-elimination and thermal cyclization) quinoline 7. Saponification and thermal decarboxylation gets rid of the now surplus carbethoxy group. The phenolic OH is converted to a chloro moiety with phosphorus oxychloride, which is displaced in turn

by methyl anthranilate to give fenamic acid 8. This undergoes ester exchange upon heating in glycerol to complete the synthesis of prodrug floctafenine (9).³




Interest in the antimicrobial properties of quinol-4-one-3-carboxylic acids continues at a significant level. The synthesis of rosoxacin (12) begins with a modified Hantzsch pyridine synthesis employing as component parts ammonium acetate, two equivalents of methyl propiolate, and one of 3-nitrobenzaldehyde. Oxidation of the resulting dihydropyridine (10) with nitric acid followed by saponification, decarboxylation, and reduction of the nitro group with iron and hydrochloric acid gives aniline 11. This undergoes the classic sequence of Gould-Jacobs reaction with methoxymethylenemalonate ester to form the 4-hydroxyquinoline ring, and then alkylation with ethyl iodide and saponification of the ester to complete the synthesis of the antibacterial agent rosoxacin (12).⁴




Droxacin (16) is a carbabioisostere of the clinically useful antimicrobial agent, oxolinic acid. Its synthesis

begins with nitrobenzofuran 13 which is reduced with H_2 and a palladized charcoal catalyst to give aniline 14. Gould-Jacobs reaction with diethyl ethoxymethylenemalonate gives hydroxyquinoline 15 along with some of the alternative cyclization isomer. The synthesis is then completed in the usual way by N-alkylation and saponification to droxacin (16).⁵



An interestingly complex analogue in this family is flumequine (17). As might be expected from the knowledge that the bacterial target is an enzyme (DNATopoisoemrase II), one of the enantiomers is quite potent but the other is not.⁶




Quinfamide (19) is one of a relatively small family of antiamoebic compounds containing a dichloroacetamide function. The synthesis begins by amidation of 6-hydroxytetrahydroquinoline with dichloroacetyl chloride to give 18. The sequence is completed by acylation with 2-furoyl chloride to give quinfamide (19).⁷

2. ISOQUINOLINE DERIVATIVES

Nantradol (25) is an especially interesting agent in that it is a potent analgesic that does not act at the morphine receptors.

It is quickly deacylated in vivo and may qualify as a prodrug. The published synthesis is rather long and bears conceptual similarities to the synthesis of cannabinoids. It has some five asymmetric centers. Dane salt formation between 3,5-dimethoxyaniline and ethyl acetoacetate followed by borohydride reduction gives synthon 20. The amino group is protected by reaction with ethyl chlorocarbonate, the ester group is saponified, and then cyclodehydration with polyphosphoric acid leads to the dihydroquinolone ring system of 21. Deblocking with HBr is followed by etherification of the nonchelated phenolic hydroxyl to give 22. Treatment with sodium hydride and ethyl formate results in both N-formylation and C-formylation of the active methylene to give 23. Michael addition of methyl vinyl ketone is followed by successive base treatments to remove the




activating C-formyl group and then to complete the Robinson annulation to give 24. Lithium in liquid ammonia reduces the olefinic linkage and successive acetylation and sodium borohydride reductions complete the synthesis of nantrado1 (25).⁸

The l-form is much the more potent, being two to seven times more potent than morphine as an analgesic. It is called levonantradol.

3. BENZOPYRAN DERIVATIVES

The enzyme aldose reductase catalyzes the reduction of glucose to sorbitol. Excess sorbitol is believed to contribute to cataracts and to neuropathy by deposition in the lens and nerves of the eyes in the latter stages of diabetes mellitus. Spirohydantoins have been found to inhibit this enzyme and so are of potential value in preventing or delaying this problem. The S enantiomers are the more potent. The synthesis of sorbinil (32) illustrates a method developed for their chiral synthesis. A chiral imine (28) is prepared by titanium tetrachloride-mediated condensation of 6-fluorodihydrobenzopyran-4-one (26) with S- α -methylbenzylamine (27) and this is reacted with hydrogen cyanide to give 29 with a high degree of chirality transfer. The basic nitrogen is next converted to the urea (30) with




highly reactive chlorosulfonyl isocyanate. Treatment with hydrogen chloride results in cyclization to the spirohydantoin


31 whose extraneous atoms are removed by hydrogen bromide treatment to give 4-(S)-sorbinil (32).⁹

Cannabinoids were used in medicine in the form of their crude extracts many centuries ago. Lately the use of cannabis for so-called recreational purposes has become a national vice of substantial proportions. Several attempts have been made to focus the potentially useful pharmacological properties of marijuana into drug molecules with no abuse potential.



Nabilone (37) is a synthetic 9-ketocannabinoid with anti-emetic properties. One of the best of the various published routes to nabilone starts with the enolacetate of nopinone (33), which on short heating with lead tetraacetate undergoes allylic substitution to give 34. Treatment with p-toluene-sulfonic acid in chloroform at room temperature in the presence of the modified olivetol derivative 35 leads to condensation to 36. Finally, treatment with stannic chloride at room temperature opens the cyclobutane ring and allows subsequent phenol capture to give optically active nabilone (37).¹⁰



Nabitan (39) is a cannabis-inspired analgesic whose nitrogen atom was introduced in order to improve water solubility and perhaps to affect the pharmacological profile as well. The phenolic hydroxyl of benzopyran synthon 38 is esterified with 4-(1-piperidino)butyric acid under the influence of dicyclohexylcarbodiimide.¹¹ In addition to being hypotensive and sedative-hypnotic, nabitrane (39) is a more potent analgesic than codeine. The preparation of synthon 38 begins with aceto-




phenone 40, which undergoes a Grignard reaction and subsequent



hydrogenolysis to put the requisite alkyl side chain in place in 41. Ether cleavage (HBr/HOAc) is followed by condensation with piperidone 42 to give tricyclic 43. Reaction with methylmagnesium bromide and hydrogenolysis of the benzylamine linkage followed by alkylation gives 38.¹²

4. BENZODIOXANE DERIVATIVES


In the β -adrenergic blocking drug pyrroxan (48), the catechol moiety is masked in a doxane ring. The synthesis begins by alkylation of phenyl acetonitrile by 2-chloroethanol to produce alcohol 44. Reduction converts this to amino alcohol 45 which undergoes thermal cyclization to 3-phenylpyrrolidine (46).



Finally, a Mannich reaction of 46 with formaldehyde and 4-acetyl-p-benzodioxane (47) leads to pyrroxan (48).¹³

5. BENZOXAZOLINONE DERIVATIVES

One of a variety of syntheses of the antipsychotic agent brofoxine (50) begins with a Grignard reaction on methyl anthranilate. The resulting product (49) is reacted with phosgene in pyridine and the synthesis is completed by bromination in acetic acid to give brofoxine.¹⁴




Another CNS active agent in this structural class is the tranquilizer-antidepressant caroxazone (52). Its published synthesis begins by reductive amination of salicylaldehyde and glycine to give 51. The synthesis is completed by reaction with phosgene and sodium bicarbonate.¹⁵




6. QUINAZOLINONE DERIVATIVES


The clinical acceptance of the dihydrochlorothiazide diuretics led to the synthesis of a quinazolinone bioisostere, fenuizone (54). The synthesis follows the usual pattern of heating anthranilamide (53) with benzaldehyde whereupon aminal formation takes place, presumably via the intermediacy of the Schiff's base.¹⁶




Synthesis of the CNS depressant/tranquilizer tioperidone (59) begins by alkylation of piperazine derivative 55 with 4-chlorobutyronitrile to give 56. Lithium aluminum hydride reduction gives primary amine 57, which is next reacted with isatoic anhydride to give anthranilamide analogue 58. Finally, reaction with phosgene gives tioperidone (59).¹⁷




An apparently unexpected by-product of studies on 1,4-benzodiazepines is the antiinflammatory agent fluquazone (63). The synthesis begins by reaction of typical benzodiazepine synthon 60 with trifluoroacetyl chloride to give intermediate 61. Reaction of this last with ammonium acetate leads to cyclization and cleavage to fluquazone (63). This occurs, presumably, through a variant of a scheme involving facile cleavage of the labile trichloromethyl group, perhaps via 62, followed by cyclodehydration.¹⁸



A related antiinflammatory agent prepared via a more traditional route is fluproquazone (65). Heating with urea in acetic acid results in transamidation by synthon 64 and subsequent cyclodehydration completes the synthesis.¹⁹




An antipsychotic agent with a chemical structure somewhat similar to that of tioperidone (59) is ketanserine (68). The synthesis involves the straightforward thermal alkylation of




N₃-(2-chloroethyl)quinazolidinedione (66) with piperidinylketone 67.²⁰

Alteration of the structural pattern produces a pair of adrenergic α -blocking agents which serve as antihypertensives. These structures are reminiscent of prazosin. Reaction of piperazine with 2-furoyl chloride followed by catalytic reduction of the furan ring leads to synthon 69. This, when heated



in the presence of 2-chloro-4-aminoquinazoline derivative 70, undergoes direct alkylation to terazocin (71).²¹ On the other hand, acylation of quinazoline 72 with oxadiazole derivative 73 gives the antihypertensive tiodazocin (74).²²




7. PHTHALAZINES

Phthalazines commonly possess adrenergic activity. One such, carbazeran (77), is a cardiostonic agent. Its patented synthesis involves nucleophilic aromatic displacement of chlorophthalazine derivative 75 with piperidiny carbamate 76 to give carbazeran (77).²³


8. BENZODIAZAPINES AND RELATED SUBSTANCES

The huge clinical success of drugs in this class has spawned an enormous list of congeners. Synthetic activity has, however, now slowed to the point that a separate chapter dealing with these heterocycles is no longer warranted.

Elfazepam (80) not only is a tranquilizer, but also stimulates feeding in satiated animals. One of several syntheses involves reaction of benzophenone derivative 78 with a glycine equivalent masked as an oxazolidine-2,5-dione (79).²⁴





A water-soluble phosphine derivative of diazepam allows for more convenient parenteral tranquilizer therapy and avoids some complications due to blood pressure lowering caused by the propylene glycol medium otherwise required for administration. Fosazepam (82) is prepared from benzodiazepine 81 by sodium hydride-mediated alkylation with chloromethyldimethylphosphine oxide.²⁵



Lormetazepam (84) is readily synthesized by Polonovski rearrangement of benzodiazepine oxide derivative 83 by heating with acetic anhydride followed by saponification of the resulting rearranged ester.²⁶ The mechanism of this rearrangement to analogous tranquilizers has been discussed previously in this series.²⁷


Quazepam (88) has a highly fluorinated sidechain so as to make this tranquilizer resistant to dealkylation. It also incorporates a lipid-solubilizing 2-thione moiety. The synthesis begins with biarylketone derivative 85 by N-alkylation with 2,2,2-trifluoroethyltrichloroacetate to give 86.




Next the product is acylated with bromoacetyl chloride and the glycine equivalent is constructed in place by a Gabriel amine synthesis (phthalamide anion followed by hydrazine) subsequent to which cyclization to benzodiazepine 87 occurs. The synthesis of the tranquilizer quazepam (88) is finished by thioamide conversion with phosphorus pentasulfide.²⁸

A number of benzodiazepines have heterocyclic rings annelated to them. One such is the tranquilizer midazolam (94). Nitrosation (HONO) of secondary amine 89 leads to the N-nitroso analogue 90. Nitrosoamidines, in the presence of carbanions, undergo carbon-carbon bond formation. Treatment of 90 with nitromethane and potassium *t*-butoxide results in formation of 91. Raney nickel-catalyzed treatment reduces both the double bond and the alkyl nitro group to give saturated amine 92. Treatment with either ethyl orthoacetate or acetic anhydride and polyphosphoric acid results in cyclization to 93 which is converted to the fused imidazole 94, midazolam, on dehydrogenation with manganese dioxide.²⁹

Another alprazolam (95) analogue is adinazolam (98). This




substance is prepared from benzodiazepine synthon 96 by amidation of the hydrazine moiety with chloroacetyl chloride followed by thermal cyclization in acetic acid to 97. Reaction with potassium iodide and diethylamine results in net displacement of the allylic halogen and formation of the tranquilizer and antidepressant, adinazolam (98).³⁰




9. MISCELLANEOUS

The antianginal agent diltiazem (104) is synthesized starting with opening of the epoxide moiety of 99 with the anion of 2-nitrothiophenol to give 100. This is resolved with cinchoni-



dine and reduced to the amine (101) before cyclodehydration to lactam 102. This was alkylated with 2-chloroethyldimethylamine, using dimethylsulfinyl sodium as base, to give 103. The synthesis of the more active d-form of cardioactive diltiazem (104) is concluded by acetylation with acetic anhydride and pyridine.³¹



REFERENCES

1. K. Nakagawa, N. Murakami, S. Yoshizaki, M. Tominaga, H. Mori, Y. Yabuuchi, and S. Shintani, J. Med. Chem., **17**, 529 (1974).
2. S. Yoshizaki, K. Tanimura, S. Tamada, Y. Yabuuchi, and K. Nakagawa, J. Med. Chem., **19**, 1138 (1976).
3. A. Allais, Chim. Ther., **8**, 154 (1973).
4. Y. Lescher and P. M. Carabateas, U.S. Patent 3,907,808 (1975); Chem. Abstr., **84**, 43880p (1975).
5. R. Albrecht, Eur. J. Med. Chem., **12**, 231 (1977); Ann., **762**, 55 (1972).
6. J. F. Gerster, S. R. Rohlfing, and R. M. Winandy, Abstr. N. Am. Med. Chem. Symp. 20-24 June, 1982, p. 153.

7. D. M. Bailey, E. M. Mount, J. Siggins, J. A. Carlson, A. Yarinsky, and R. G. Slighter, J. Med. Chem., 22, 599 (1979).
8. M. R. Johnson and G. M. Milne, J. Heterocycl. Chem., 17, 1817 (1980).
9. R. Sarges, H. R. Howard, Jr., and P. R. Kelbaugh, J. Org. Chem., 47, 4081 (1982).
10. R. A. Archer, W. B. Blanchard, W. A. Day, D. W. Johnson, E. R. Lavagnino, C. W. Ryan, and J. E. Baldwin, J. Org. Chem., 42, 2277 (1977).
11. R. K. Razdan, B. Z. Terris, H. K. Pars, N. P. Plotnikoff, P. W. Dodge, A. T. Dren, J. Kyncl, and P. Somani, J. Med. Chem., 19, 454 (1976).
12. M. Winn, D. Arendsen, P. Dodge, A. Dren, D. Dunnigan, R. Hallas, H. Hwang, J. Kyncl, Y.-H. Lee, N. Plotnikoff, P. Young, H. Zaugg, H. Dalzell, and R. K. Razdan, J. Med. Chem., 19, 461 (1976).
13. V. A. Dobrina, D. V. Ioffe, S. G. Kuznetsov, and A. G. Chigarev, Khim. Pharm. Zh., 8, 14 (1974); Chem. Abstr., 81, 91445k (1974).
14. L. Bernardi, S. Coda, A. Bonsignori, L. Pegrassi, and G. K. Suchowsky, Experientia, 24, 774 (1968).
15. L. Bernardi, S. Coda, V. Nicoletta, G. P. Vicario, A. Minghetti, A. Vigevari, and F. Arcamone, Arzneim. Forsch., 29, 1412 (1979); L. Bernardi, S. Coda, L. Pegrassi, and G. K. Suchowsky, Experientia, 24, 74 (1968).
16. G. Cantarelli, Il Farmaco, Sci. Ed., 25, 761 (1970).
17. R. F. Parcell, U.S. Patent 3,819,630; Chem. Abstr., 80, 146190k (1974).
18. L. Bernardi, S. Coda, V. Nicoletta, G. P. Vicario, A. Minghetti, A. Vigevari, and F. Arcamone, Arzneim. Forsch.,

- 29, 1412 (1979); L. Bernardi, S. Coda, L. Pegrassi, and K. G. Suchowsky, Experientia, 24, 774 (1968).
19. P. G. Mattner, W. G. Salmond, and M. Denzer, French Patent 2,174,828 (1973).
20. J. Vandenberg, L. E. J. Kennis, M. J. M. C. Van der Aa, and A. H. M. T. van Heertum, Eur. Patent Appl. 13,612 (1980); Chem. Abstr., 94; 65718a (1981).
21. M. Winn, J. Kyncl, D. A. Dunnigan, and P. H. Jones, U.S. Patent 4,026,894 (1977); Chem. Abstr., 87; 68411m (1977).
22. R. A. Partyka and R. R. Crenshaw, U.S. Patent 4,001,237 (1977); Chem. Abstr., 86; 140028r (1977).
23. S. F. Campbell, J. C. Danilewicz, A. G. Evans, and A. L. Ham, British Patent Appl. GB 2,006,136 (1979); Chem. Abstr., 91; 193331u (1979).
24. Anon., Japanese Patent JP 709,691 (1970); Chem. Abstr., 80; 133497r (1974).
25. E. Wolfe, H. Kohl, and G. Haertfelder, German Patent DE 2,022,503 (1971); Chem. Abstr., 76; 72570c (1972).
26. S. C. Bell, R. J. McCaully, C. Gochman, S. J. Childress, and M. I. Gluckman, J. Med. Chem., 11, 457 (1968); S. C. Bell and J. C. Childress, J. Org. Chem., 27, 1691 (1962).
27. D. Lednicher and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Wiley, New York, Vol. 1, p. 365; Vol. 2, p. 402.
28. M. Steinman, J. G. Topliss, R. Alekel, Y.-S. Wong, and E. E. York, J. Med. Chem., 16, 1354 (1973).
29. A. Walser, L. E. Benjamin, Sr., T. Flynn, C. Mason, R. Schwartz, and R. I. Fryer, J. Org. Chem., 43, 936 (1978); R. I. Fryer, J. Blount, E. Reeder, E. J. Trybulski, and A. Walser, J. Org. Chem., 43, 4480 (1978).

30. J. B. Hester, Jr., A. D. Rudzik, and P. F. Von Voigtlander, J. Med. Chem., 23, 392 (1980).
31. H. Inoue, S. Takeo, M. Kawazu, and H. Kugita, Yakugaku Zasshi, 93, 729 (1973); H. Kugita, H. Inoue, M. Ikezaki, M. Konda, and S. Takeo, Chem. Pharm. Bull., 19, 595 (1971).


12 Beta-Lactams

After 40 years of clinical use, benzylpenicillin (1) remains an extremely effective and useful drug for the treatment of infections caused by bacteria susceptible to it. It fails, however, to be a perfect drug on several grounds. Its activity spectrum is relatively narrow; it is acid and base unstable and so must be given by injection; increasingly strains carry enzymes (β -lactamases) that inactivate the drug by hydrolysis; and it is haptenic so that many patients become allergic to it. Many analogues have been synthesized in order to overcome these drawbacks and a substantial number of semisynthetic penicillins, cephalosporins, cephamycins, and so forth, have subsequently been marketed. Recently new impetus has been added to the field by the discovery of new ring systems in fermentation liquors and through development of novel synthetic approaches so that the field of β -lactam chemistry is now characterized by the feverish activity reflected in the number of entries in this chapter.

1. PENICILLINS

One of the most successful penicillin analogues has been


ampicillin (2). The relatively small chemical difference between ampicillin and benzylpenicillin not only allows for substantial oral activity but also results in a substantial broadening of antimicrobial spectrum so as to allow for use against many Gram-negative bacteria. Many devices have been employed in order to enhance still further the oral absorption of ampicillin. Bacampicillin (6) is a prodrug of ampicillin



designed for this purpose. An azidopenicillin sodium salt (3) is reacted with mixed carbonate ester 4 (itself prepared from acetaldehyde and ethyl chlorocarbonate) to give ester 5. Reduction of the azido linkage with hydrogen and a suitable catalyst produces bacampicillin (6). Both enantiomers [starred (*) carbon] are active. The drug is rapidly and efficiently absorbed from the gastrointestinal tract and is quickly cleaved by serum esterases to bioactive ampicillin (2), acetaldehyde, carbon dioxide, and ethanol.¹

Sarpicillin (10) is a double prodrug of ampicillin in that not only is the carboxy group masked as an ester, but a

hetacillin-like acetonide has been added to the C-6 amide side chain. Its synthesis begins with the potassium salt of




penicillin V (phenoxymethylpenicillin, 7) which is esterified with methoxymethyl chloride to give 8. This is reacted with phosphorus pentachloride and the resulting imino chloride is cleaved to the free amine with N,N-dimethylaniline. Amine 9 then reacts with phenylglycyl chloride in acetone to complete the synthesis of sarpacillin (10).² The acetonide is presumably formed after acylation. A closely related prodrug of amoxycillin, known as sarmoxicillin (11), is made in the same way but with reaction of amine 9 with the hydrochloride of p-hydroxyphenylglycyl chloride in acetone being involved instead.³


An important molecular target of the β -lactam antibiotics is an enzyme that acts as a transpeptidase in the stepwise polymerization leading to a thickened, strong bacterial cell wall. Several amino acids are present in addition to the terminal D-alanyl-D-alanyl unit which the Strominger hypothesis suggests has the same overall shape and reactivity as ampicillin. This suggests that acylation of the amino group of ampicillin might lead to enhanced affinity or at least would be sterically allowable. It is interesting to find, therefore, that such acylation broadens the antimicrobial spectrum of the corresponding penicillins so that they now include the important

Gram-negative pathogen *Pseudomonas aeruginosa*.

Mezlocillin (13), one such agent, can be made in a variety of ways including reaction of ampicillin with chlorocarbamate 12 in the presence of triethylamine.⁴ Chlorocarbamate 12




itself is made from ethylenediamine by reaction with phosgene to form the cyclic urea followed by monoamide formation with methanesulfonyl chloride and then reaction of the other nitrogen atom with phosgene and trimethylsilyl chloride. The closely related analogue azlocillin (14) is made in essentially the same manner as for mezlocillin, but with omission of the mesylation step.⁵ Interestingly, azlocillin is the more active of the two against many *Pseudomonas aeruginosa* strains in vitro. An interesting alternative synthesis of azlocillin involves activation of the substituted phenylglycine analogue 15 with 1,3-dimethyl-2-chloro-1-imidazolinium chloride (16) and then condensation with 6-aminopenicillanic acid.^{5,6}




Another acylated ampicillin derivative with expanded antimicrobial spectrum is piperacillin (19). Its synthesis begins with 1-ethyl-2,3-diketopiperazine (17, which itself is made from N-ethylethylenediamine and diethyl oxalate), which is activated by sequential reaction with trimethylchlorosilane and then trichloromethyl chloroformate to give 18. This last reacts with ampicillin (2) to give piperacillin (19) which is active against, among others, the Enterobacteriaceae and Pseudomonads that normally are not sensitive to ampicillin.⁷

Continuing this theme, pirbenicillin (22) is another N-acylated antipseudomonal ampicillin analogue. Its synthesis begins by acylating 6-aminopenicillanic acid with N-carbobenzoxypyrenylglycine by reaction with dicyclohexylcarbodiimide and N-hydroxysuccinimide to activate the carboxyl group. The protecting CBZ group is removed from 19 on treatment with sodium carbonate to give 20. The synthesis of pirbenicillin is completed by reaction with 4-pyridoisminomethyl ether (21) (itself prepared from 4-cyanopyridine and anhydrous methanolic hydrogen chloride).⁸



Piridicillin (27) is made by N-acylating amoxycillin with a rather complex acid. The synthesis begins by reacting N,N-diethanolamine with p-acetylbenzene-sulfonyl chloride to give 23. Conversion (to 24) with ethyl formate and sodium


methoxide is followed by base-catalyzed addition-elimination with cyanoacetamide, during the course of which reaction cyclodehydration occurs to produce the pyridone (25). Saponification of the carboxylic acid 26 is followed by carboxy



activation using the active ester method (dicyclohexylcarbodiimide and N-hydroxysuccinimide) and condensation with amoxycillin to produce the broad spectrum antibiotic, piridicillin (27).⁹

There is only one clinically significant penicillin at present that does not have an amide side chain. Mecillinam (amidinocillin, 29) has, instead, an amidine for a side chain. It has very little effective anti-Gram positive activity but it is quite effective against Gram-negative microorganisms. Its synthesis begins by reacting N-formyl-1-azacycloheptane with oxalyl chloride to form the corresponding imino chloride (28). This is then reacted with 6-aminopenicillanic acid to produce mecillinam.¹⁰ A prodrug form, amidinocillin pivoxyl (30), is

made in the same manner by reaction of 28 with pivaloyloxymethyl 6-aminopenicillanoate instead.¹¹




2. CEPHALOSPORINS

Widespread clinical acceptance continues to be accorded to the cephalosporins, and the field is extremely active as firms search for the ultimate contender. Among the characteristics desired is retention of the useful features of the older members (relatively broad spectrum, less antigenicity than the penicillins, relative insensitivity toward β -lactamases, and convenience of administration) while adding better oral activity and broader antimicrobial activity (particularly potency against Pseudomonas, anaerobes, meningococci, cephalosporinase-carrying organisms, and the like). To a considerable extent these objectives have been met, but the price to the patient has been dramatically increased.


Cephachlor (35) became accessible when methods for the preparation of C-3 methylenecephalosporins became convenient. The allylic C-3-acetoxy residue characteristic of the natural cephalosporins is activated toward displacement by a number of oxygen- and sulfur-containing nucleophiles. Molecules such as 31 can therefore be prepared readily. Subsequent reduction with chromium(II) salts leads to the desired C-3 methylenecephems (32), which can in turn be ozonized at low temperatures to produce the C-3 keto analogues. These are

isolated in the form of the C-3 hydroxycephem enolates (33). Next, treatment with a variety of chlorinating agents (SOCl_2 , PCl_3 , POCl_3 , $(\text{COCl})_2$, and COCl_2) in dry DMF solvent produces the C-3 chloro analogues (34). The reaction can be carried out so that the C-7 side chain is removed by the imino chloride method so as to allow installation of the 7-D-2-amino-2-phenylacetamido side chain of cephaclor (35).¹²




Conceptually closely related, cefroxadine (40) can be prepared by several routes, including one in which the enol (33) is methylated with diazomethane as a key step. A rather more involved route starts with comparatively readily available phenoxymethylpenicillin sulfoxide benzhydryl ester (36). This undergoes fragmentation when treated with benzothiazole-2-thiol to give 37. Ozonolysis (reductive work-up) cleaves the olefinic linkage and the unsymmetrical disulfide moiety is converted to a tosyl thioester (38). The enol moiety is methylated with diazomethane, the six-membered ring is closed by reaction with 1,5-diazabicyclo[5.4.0]undec-5-ene (DBU), and the ester protection is removed with trifluoroacetic acid to


give 39. The amide side chain is removed by the usual phosphorus pentachloride/dimethylaniline sequence followed by reamidation with the appropriate acid chloride. The result of all this is cefroxadine (40).¹³



Possessing a side chain at C-7 reminiscent of that of amoxicillin and a more typical sulfur containing C-3 moiety, cefatrizine (44) can be synthesized by the active ester mediated condensation of t-BOC-2-p-hydroxyphenylglycine (41) with N-silyl 3-cephem synthon 42. The t-butyloxycarbonyl protecting group of intermediate 43 is removed with formic acid in order to complete the synthesis of cefatrizine (44), a broad spectrum cephalosporin.¹⁴ The thiol necessary for synthesis of intermediate 42 from 7-aminocephalosporanic acid can be prepared from 1-N-benzyl-2-azoimidazole (45) by lithiation (n-butyllithium) followed by thiolation (hydrogen sulfide) to give intermediate 46. The protecting benzyl moiety is then removed reductively with sodium in liquid ammonia to give synthon 47.




Structurally rather similar to cefatrizine is cefaparole (49). It is prepared in quite an analogous manner by active ester condensation of 41 and 7-aminocephalosporanic acid analogue 48. The blocking group is removed with trifluoroacetic acid in anisole (9:1) to give cefaparole. Racemization does not take place during the synthetic sequence so the desired R stereochemistry of the side chain amino group is retained.¹⁴




Analogous to azlocillin-mezlocillin, acylation of the amino group of 2-phenylglycine containing cephalosporins is consistent with antipseudomonal activity. There are many


routes to cefoperazone (52). One of the more obvious is the condensation of cephalosporin antibiotic 50 with 2,3-diketopiperazine 51 under modified Schotten-Baumann conditions.¹⁵



Cefonicid (55) is synthesized conveniently by nucleophilic displacement of the C-3 acetoxy moiety of 53 with the appropriately substituted tetrazole thiol (54).¹⁶ The mandelic acid amide C-7 side chain is reminiscent of cefamandole.




Cefazaflur (58) stands out among this group of analogues because it lacks an arylamide C-7 side chain (see cephacetrile for another example).¹⁷ Cefazaflur (58) is synthesized by reaction of 3-(1-methyl-1H-tetrazol-5-ylthiomethylene)-7-aminocephem-4-carboxylic acid (56) with trifluoromethylthioacetyl chloride (57).¹⁸




Cefsulodin (60) has a sulfonic acid moiety on the C-7 acyl side chain. This moiety conveys antipseudomonal activity to certain penicillins, and it is interesting to note that this artifice works with cefsulodin as well. It is also interesting that, in contrast to the other so-called third-generation cephalosporins, the spectrum of cefsulodin is rather narrow and its clinical success will place a premium upon accurate diagnosis. Its synthesis begins by acylation of 7-aminocephalosporanic acid with the acid chloride of 2-sulfonylphenylacetyl chloride to give cephalosporin 59. Reaction of that intermediate with aqueous potassium iodide and isonicotinic acid amide results in acetoxyl displacement from C-3 and formation of cefsulodin (60).²¹ The quaternary base at C-3 is reminiscent of the substitution pattern of cephaloridine.

The structural feature of ceforanide that is of particular interest is the movement of the C-7 side chain amino moiety from the position α to the amide carbonyl, where it normally resides in ampicillin/cephalexin analogues, to lodgement on a methylene attached to the ortho position on the aromatic ring.




This puts it geometrically in the same general area but considerably alters the electronic character of the molecule upon protonation. The synthesis of ceforanide (64) begins with a Beckmann rearrangement of the oxime of 2-indanone (61) to give lactam 62. Hydrolysis followed by protection of the amino group as the enamine (63) allows for subsequent mixed anhydride (isobutylchlorocarbonate)-mediated amide formation with the corresponding 7-aminocephalosporin synthon to give ceforanide (64). The requisite nucleophile for the C-3 moiety is prepared simply by carbonation of the lithio derivative of 1-methyl-1-H-tetrazol-5-ylthiol.²²

Cefotiam (67) has an acyl aromatic C-7 side chain bioisosteric with an anilino ring. It can be prepared by acylation of the suitable acid moiety with 4-chloroacetoacetyl





chloride to give amide 65. Chloride displacement with thiourea leads to cyclodehydration to the aminothiazole cefotiam (67), probably via intermediate 66.²³

The interposition of a syn-oximino ether moiety between the amide carbonyl and the aromatic ring has proved richly





rewarding in that substantial resistance to β -lactamases results from this steric hindrance. A large number of analogues now bear this feature, for example, cefuroxime (71). The synthesis can be accomplished in a variety of ways. Benzhydryl ester 68 (preparable from cephalothin²⁴) is acylated with trichloroacetyl isocyanate and the side chain at C-7 is removed via the imino chloride method (pyridine and phosphorus pentachloride followed by tosic acid) to produce 7-aminocephalosporanic acid analogue 69. The carbamate moiety is partially hydrolyzed to give 70 through use of anhydrous methanolic hydrogen chloride (generated with methanol and acetyl chloride). Next, the benzhydryl ester is cleaved with trifluoroacetic acid and the synthesis is concluded by a Schotten-Baumann acylation with the appropriate syn-oximinoether-bearing acid chloride so as to produce cefuroxime.²⁵

An analogous third-generation cephalosporin whose synthesis illustrates one of the methods of preparing the requisite oximino acid side chains is cefotaxime (76). syn-Oxime 72 is methylated stepwise with dimethyl sulfoxide and base and then chlorinated (Cl_2 in chloroform) to produce 73. Alkylation of thiourea with this product results in concomitant cyclodehydration to produce aminothiazole 74. The primary amino group is then blocked with chloroacetyl chloride, the ester group is saponified, and then the intermediate is used to acylate 7-aminocephalosporanic acid in the usual way via the acid chloride. The blocking chloroacetamide moiety of 75 is then cleverly removed by reaction with thiourea in order to unmask β -lactamase stable cefotaxime (76).²⁶ Ceftazidime (81) carries the theme of bulky oximino ethers much further. Its synthesis begins with nitrous acid treatment of ethyl acetoacetate to produce oxime 77. This is next converted to




a 2-aminothiazole (78) by halogenation with sulfonyl chloride followed by thiourea displacement. The amino group is



Pieces of various routes to moxalactam have been published from which the following may be assembled as one of the plausible pathways. The benzhydryl ester of 6-aminopenicillanic acid (84) is S-chlorinated and treated with base whereupon the intermediate sulfonyl chloride fragments (to 85). Next, displacement with propargyl alcohol in the presence of zinc chloride gives predominantly the stereochemistry represented by diastereoisomer 86. The side chain is protected as the phenyl-acetylamide; the triple bond is partially reduced with a 5% Pd-CaCO₃ catalyst and then epoxidized with m-chloroperbenzoic acid to give 87. The epoxide is opened at the least hindered end with the lithium salt of 1-methyl-1H-tetrazol-5-ylthiol to put in place the future C-3 side chain and give intermediate 88. Jones oxidation followed in turn by ozonolysis (reductive work-up with zinc-acetic acid) and reaction with thionyl chloride and pyridine give halide 89. The stage is now set for an intramolecular Wittig reaction. Displacement with triphenylphosphine and Wittig olefination gives 1-oxacephem 90. Next a sequence is undertaken of side chain exchange and introduction of a C-7 methoxyl group analogous to that which is present in the cephamycins and gives them enhanced β -lactamase stability. First 90 is converted to the imino chloride with PCl₅ and then to the imino methyl ether (with methanol) and next hydrolyzed to the free amine. Imine formation with 3,5-di-t-butyl-4-hydroxybenzaldehyde is next carried out leading to 91. Oxidation with nickel peroxide gives iminoquinone methide 92, to which methanol is added in a conjugate sense and in the stereochemistry illustrated in formula 93. The imine is exchanged away with Girard reagent T to give 94, and this is acylated by a suitable protected arylmalonate, as the hemiester hemiacid chloride, so as to give 95. Deblocking with aluminum chloride and anisole gives moxalactam (96).

Moxalactam is a synthetic antibiotic with good activity against Gram-negative bacteria including pseudomonads and has excellent stability against β -lactamases.²⁷



REFERENCES

1. B. A. Ekstrom, O. K. J. Kovacs, and B. O. H. Sjoberg, German Offen. 2,311,328 (1973); Chem. Abstr., 80, 14921q (1974).
2. P. Sleezer and D. A. Johnson, German Offen. 2,244,915 (1973); Chem. Abstr., 78, 159590z (1973).
3. P. Sleezer and D. A. Johnson, South African 75 04,722 (1976); Chem. Abstr., 86, 171440y (1977).
4. W. Schroeck, H. R. Furtwaengler, H. B. Koenig, and K. G. Metzger, German Offen. 2,318,955 (1973); Chem. Abstr., 82, 31313b (1975).
5. H. B. Koenig, K. G. Metzger, H. A. Offe, and W. Schroeck, Eur. J. Med. Chem., 17, 59 (1982).
6. V. J. Bauer and S. R. Safir, J. Med. Chem., 9, 980 (1966).
7. I. Saikawa, S. Takano, C. Yoshida, O. Takashima, K. Momonoi, T. Yasuda, K. Kasuya, and M. Komatsu, Yakugaku Zasshi, 97, 980 (1977).
8. E. S. Hamanaka and J. G. Stam, South African 74 00,509 (1973); Chem. Abstr., 83, 58808z (1975).
9. J. S. Kaltenbronn, J. H. Haskell, L. Daub, J. Knoble, D. DeJohn, U. Krolls, N. Jenesel, G.-G. Huang, C. L. Heifitz, and M. W. Fischer, J. Antibiotics, 32, 621 (1978).
10. F. Lund and L. Tybring, Nature, New Biol., 236, 135 (1972).
11. F. J. Lund, German Offen. 2,055,531 (1971); Chem. Abstr., 75, 49070k (1971).
12. R. R. Chauvette and P. A. Pennington, J. Med. Chem., 18, 403 (1975).
13. R. B. Woodward and H. Bickel, U.S. Patent 4,147,864 (1979); Chem. Abstr., 91, 74633j (1979).
14. G. L. Dunn, J. R. E. Hoover, D. A. Berges, J. J. Taggart,

- L. D. Davis, E. M. Dietz, D. R. Jakas, N. Yim, P. Actor, J. V. Uri and, J. A. Weisbach, J. Antibiotics, 29, 65 (1976).
15. I. Saikawa, S. Takano, Y. Shuntaro, C. Yoshida, O. Takashima, K. Momonoi, S. Kuroda, M. Komatsu, T. Yasuda, and Y. Kodama, German Offen., DE 2,600,880 (1977); Chem. Abstr., 87, 184533b (1977).
16. D. A. Berges, U.S. Patent 4,093,723 (1978); Chem. Abstr., 89, 180025f (1978).
17. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Wiley, New York, 1980, Vol. 2, p. 441.
18. R. M. deMarinis, J. C. Boehm, G. L. Dunn, J. R. E. Hoover, J. V. Uri, J. R. Guarini, L. Philips, P. Actor, and J. A. Weisbach, J. Med. Chem., 20, 30 (1977).
19. H. Nomura, T. Fugono, T. Hitaka, I. Minami, T. Azuma, S. Morimoto, and T. Masuda, Heterocycles, 2, 67 (1974).
20. W. J. Gottstein, M. A. Kaplan, J. A. Cooper, V. H. Silver, S. J. Nachfolger, and A. P. Granatek, J. Antibiotics, 29, 1226 (1979).
21. M. Namata, I. Minamida, M. Yamaoka, M. Shiraishi, T. Miyawaki, H. Akimoto, K. Naito, and M. Kida, J. Antibiotics, 31, 1262 (1978).
22. D. Lednicer and L. A. Mitscher, The Organic Chemistry of Drug Synthesis, Wiley, New York, 1977, Vol. 1, p. 417, 420.
23. M. C. Cook, G. I. Gregory, and J. Bradshaw, German Offen., DE2,439,880 (1975); Chem. Abstr., 83, 43354z (1975).
24. M. Ochiai, A. Morimoto, T. Miyawaki, Y. Matsushita, T. Okada, H. Natsugari, and M. Kida, J. Antibiotics, 34, 171 (1981); R. Reiner, U. Weiss, U. Brombacher, P. Lanz, M. Montavon, A. Furlenmeier, P. Angehrn, and P. J. Probst, J. Antibiotics, 33, 783 (1980).




25. C. H. O'Callaghan, D. G. H. Livermore, and C. E. Newall, German Offen. DE 2,921,316 (1979); Chem. Abstr., 92, 198413c (1980).
26. T. Takaya, H. Takasugi, K. Tsuji, and T. Chiba, German Offen. DE 2,810,922 (1978); Chem. Abstr., 90, 204116k (1979).
27. H. Otsuka, W. Nagata, M. Yoshioka, M. Narisada, T. Yoshida, Y. Harada, and H. Yamada, Med. Res. Revs., 1, 217 (1981); M. Narisada, H. Onoue, and W. Nagata, Heterocycles, 7, 839 (1977); M. Narisada, T. Yoshida, O. Onoue, M. Ohtani, T. Okada, T. Tsugi, I. Kikkawa, N. Haga, H. Satoh, H. Itani, and W. Nagata, J. Med. Chem., 22, 757 (1979).

13 Miscellaneous Fused Heterocycles


Medicinal agents discussed to this point have been roughly classifiable into some common structural groups; biological activity often followed the same rough classification. As was the case in the preceding volumes in this series, a sizable number of compounds, often based on interesting heterocyclic systems, defy ready grouping by structure. These are thus discussed below under the cover of "miscellaneous." It might be added as an aside that this section may include compounds that will someday move to new chapters. If one of these drugs proves to be a major clinical or marketing success, it will no doubt occasion a considerable amount of competitive work. Since some of this work will undoubtedly result in agents with generic names, the class may well finally grow to the point where it will

require listing as a structural group.


A rather simple derivative of imidazoimidazoline has been described as an antidepressant agent. Preparation of this compound starts with the displacement of the nitramine grouping in imidazoline derivative 1 by phenylethanolamine 2. The product of this reaction is then treated with thionyl chloride. The probable chloro intermediate (4) cyclizes under the reaction conditions to afford imafen (5).¹



The imidazothiazoline tetramisole (6) has shown quite good activity as a broad spectrum anthelmintic agent. This drug has in addition aroused considerable interest as an agent which modifies the host immune response. Further substitution on the aromatic ring has proved compatible with activity. Displacement of halogen on the phenacyl bromide 7 with aminothiazole 8 affords the alkylated product 9. Catalytic hydrogenation serves to reduce both the heterocyclic ring and the carbonyl group (10). Cyclization by means of sulfuric acid completes the synthesis of butamisole (11).²




Benzofurans of the very general structure represented by 12 have formed the basis of several quite effective drugs for treatment of cardiovascular disease. It is thus of note that replacement of the aromatic nucleus by the isosteric indolizidine system affords a compound with quite similar activity. Friedel-Crafts type acylation of indolizidine 13 with substituted benzoyl chloride 14 gives the ketone (15). Removal of the protecting group gives the free phenol. Alkylation by means of *N,N*-di(*n*-butyl)-2-chloroethylamine affords the corresponding basic ether. There is thus obtained the antiarrhythmic agent butoprizine (17).³



Aggregation of blood platelets is the requisite first event for the maintenance of intact circulation in the face of any break in a blood vessel. It is the platelet clump that starts the long and complicated process leading to closure of the broken vessel by an organized blood clot. Though this property of platelets is vital to maintenance of the circulatory system, an excessive tendency to aggregation can also lead to problems. Thus platelet clumps formed in blood vessels in the absence of injury can lead to blockade of blood circulation and subsequent injury. Strokes and some types of myocardial infarcts have thus been associated with platelet clumps. The nonsteroid antiinflammatory agents as a class show platelet antiaggregation activity in a number of test systems; however, there has been a considerable amount of effort expended to uncovering agents from other structural classes that will not share the deficits of the nonsteroid antiinflammatories. Ticlopidine (24), a drug that shows good activity in various animal models has undergone extensive clinical testing as a platelet antiaggregator.


The key intermediate 21 is in principle accessible in any of several ways. Thus reaction of thiophenecarboxaldehyde 18 with aminoacetal 19 would lead to the Schiff base 20; treatment with acid would result in formation of the fused thiophene-pyridine ring (21). Alkylation of that intermediate with benzyl chloride 22 gives the corresponding ternary iminium salt 23. Treatment with sodium borohydride leads to reduction of the quinolinium ring and thus formation of ticlopidine (24).⁴




The purines, as is well known, play a very central role in the biochemistry of life. This heterocyclic nucleus is involved in vital processes in a host of guises, from its participation in the genetic message to its part in the energy transmission system and perhaps even as a neurotransmitter. It is thus not surprising that considerable attention has been devoted to this heterocyclic system as a source for drugs; it is somewhat unexpected that so few of these efforts have met with success.


The success of antibacterial therapy hinges largely on the fact that the metabolism of bacteria differs sufficiently from that of the host so that it is possible to interfere selectively with this process. Viral infections have been much more difficult to treat because the organism in effect takes over the metabolic processes of the host cell; selectivity is thus very slight. One of the signal breakthroughs in this field of therapy is an agent that takes advantage of one of those small differences, acting as a false substrate for a biochemical process necessary for viral replication. It is pertinent that this drug, acyclovir (27) may be viewed as an analogue of the nucleoside guanosine 28, in which two of the ring carbons of ribose (or deoxyribose) have been deleted. Preparation of

this agent starts with the alkylation of guanine (25) with the chloromethyl ether 25a. Removal of the protecting group (26) by saponification affords acyclovir (27).⁵




The uric acid derivative theophylline (29) is one of the mainstays as a bronchodilator drug for the treatment of asthma. This agent's narrow therapeutic index and host of side effects has led to an active search for a safer derivative. Synthesis of one such compound starts with the condensation of amine 30 with methyl isocyanate. Acylation of the resulting urea (31) with cyanoacetic acid gives the intermediate 32; this is then cyclized to the corresponding uracil 33 by means of base. Nitrosation (34) followed by reduction of the newly introduced nitroso group gives the ortho diamine function (35). The remaining ring is constructed by first acylating 35 with acetic anhydride (to give 36); cyclization again by means of base completes the purine nucleus. There is thus obtained the bronchodilating agent verofylline (37).⁶







Beta adrenergic agonists also exert bronchodilating effects. These drugs are thus often used in conjunction with theophylline in asthma therapy. A drug that combines both moieties, reproterol (40), has interestingly proved clinically useful as an antiasthmatic agent. This compound can in principle be obtained by first alkylating theophylline with 1-bromo-3-chloropropane to give 38. Use of this halide to alkylate aminoalcohol 39 would then afford reproterol (40).



As noted earlier (see Chapter 10), 4-acylpiperidines separated from benzimidazole by a three carbon chain often show antipsychotic activity. The heterocycle can apparently be replaced by a pyridopyrimidine ring. Thus alkylation of piperidine 41 with halide 42 affords pirenperone (43).⁷




Hydrazinopyridazines such as hydralazine have a venerable history as antihypertensive agents. It is of note that this biological activity is maintained in the face of major modifications in the heterocyclic nucleus. The key intermediate keto ester 45 in principle can be obtained by alkylation of the anion of piperidone 44 with ethyl bromoacetate. The cyclic acylhydrazone formed on reaction with hydrazine (46) is then oxidized to give the aromatized compound 47. The hydroxyl group is then transformed to chloro by treatment with phosphorus oxychloride (48). Displacement of halogen with hydrazine leads to the formation of endralazine (49).⁸



Two closely related pyridotriazines have been described as antifungal agents. Displacement of halogen on nitro-


chloropyridine 50 with the monocarbamate of hydrazine affords intermediate 51. This is then first hydrolyzed to the free hydrazine (52) and the nitro group reduced to the corresponding amine (53). Condensation of this intermediate with phenylacetic acid leads to formation of the cyclic amidine derivative 54. Oxidation with manganese dioxide introduces the remaining unsaturation; there is thus obtained triafungin (55).⁹ Condensation of 53 with phenoxyacetic acid gives, after aromatization of the first formed product, the antifungal agent oxyfungin (56).¹⁰




The enormous commercial success of the benzodiazepine anxiolytic agents has spurred a correspondingly large effort in many laboratories aimed at developing novel analogues (see, for example, Chapter 11). In this case it is probably no exaggeration to say that every part of the parent molecule has been modified in the search for novel patentable analogues. In the course of such work it has been found that replacement of the fused benzene ring by a

heterocyclic ring is compatible with tranquilizing activity.


Preparation of one of the analogues in which benzene is replaced by pyrazole starts by nitration of pyrazole carboxylic acid 57. The product, 58, is then converted to the acid chloride (59). This intermediate is then used to acylate benzene in a Friedel-Crafts reaction. The nitroketone is then reduced to the corresponding amine. Reaction with ethyl glycine can be visualized as involving initially formation of the Schiffs' base (62). Displacement of ethoxide by the ring amino group leads to formation of the lactam. There is thus obtained ripazepam (63).¹¹



A somewhat different strategy is employed for preparation of the desoxy analogue containing the reversed pyrazole. Acylation of chloropyrazole 64 with *m*-chlorobenzoyl chloride affords the ketone 65. Reaction of that with ethylenediamine leads directly to the anxiolytic agent zometapine (66).¹² The overall sequence obviously involves sequential Schiff base formation and nucleophilic displacement of chlorine; the order of these steps is not clear.




In a similar vein, acylation of aminoketone 67 with chloroacetyl chloride affords the corresponding chloroamide 68. Reaction of that intermediate with ammonia serves to form the diazepine ring, possibly via the glycineamide. The product bentazepam (69) is described as a tranquilizer.¹³





Carboxylic acid derivatives of heterocycles have proved a source of compounds that show the same allergic mediator release inhibiting activity as sodium cromoglycate. A number of these agents have been taken to the clinic for trial as antiallergic agents.

Friedel-Crafts cyclization of phenoxy ether 70 leads to the corresponding xanthone 71. Exhaustive oxidation of the methyl group leads to the carboxylic acid, xanoxate (72).¹⁴




Preparation of the analogue in which isopropoxy is replaced by a methylsulfoxide involves a somewhat more complex scheme. Aromatic nucleophilic displacement of halogen in dicarboxyl ester 73 leads to diphenyl ether 75. The product is then saponified (76), cyclized to the xanthone and again esterified (78). The aromatic ether is then demethylated to the free phenol (79). This group is converted to the thiocarbamate 80 by means of dimethylthiocarbamoyl chloride. Thermal rearrangement of the thiocarbamate function by the method of Newman results in overall exchange of sulfur for oxygen to afford thiocarbamate 81. This is then converted to the free thiol, with accompanying saponification (82). Methylation of the thiol group (83) followed by controlled oxidation of the thioether leads to the sulfoxide. There is thus obtained the antiallergic agent tixanox (84).¹⁵






It has by now been well established that the tricyclic ring system of the phenothiazine tranquilizers is not an absolute requirement for antipsychotic activity; that moiety has been successfully replaced by ring systems as diverse as acridine and even dihydroanthracene. It should thus not be surprising to note that dibenzopyran derivatives also lead to active compounds. Thus reaction of xanthone 85 with the Grignard reagent from chloropiperidine 86 gives after dehydration the antipsychotic agent clonipazam (87).¹⁶




It is by now apparent that the nature of the aryl group in the arylacetic and arylpropionic acid antiinflammatory agents can be varied quite widely without loss of activity. The corresponding derivatives of homologous xanthenes and thioxanthenes thus both show activity as nonsteroid antiinflammatory agents.


Starting material for the first of these agents can in principle be obtained by alkylation of phenol 88 with benzyl chloride 89. Cyclization of the product (90) under Friedel-Crafts conditions leads directly to isoxepac (91).¹⁷




Preparation of the sulfur analogue involves as the first step cyclization of the terephthalic acid derivative 92. The acid is then converted to the acid chloride and this is allowed to react with diazomethane. Rearrangement of the resulting diazoketone (95) under the conditions of the Arndt-Eistert reaction leads to the homologated acid. There is thus obtained tiopinac (96).¹⁸



Antidepressant agents show almost the same degree of tolerance as to the nature of the tricyclic moiety as do the antipsychotic agents. Thus the dehydration product(s) from the condensation of ketone 97 with the Grignard reagent from 3-chloroethyl-N,N-dimethylamine affords the antidepressant diothiepin (98).¹⁹



Historically, both the tricyclic antipsychotic and antidepressant agents are derived in almost direct line from a series of tricyclic antihistaminic compounds (see 104 below). Minor changes in structure in some of the newer compounds in fact lead to drugs in which antihistaminic activity predominates. Thus ketotifen, which differs from antipsychotic compounds such as 87 only in detail, is a rather potent antihistamine. Bromination of ketone 99 occurs on the ethylene bridge to afford the 1,2 dibromide as a mixture of isomers (100); dehydrohalogenation by means of strong base gives the vinyl bromide 101 apparently as a single regioisomer. Reaction with the Grignard reagent from N-methyl-4-bromopiperidine gives the alcohol 102. Exposure




Attachment of the basic side chain to the phenothiazine nucleus by means of a carbonyl function apparently abolishes the usual CNS or antihistamine effects shown by most compounds in this class. The product azaclorzine instead is described as an antianginal agent. Reduction of proline derivative 106 with lithium aluminum hydride gives the corresponding fused piperazine 107. Use of that base to alkylate the chloroamide 109, obtained from acylation of phenothiazine with 3-chloropropionyl chloride, leads to azaclorzine (110).²¹




Fluorobutyrophenone derivatives of 4-arylpiperidines are well-known antipsychotic agents. It is thus interesting to note that the piperidine can in fact be fused onto an


indole moiety with retention of activity. Fischer indole condensation of 4-piperidone 111 with phenylhydrazine 112 leads to the indole 113. Alkylation of the anion from the indole with p-bromofluorobenzene gives the corresponding N-arylated derivative (114). Removal of the protecting group followed by alkylation on nitrogen with the acetal from p-p-fluorobutyryl chloride gives intermediate 116. Hydrolysis of the acetal followed by reduction of the ketone by means of sodium borohydride gives the antipsychotic agent flutroline (118).²²




A remarkably simple fused indole devoid of the traditional side chains is described as an antidepressant agent. Michael addition of the anion from indole ester 119 to acrylonitrile affords the cyanide 120. Selective reduction of the nitrile leads to the aminoester 121. This is then cyclized to the lactam (122). Reduction of the carbonyl group by means of lithium aluminum hydride leads to azepindole (123).²³



A fused pyrazoloquinolone provides an exception to the rule that antiallergic agents must contain a strongly acidic proton. Entry to the ring system is gained by electrocyclic reaction of diazoindolone 124 (possibly obtained by reaction of the anion from indolone with p-toluenesulfonyl azide) with propargylaldehyde. The initial adduct to the 1,3-dipole represented by the diazo group can be formulated as the spiro intermediate 125. Bond reorganization would then lead to the observed product (126). Reduction of the carbonyl with sodium borohydride leads to the corresponding alcohol, and thus pirquinozol (127).²⁴




An imidazoquinazoline constitutes still another compound that does not fall in the classification of a nonsteroid antiinflammatory agent yet shows good platelet antiaggregating activity. Condensation of benzyl chloride 128 with the ethyl ester of glycine gives alkylated product 129. Reduction of the nitro group leads to aniline 130. Reaction with cyanogen bromide possibly gives cyanamide 131 as the initial intermediate. Addition of aliphatic nitrogen would then lead to formation of the quinazoline ring (132). Amide formation between the newly formed imide and the ester would then serve to form the imidazolone ring. Whatever the details of the sequence, there is obtained in one step anagrelide (133).²⁵




A seemingly complex heterocycle which on close examination is in fact a latentiated derivative of a salicylic acid shows antiinflammatory activity. It might be speculated that this compound could quite easily undergo metabolic transformation to a salicylate and that this product is in fact the active drug. Condensation of acid 134 with hydroxylamine leads to the hydroxamic acid 135. Reaction of that with the ethyl acetal from 4-chlorobutyraldehyde then leads to the cyclic carbinolamine derivative 136. Treatment


with mild base causes internal alkylation and consequent formation of the last ring. There is thus obtained meseclazone (137).²⁶




The observation that a carboxyl derivative of a pyrimidinoquinoline shows mediator release inhibiting activity is in consonance with the earlier generalization. Knoevenagel condensation of nitroaldehyde 138 with cyanoacetamide gives the product 139. Treatment with iron in acetic acid leads to initial reduction of the nitro group (140). Addition of that function to the nitrile leads to formation of the quinoline ring (141). Reaction of that compound with ethyl oxalate results in formation of the quinazoline ring. The product, pirolate (142), is described as an antiallergy agent.²⁷




A tetracyclic heterocycle that bears little relation to any clinically used drug has been described as an anti-inflammatory agent. The compound is prepared in rather straightforward manner by initial condensation of dihalide 143 with 1,2-diaetylhydrazine. Hydrolysis of this product gives cyclic hydrazone 145. Exposure to a second mole of dihalide leads to diftalone (146).²⁸ The regiochemistry may be rationalized by assuming that the more reactive acid chloride attacks the more nucleophilic unacylated nitrogen.




The tricyclic antidepressants (as well as, incidentally the antipsychotic drugs) are characterized by a three carbon chain between the ring system and the basic nitrogen. Incorporation of one of those carbon atoms into an additional fused ring is apparently consistent with activity. Preparation of this compound involves first homologation of the side chain. Thus the carboxylic acid 147 is first converted to the acid chloride (148); reaction with diazomethane leads to the diazoketone 149. This is then subjected to photolytic rearrangement to afford the corresponding acetic acid (150). Condensation with methylaniline then gives the amide 151. Reduction with lithium aluminum hydride affords

azipramine (152).²⁹


Any migraine sufferer will willingly testify that this condition has little in common with the headaches to which the rest of mankind are subject. Recent medical studies too have shown fairly conclusively that, whatever the etiology of migraine, it is a condition quite distinct from the common headache. The syndrome is in fact so distinct as to be untouchable by the common headache cures such as aspirin. Drugs for treatment of migraine are unfortunately almost nonexistent. (The lack of appropriate animal models in no small way hinders the search for a treatment.) A benzofuranbenzoxepin has interestingly been described as an antimigraine agent. Bromination of benzofuran 153 proceeds on the methyl group to give the arylmethyl bromide 154. Displacement by phenoxide then leads to intermediate 155. Saponification (156) followed by Friedel-Crafts cyclization serves to form the seven-membered ring (157). Condensation of the ketone with the Grignard reagent from 3-chloropropyl-N,N-dimethylamine gives the olefin on dehydration, possibly as a mixture of isomers. There is thus obtained oxetorene (158).³⁰



In the steroid series, hormone antagonists usually bear some structural resemblance to the endogenous agonists. That is to say, antagonists are almost always steroids themselves. Even in the case of the nonsteroid estrogen antagonists, there is a fairly clear structural resemblance to estradiol. It is thus somewhat surprising to note a clearly nonsteroidal androgen antagonist. The compound in question, pentomone (163) is, as a result of this activity, a potential drug for treatment of prostate enlargement. Condensation of salicylaldehyde 159 with cyclohexanone 160 proceeds twice to give directly the pentacyclic intermediate 161. The reaction may be visualized as initial conjugate addition of phenoxide to the enone followed by interception of the resulting anion by the aldehyde carbonyl group. Hydrogenation of the intermediate reduces both the double bonds and the carbonyl group (162). Back oxidation of the alcohol thus formed with pyridinium chlorochromate affords pentomone (163).



The ergolines have provided a number of drugs that show interaction with neurotransmitters. Depending on the substitution pattern, they may be dopamine agonists or antagonists, α -adrenergic blockers, or inhibitors of the release of prolactin. A recent member of the series, pergolide (167), shows activity as a dopamine antagonist. Reduction of ester 164³¹ by means of lithium aluminum hydride gives the corresponding alcohol; this is then converted to its mesylate (166). Displacement with methanethiol affords pergolide (167).³²



REFERENCES

1. J. L. H. Van Gelder, A. H. M. Raeymaekers, L. F. C. Roevens and W. J. Van Laerhoven, U.S. Patent 3,925,383; Chem. Abstr., 84, 180264e (1976).
2. L. D. Spicer and J. J. Hand, French Demande 2,199,979; Chem. Abstr., 82, 16835e (1975).
3. J. Gubin and G. Rosseels, German Offen 2,707,048; Chem. Abstr., 88, 6719e (1978).
4. J. P. Maffrand and F. Elloy, Eur. J. Med. Chem., 9, 483 (1974).
5. H. J. Shaeffer, German Offen., 2,539,963 (1976).
6. J. Diamond, German Offen., 2,713,389; Chem. Abstr., 88, 22984t (1978).
7. L. E. J. Kennis and J. C. Mertens, U.S. Patent 4,347,287; Chem. Abstr., 98, 16716 (1983).
8. E. Schenker, Swiss Patent 565,797; Chem. Abstr., 83, 206311z (1975).
9. G. C. Wright, A. V. Bayless, and J. E. Gray, German Offen., 2,427,382; Chem. Abstr., 82, 171087f (1975).
10. G. C. Wright, A. V. Bayless, and J. E. Gray, German Offen., 2,427,377; Chem. Abstr., 82, 171088g (1975).
11. I. C. Nordin, U.S. Patent 3,553,207; Chem. Abstr., 75, 5972b (1971).
12. H. A. DeWald and S. J. Lobbestael, South African Patent 73 07696; Chem. Abstr., 84, 59594j (1976).

13. F. J. Tinney, U.S. Patent 3,558,606; Chem. Abstr., 74, 141896m (1971).
14. D. E. Bays, German Offen., 2058295; Chem. Abstr., 75, 98447x (1971).
15. J. R. Pfister, I. T. Harrison, and J. H. Fried, Chem. Abstr., 85, 21108g (1976).
16. C. L. Zirkle, German Offen., 2,549,841; Chem. Abstr., 85, 78025m (1976).
17. P. Herbst and D. Hoffmann, German Offen., 2,600,768; Chem. Abstr., 85, 143001s (1976).
18. J. Ackrell, Y. Antonio, F. Fidenico, R. Landeros, A. Leon, J. M. Muchowski, M. L. Maddox, P. H. Nelson, and W. H. Rooks, J. Med. Chem. 21, 1035 (1978).
19. C. L. Zirkle, U.S. Patent 3,609,167; Chem. Abstr., 75, 151694d (1971).
20. J. P. Bourquin, G. Schwarb, and E. Waldvogel, German Offen. 2,144,490; Chem. Abstr., 77, 34296f (1972).
21. N. V. Kaverina, G. A. Markova, G. G. Chichkanov, L. S. Nazarovo, A. M. Likhosherstor, and A. P. Skoldinov, Khim. Pharm. Zh., 12, 97 (1978).
22. J. J. Plattner, C. A. Harbert, J. R. Tretter, and W. M. Welch Jr., German Offen., 2,514,084; Chem. Abstr., 84, 44008x (1976).
23. B. Reynolds and J. Carson, German Offen. 1928726; Chem. Abstr., 72, 55528v (1970).
24. B. R. Vogt, German Offen. 2,726,389; Chem. Abstr., 88, 121240d (1978).
25. W. N. Beverung and R. A. Partyka, U.S. Patent 3,932,407; Chem. Abstr., 84, 10564 (1976).
26. D. B. Reisner, B. J. Ludwig, H. M. Bates, and F. M. Berger, German Offen., 2010418; Chem. Abstr., 73, 120644s (1970).

27. T. H. Althuis, L. J. Czuba, H. J. E. Hess, and S. B. Kadin, German Offen., 2,418,498; Chem. Abstr., 82, 73015m (1975).
28. E. Bellasio and E. Testa, Il Farmaco, Ed. Sci., 25, 305 (1970).
29. M. Riva, L. Toscano, A. Bianchetti, and G. Grisanti, German Offen., 2,529,792; Chem. Abstr., 84, 150530w (1976).
30. F. Binon and M. Descamps, German Offen. 1963205; Chem. Abstr., 73, 77221n (1970).
31. D. Lednicer and L. A. Mitscher, "The Organic Chemistry of Drug Synthesis", Vol. 2, Wiley, New York, 1980, p. 479.
32. E. C. Kornfeld and N. J. Bach, Eur. Patent Appl., 3,667; Chem. Abstr., 92, 181450q (1980).

Cross Index of Drugs

Aldosterone Antagonist

Spirorenone

Analgesic

Alfentanil
Anilopam
Bicifadine
Carfentanil
Ciprefadol
Ciramadol
Codorphone
Doxpicomine
Drinden

Levonantradol
Lofentanil
Moxazocine
Nabitan
Natnradol
Proxorphan
Sulfentanil
Tonazocine
Verilopam

Anaesthetic

Etomidate

Minoxalone

Androgen

Nistermine Acetate

Androgen Antagonist

Azastene
Flutamide

Pentomone
Topterone

Antiacne

Etretinate
Isotretinoin

Montretinide
Piroctone

Antiallergic

Isamoxole
Lodoxamide ethyl
Nivimedone
Oxatomide

Pirolate
Pirquinazol
Tixanox
Xanoxate

Antamebic

Quinfamide

Antianginal

Azaclorzine
Bepridil
Cinepazet
Diltiazem
Drophenilamine

Molsidomine
Nicarpidine
Nicordanil
Nimodipine
Tosifen

Antiarrhythmic

Butoprozine
Clofilium Phosphate
Disobutamine
Drobuline
Encainide
Emilium Tosylate
Flecainide

Lorcanide
Meobentine
Oxiramide
Pirmenol
Ropitoin
Tocainide

Antibacterial

Droxacin
Fludalanine
Flumequine

Metioprim
Rosoxacin
Tetroxoprim

Antibiotic

Amidinocillin
Amidinocillin Pivoxyl
Azolocillin

Cefroxadine
Cefsulodin
Ceftazidine

Bacampicillin
Cefaclor
Cefaparole
Cefatrizine
Cefazafur
Cefoperazone
Cefonicid
Ceforanide
Cefotaxime
Cefotiam

Ceftizoxime
Cefuroxime
Mezlocillin
Moxlactam
Piperacillin
Pirbencillin
Piridicillin
Sarmoxicillin
Sarpicillin

Anticonvulsant

Cinromide

Antidepressant

Azepindole
Azipramine
Cycloenzaprine
Cyclindole
Dothiepin
Fluotracer
Fluoxetine
Imafen

Napactidine
Nisoxetine
Nitrafudam
Pridefine
Tametraline
Viloxazine
Zimelidine

Antidiarrheal

Nufenoxole

Antiemetic

Domperidone

Nabilone

Antifungal

Azoconazole
Butoconazole
Doconazole
Ketoconazole
Naftidine
Orconazole
Oxifungin

Parconazole
Sulconazole
Terconazole
Tioconazole
Tolciclate
Triafungin

Anthelmintic

Butamisol

Frentizole

Carbantel
Felsantel
Fenbendazole

Nocodazole
Tioxidazole

Antihistamine

Astemizole
Ketotifen

Thiazinium Chloride

Antihypertensive

Captopril
Endralazine
Guanfacine
Indorenate
Ketanserin

Proroxan
Terazocin
Tiamenidine
Tiodazocin

Antihypertensive - β -Blocker

Bucindolol
Diacetolol
Exaprolol
Pamatolol

Penbutolol
Primidolol
Prizidolol

Antihypertensive - α, β -Blocker

Bevantolol
Labetolol

Medroxalol
Sulfinalol

Anti-inflammatory - Steroid

Acolmethasone Dipropionate
Budesonide
Ciprocinnonide
Flumoxonide

Haloprednone
Meclorisone Dibutyrate
Procinnonide

Anti-inflammatory - Non-Steroid

Anitrazafen
Amefenac
Bromperamole
Carprofen
Diftalone
Epirizole
Fencolofenac
Fenclosal
Floctafenine

Indoprofen
Isoxepace
Oxarbazole
Mesecclazone
Morniflumate
Orpanoxin
Pirazolac
Sermetacin
Talniflumate

Fluquazone
Fluproquazone
Fluretofen

Tiopinac
Zidometacin
Zomepirac

Antimalarial

Halofantrine

Antimigraine

Oxetorone

Antineoplastic

Ametantrone
Cyclophosphamide
Estramustine
Etoprine
Ifosfamide

Metoprine
Mitoxantrone
Prednimustine
Tegafur
Trofosfamide

Antiprotozoal

Bamnidazole
Ornidazole

Misonidazole

Antipsychotic

Clopipazam
Cloroperone
Declenperone
Flucindole

Fluotracen
Flutroline
Halopemide
Pipenperone

Antiulcer

Arbaprostil
Etintidine
Oxmetidine

Ranitidine
Tolimidone

Antiviral

Acyclovir
Arildone

Enviroxime

Anxiolytic

Adinazolam
Bentazepam

Lormetazepam
Midazolam

Brofoxine
Caroxazone
Elfazepam
Fosazepam

Quazepam
Ripazepam
Tioperidone
Zometapine

Bronchodilator

Ipatropium Bromide

Verofylline

Brochoclilator - β -Adrenergic

Bitolterol
Colterol
Carteolol
Dipirefrin

Nisobutero1
Prenalterol
Reproterol

Cardiotanic

Actodigin
Amrinone

Butopamine
Carbazeran

Catract Inhibitor

(Aldose Reductase Inhibitors)

Alrestatin

Sorbini1

Cognition Enhancer

Amacetam

Diagnostic Aid (Pancreatic Function)

Bentiromide

Diuretic

Azosemide
Fenquizone
Indacrinone

Muzolimine
Ozolinone
Piretanide

Dental Carries Prohylactic

Ipexidine

Dopamine Antagonist

Pergolide

Estrogen AntagonistNitromifene
Tamoxifen

Trioxifene

HypoglycemicGlicetanile
Gliflumide

Piriglriride

HypolipidemicBenzafibrate
Cetaben
CiprofibrateGemcadiol
Gemfibrozil**Immunomodulator**

Azarole

Muscle RelaxantClodanolene
Lidamidine

Xilobam

Uterine Stimulant/OxytocicCarboprost
Mefenprost

Sulprostone

Peripheral VasodilatorButerizine
CetiedilSuloctidil
Tipropidil**Platelet Agregation Inhibitor**Anagrelide
Eprostenol

Ticlopidine

Progestin

Gestodene

Gestrinone

Sedative

Fenobam

Milenperone

Vasosilator

Alprostadi1

Epoprostenol

Vitamin (D₃)

Calcifediol

Calcitriol

Cumulative Index, Vols. 1-3

- Acebutolol 2, 109
 Aceclidine 2, 295
 Acedapsone 2, 112
 Acenocoumarole 1, 331
 Aceperone 2, 332
 Acephylline 1, 425
 Acetaminophen 1, 111
 Acetanilide 1, 111; 2, 97
 Acetazolamide 1, 249
 Acetohexamide 1, 138
 Acetophenazine 1, 383
 Acetyl methoxprazine 1, 131
 Acetylmethadol 1, 81
 Aclomethasone 3, 96
 Actodigin 3, 99
 Acyclovir 3, 229
 Adinazolam 3, 197
 Adiphenine 1, 81
 Adrenalone 2, 38
 Albendazole 2, 353
 Albuterol 2, 43
 Albutoin 2, 261
 Alclofenac 2, 68
 Aldosterone 1, 206
 Aletamine 2, 48
 Alfentanyl 3, 118
 Algestone acetone 2, 171
 Algestone acetophenide 2, 171
 Alipamide 2, 94
 Allobarbitol 1, 269
 Allopurinol 1, 152, 269
 Allylestrenol 1, 172
 Alonimid 2, 295
 Aloxidone 1, 232
 Alpertine 2, 342
 Alpha eucaine 1, 8
 Alphaprodine 1, 304; 2, 328
 Alprazolam 3, 197
 Alprenolol 1, 177
 Alprostadil 3, 2
 Alrestatin 3, 72
 Althiazide 1, 359
 Alverine 2, 55
 Amacetam 3, 127
 Amantidine 2, 18
 Ambucaine 1, 11
 Ambucetamide 1, 94
 Ambuside 2, 116
 Amcinafal 2, 185
 Amcinafide 2, 185
 Amedaline 2, 348
 Ametantrone 3, 75
 Amfenac 3, 38
 Amicibone 2, 11

- Amicycline 2, 228
 Amidephrine 2, 41
 Amidinocillin 3, 208
 Amidoquine 1, 342
 Amiloride 1, 278
 Aminitrozole 1, 247
 Aminoglutethimide 1, 257
 Aminometetradine 1, 265
 Aminophenazole 1, 248
 Aminophylline 1, 427
 Aminopromazine 1, 390
 Aminopropylon 1, 234
 Aminopyrine 1, 234; 2, 262
 Aminorex 2, 265
 Amiquinsin 2, 363
 Amisometradine 1, 266
 Amitriptyline 1, 151, 404
 Amobarbital 1, 268
 Amoproxan 2, 91
 Amopyroquine 1, 342
 Amoxapine 2, 428
 Amoxycillin 1, 414
 Amphedloral 2, 48
 Amphetamine, 1, 37, 70; 2, 47
 Amphetaminil 2, 48
 Ampicillin 1, 413; 2, 437, 438
 Amprolium 1, 264
 Ampyzine 2, 298
 Amquinat 2, 370
 Amrinone 3, 147
 Anagesterone acetate 2, 165
 Anagrelide 3, 244
 Androstanolone 1, 173
 Anidoxime 2, 125
 Anileridine 1, 300
 Anilopam 3, 121
 Anisindandione 1, 147
 Anitrazafen 3, 158
 Antazoline 1, 242
 Antipyrine 1, 234
 Apazone 2, 475
 Aprindene 2, 208
 Aprobarbital 1, 268
 Aprophen 1, 91
 Arbaprostil 3, 8
 Arildone 3, 45
 Astemizol 3, 177
 Atenolol 2, 109
 Atropine 1, 35, 71, 89, 93; 2, 71
 Azabon 2, 115
 Azacosterol 2, 161
 Azaclozine 3, 241
 Azacyclonol 1, 47
 Azanator 2, 457
 Azaperone 2, 300
 Azarole 3, 129
 Azastene 3, 89
 Azatadine 2, 424
 Azathioprine 2, 464
 Azepinamide 1, 137
 Azepindole 3, 242
 Azipramine 3, 246
 Azlocillin 3, 206
 Azoconazole 3, 137
 Azolimine 2, 260
 Azomycin 1, 238
 Azosemide 3, 27
 Bacampicillin 3, 204
 Baclofen 2, 121
 Bamethan 2, 39
 Bamiphylline 1, 426
 Bamipine 1, 51
 Bamnidazole 3, 132
 Barbitol 1, 267
 BAS 2, 96
 BCNU 2, 12
 Becanthon 2, 413
 Beloxamide 2, 56
 Bemidone 1, 305
 Bemigrade 1, 258
 Benactyzine 1, 93
 Benapryzine 2, 74
 Bendazac 2, 351
 Bendroflumethiazide 1, 358
 Benfurodil 2, 355, 356
 Benorterone 2, 156
 Benoxaprofen 2, 356
 Benperidol 2, 290

- Benproperine 2, 100
 Bentazepam 3, 235
 Bentiromide 3, 60
 Benzbromarone 2, 354
 Benzestrol 1, 103
 Benzetimide 2, 293
 Benzilium bromide 2, 72
 Benzindopyrine 2, 343
 Benziodarone 1, 313
 Benzocaine 1, 9
 Benzocetamine 2, 220
 Benzodepa 2, 122
 Benzphetamine 1, 70
 Benzquinamide 1, 350
 Benztriamide 2, 290
 Benzylamine 1, 323; 2, 350
 Benzylpenicillin 1, 408
 Bepridil 3, 46
 Beta eucaine 1, 9
 Betahistine 2, 279
 Betamethasone 1, 198
 Bethanidine 1, 55
 Bevantolol 3, 28
 Bezafibrate 3, 44
 Bicifadine 3, 120
 Biperiden 1, 47
 Bitolterol 3, 22
 Bishydroxycoumarin 1, 331
 Bolandiol diacetate 2, 143
 Bolasterone 1, 173; 2, 154
 Boldenone 2, 153
 Bolmantalate 2, 143
 Boxidine 2, 99
 Bretylium tosylate 1, 55
 Brofoxine 3, 191
 Bromperidol 2, 331
 Bromhexine 2, 96
 Bromindione 2, 210
 Bromisovalum 1, 221
 Bromodiphenhydramine 1, 42
 Bromoxanide 2, 94
 Bromperidol 2, 331
 Broperamole 3, 139
 Brompheniramine 1, 77
 Bucainide 2, 125
 Bucindolol 3, 28
 Buclizine 1, 59
 Bucloxic acid 2, 126
 Budesonide 3, 95
 Buformin 1, 221; 2, 21
 Bufuralol 2, 110
 Bumetanide 2, 87
 Bunaftine 2, 211
 Bunamidine 2, 212
 Bunitridine 2, 215
 Bunitrolol 2, 106, 110
 Bunolol 2, 110, 215
 Bupicomide 2, 280
 Bupivacaine 1, 17
 Buprenorphine 2, 321
 Bupropion 2, 124
 Buquinolate 1, 346
 Burimamide 2, 251
 Buspirone 2, 300
 Butabarbital 1, 268
 Butacaine 1, 12
 Butacetin 2, 95
 Butaclamol 2, 226
 Butalbital 1, 268
 Butamirate 2, 76
 Butamisole 3, 226
 Butaperazine 1, 381
 Butazolamide 1, 249
 Buterizine 3, 175
 Butethal 1, 268
 Butoconazole 3, 134
 Butorphanol 2, 325
 Butoxamine 1, 68
 Butriptyline 1, 151
 Butropium bromide 2, 308
 Butylallonal 1, 269
 Butylvinal 1, 269
 Caffeine 1, 111, 423
 Calcifediol 3, 101
 Calcitriol 3, 103
 Calusterone 2, 154
 Cambendazole 2, 353
 Canrenoate 2, 174
 Canrenone 2, 174
 Capobenic acid 2, 94
 Captodiamine 1, 44

- Captopril 3, 128
 Caramiphen 1, 90
 Carbacephalothin 1, 420
 Carbadox 2, 390
 Carbamazepine 1, 403
 Carbantel 3, 57
 Carbaprost 3, 7
 Carbazeran 3, 195
 Carbenicillin 1, 414; 2, 437
 Carbetidine 1, 90
 Carbidopa 2, 119
 Carbimazole 1, 240
 Carbinoxamine 1, 43; 2, 32
 Carbiphen 2, 78
 Carbromal 1, 221
 Carbubarbital 1, 269, 273
 Carbutamide 1, 138
 Carbuterol 2, 41
 Carfentanyl 3, 117
 Carisoprodol 1, 219
 Carmantadine 2, 20
 Carmustine 2, 12
 Carnidazole 2, 245
 Caroxazone 3, 191
 Carphenazine 1, 383
 Carpipramine 2, 416
 Carprofen 3, 169
 Cartazolate 2, 469
 Carteolol 3, 183
 Cefaclor 3, 209
 Cefadroxil 2, 440
 Cefamandole 2, 441
 Cefaparole 3, 212
 Cefatrizine 3, 211
 Cefazaflur 3, 213
 Cefazolin 3, 442
 Cefonicid 3, 213
 Ceforanide 3, 214
 Cefotaxime 3, 216
 Cefotiam 3, 215
 Cefoxitin 2, 435, 443
 Cefroxadine 3, 210
 Cefsulodin 3, 214
 Ceftazidine 3, 216
 Ceftizoxime 3, 218
 Cefuroxime 3, 216
 Cephalixin 1, 417; 2, 439
 Cephaloglycin 1, 417
 Cephaloridine 1, 417
 Cephalothin 1, 417 420
 Cephapirin 2, 441
 Cephradine 2, 440
 Cetaben 3, 60
 Cetophenicol 2, 46
 Cetiedil 3, 42
 Chloraminophenamide 1, 133
 Chlofluperidol 1, 306
 Chloramphenicol 1, 75; 2, 28, 45
 Chlorazaniol 1, 281
 Chlorbenzoxamine 1, 43
 Chlorcyclizine 1, 58
 Chlordiazepoxide 1, 365; 2, 401
 Chlorexolone 1, 321
 Chlorguanide 1, 115
 Chlorimpiphenine 1, 385
 Chlorindandione 1, 147
 Chlormadinone acetate 1, 181; 2, 165
 Chlormidazole 1, 324
 Chlorophenylalanine 2, 52
 Chloroprocaine 1, 11
 Chloropyramine 1, 402
 Chloroquine 1, 341
 Chlorothen 1, 54
 Chlorothiazide 1, 321, 355; 2, 395
 Chlorotrianisene 1, 104
 Chlorphenamide 1, 133, 358
 Chlorphendianol 1, 46
 Chlorphenesin 1, 118
 Chlorpheniramine 1, 77
 Chlorphenoxamine 1, 44
 Chlorphentermine 1, 73
 Chlorproethazine 1, 379
 Chlorproguanil 1, 115
 Chlorpromazine 1, 319 378; 2, 409; 3, 72
 Chlorpropamide 1, 137
 Chlorprothixene 1, 399
 Chlorpyramine 1, 51

- Chlortetracycline 1, 212
 Chlorthalidone 1, 322
 Chlorzoxazone 1, 323
 Chromoglycate 1, 313, 336
 Chromonor 1, 331
 Ciclafrine 2, 266
 Ciclopirox 2, 282
 Cicloprofen 2, 217
 Cimetidine 2, 253
 Cinanserin 2, 96
 Cinepazet 3, 157
 Cinepazide 2, 301
 Cingestol 2, 145
 Cinnamedrine 2, 39
 Cinnarizine 1, 58
 Cinoxacin 2, 388
 Cinromide 3, 44
 Cintazone 2, 388, 474
 Cintriamide 2, 121
 Ciprefadol 3, 119
 Ciprocinonide 3, 94
 Ciprofibrate 3, 44
 Ciramadol 3, 122, 123
 Citenamide 2, 221
 Clamoxyquin 2, 362
 Clazolam 2, 452
 Clazolimine 2, 260
 Clemastine 2, 32
 Clemizole 1, 324
 Clioxanide 2, 94
 Cliprofen 2, 65
 Clobazam 2, 406
 ClobutinoI 2, 121
 Clocapramine 2, 416
 Clocental 1, 38
 Clocortolone acetate 2, 193
 Clodanole 3, 130
 Clodazon 2, 354
 Clofenpyride 2, 101
 Clofibrate 1, 119; 2, 79, 101, 432
 Clofilium phosphate 3, 46
 Clogestone 2, 166
 Clomacran 2, 414
 Clomegestone acetate 2, 170
 Clometherone 2, 170
 Clomifene 1, 105, 148; 2, 127
 Clominorex 2, 265
 Clonidine 1, 241
 Clonitazene 1, 325
 Clonixeril 2, 281
 Clonixin 2, 281
 Clopamide 1, 135; 2, 93
 Clopenthixol 1, 399
 Cloperidone 2, 387
 Cloperone 3, 150
 Clopimozide 2, 300
 Clopipazam 3, 237
 Clopirac 2, 235
 Cloprednol 2, 182
 Cloprostenol 2, 6
 Clorprenaline 2, 39
 Closantel 3, 43
 Closiramine 2, 424
 Clothiapine 1, 406; 2, 429
 Clothixamide 2, 412
 Cloxacillin 1, 413
 Cloxazepam 1, 370
 Clozapine 2, 425
 Codeine 1, 287; 2, 317
 Codorphone 3, 112
 Codoxime 2, 318
 Colchicine 1, 152, 426
 Colterol 3, 21
 Cormethazone acetate 2, 194, 196
 Cortisone 1, 188; 2, 176, 179
 Cortisone acetate 1, 188, 190
 Cortivazol 2, 191
 Cotinine 2, 235
 Cromoglycate 3, 66, 235
 Cyclacillin 2, 439
 Cyclandelate 1, 94
 Cyclazocine 1, 298; 2, 327
 Cyclindole 3, 168
 Cyclizine 1, 58
 Cyclobarbitol 1, 269
 Cyclobendazole 2, 353
 Cyclobenzaprine 3, 77
 Cycloguanil 1, 281
 Cyclomethycaine 1, 14
 Cyclopal 1, 269

- Cyclopentolate 1, 92
 Cyclopenthiiazide 1, 358
 Cyclophosphamide 3, 161
 Cyclopryazate 1, 92
 Cycloserine 3, 14
 Cyclothiazide 1, 358
 CyCRMine 1, 47
 Cyheptamide 2, 222
 Cypenamine 2, 7
 Cyprazepam 2, 402
 Cyproheptadine 1, 151
 Cyprolidol 2, 31
 CyproquinatE 2, 368
 Cyproterone acetate 2, 166
 Cyrapoximide 2, 293

 Dacarbazine 2, 254
 Daledanin 2, 348
 Danazol 2, 157
 Dantrolene 2, 242
 Dapsone 1, 139; 2, 112
 Dazadrol 2, 257
 Debrisoquine 1, 350; 2, 374
 Declenperone 3, 172
 DecoquinatE 2, 368
 Delmadinone acetate 2, 166
 Demoxepam 2, 401
 Deprostit 2, 3
 Descinolone acetonide 2, 187
 Deserpidine 1, 320
 Desipramine 1, 402
 Desonide 2, 179
 Deterenol 2, 39
 Dexamethasone 1, 199
 Dexbrompheniramine 1, 77
 Dexchlorpheniramine 1, 77
 Dexivacaine 2, 95
 Dexnorgestrel acetate 2, 152
 Dextroamphetamine 1, 70
 Dextromoramide 1, 82
 Dextromorphan 1, 293
 Dextrothyroxine 1, 92
 Diacetolol 3, 28
 Diamocaine 2, 336
 Dianithazole 1, 327
 Diapamide 2, 93

 Diaveridine 2, 302
 Diazepam 1, 365; 2, 452
 Diazoxide 1, 355; 2, 395
 Dibenamine 1, 55
 Dibenzepin 1, 405; 2, 424, 471
 Dibucaine 1, 15
 Dichlorisone 1, 203
 Dichloroisoproterenol 1, 65;
 2, 106
 Dichlorophenamide 1, 133
 Diclofenac 2, 70
 Dicloxacillin 1, 413
 Dicoumarol 1, 147
 Dicyclomine 1, 36
 Dienestrol 1, 102
 Diethyl carbamazine 1, 278
 Diethylstilbestrol 1, 101
 Diethylthiambutene 1, 106
 Difenoximide 2, 331
 Difenoxin 2, 331
 Diflucortolone 2, 192
 Diflumidone 2, 98
 Diflunisal 2, 85
 Difluprednate 2, 191
 Diftalone 3, 246
 Dihexyverine 1, 36
 Dihydralazine 1, 353
 Dihydrocodeine 1, 288
 Diltiazem 3, 198
 Dimefadane 2, 210
 DimeflinE 2, 391
 Dimetacrine 1, 397
 Dimethisoquine 1, 18
 Dimethisterone 1, 176, 187
 Dimethothiazine 1, 374
 Dimethoxanate 1, 390
 Dimethylpyrindene 1, 145
 Dimethylthiambutene 1, 106
 Dimetridazole 1, 240
 Dinoprost 1, 27, 33
 Dinoprostone 1, 27, 30, 33, 35
 Dioxadrol 2, 285
 Dioxyline 1, 349
 Diphenhydramine 1, 41
 Diphenidol 1, 45
 Diphenoxylate 1, 302; 2, 331

- Diphenylhydantion 1, 246
 Diphepanol 1, 46
 Dipipanone 1, 80
 Dipipoverine 1, 94
 Dipirefrin 3, 22
 Dipyridamole 1, 428
 Dipyrone 2, 262
 Disobutamide 3, 41
 Disopyramide 2, 81; 3, 41
 Disulfiram 1, 223
 Dithiazanine 1, 327
 Dixyrazine 1, 384
 Dobutamine 2, 53
 Doconazole 3, 133
 Domazoline 2, 256
 Domperidone 3, 174
 Dopamantine 2, 52
 Dorastine 2, 457
 Dothiepin 3, 239
 Doxapram 2, 236
 Doxaprost 2, 3
 Doxepin 1, 404
 Doxpicomine 3, 122
 Doxylamine 1, 44
 Drindene 3, 65
 Drobuline 3, 47
 Drocinnonide 2, 186
 Dromostanolone 1, 173
 Droprenylamine 3, 47
 Droperidol 1, 308
 Droxacin 3, 185
 Dydrogesterone 1, 185
- Econazole 2, 249
 Ectylurea 1, 221
 Elantrine 2, 418
 Elfazepam 3, 195
 Elucaine 2, 44
 Emilium tosylate 3, 47
 Encainide 3, 56
 Encyprate 2, 27
 Endralazine 3, 232
 Endrysone 2, 200
 Enviroxime 3, 177
 Ephedrine 1, 66
 Epimestrol 2, 13
- Epinephrine 1, 95, 241
 Epirazole 3, 152
 Epithiazide 1, 359
 Epoprostenol 3, 10
 Eprazinone 1, 64
 Eprozinol 2, 44
 Eritadenine 2, 467
 Erythryptamine 1, 317
 Esproquin 2, 373
 Estramustine 3, 83
 Estrazinol 2, 142
 Estradiol 1, 162; 2, 136
 Estradiol benzoate 1, 162
 Estradiol cypionate 1, 162
 Estradiol dipropionate 1, 162
 Estradiol hexabenzoate 1, 162
 Estradiol valerate 1, 162
 Estramustine 3, 83
 Estrofurate 2, 137
 Estrone 1, 156, 161
 Etafedrine 2, 39
 Etazolate 2, 469
 Eterobarb 2, 304
 Ethacrynic acid 1, 120; 2, 103
 Ethambutol 1, 222
 Ethamivan 2, 94
 Ethionamide 1, 255
 Ethisterone 1, 163, 172
 Ethithiazide 1, 358
 Ethoheptazine 1, 303
 Ethonam 2, 249
 Ethopropazine 1, 373
 Ethosuximide 1, 228
 Ethotoin 1, 245
 Ethoxzolamide 1, 327
 Ethylestrenol 1, 170
 Ethylmorphine 1, 287
 Ethynerone 2, 146
 Ethynodiol diacetate 1, 165, 186
 Ethynodrel 1, 164
 Ethynylestradiol 1, 162
 Etidocaine 2, 95
 Etingidine 3, 135
 Etoclofen 2, 89
 Etomidate 3, 135

- Etonitazene 1, 325
 Etoprine 3, 153
 Etorphine 2, 321
 Etoxidrol 2, 285
 Etapralol 3, 28

 Famotine 2, 37
 Fantridone 2, 421
 Felsantel 3, 57
 Fenalamide 2, 81
 Fenbendazole 3, 176
 Fenbufen 2, 126
 Fencamfamine 1, 74
 Fenclofenac 3, 37
 Fenclorac 2, 66
 Fenclozic acid 2, 269
 Fendosal 3, 170
 Fenestrel 2, 9
 Fenethylline 1, 425
 Fenfluramine 1, 70
 Fenimide 2, 237
 Fenisorex 2, 391
 Fenmetozole 2, 257
 Fenobam 3, 136
 Fenoprofen 2, 67
 Fenoterol 2, 38
 Fenpipalone 2, 293
 Fenquizon 3, 192
 Fenspiriden 2, 291
 Fentanyl 1, 299, 306, 309; 3, 116
 Fenyripol 2, 40
 Fetoxylate 2, 331
 Flavoxate 2, 392
 Flazalone 2, 337
 Flecainide 3, 59
 Fletazepam 2, 403
 Floctafenine 3, 184
 Floxacillin 1, 413
 Fluanisone 1, 279
 Fluazepam 1, 366
 Flubanilate 2, 98
 Flubendazole 2, 354
 Flubiprofen 1, 86
 Flucindol 3, 168
 Flucinolone 3, 94
 Flucinolone acetone 1, 202
 Flucloronide 2, 198
 Fludalanine 3, 14
 Fludorex 2, 44
 Fludrocortisone 1, 192
 Fludroxycortide 1, 202
 Flufenamic acid 1, 110; 2, 69
 Flumequine 3, 186
 Flumethasone 1, 200
 Flumethiazide 1, 355; 2, 355
 Flumetramide 2, 306
 Fluminorex 2, 265
 Flumizole 2, 254; 3, 158
 Flumoxonide 3, 95
 Flunarizine 2, 31
 Flunidazole 2, 246
 Flunisolid 2, 181
 Flunitrazepam 2, 406
 Flunixin 2, 281
 Fluproquazone 3, 193
 Fluorocortolone 1, 204
 Fluorometholone 1, 203
 Fluoroprednisolone, 9 α , 1, 292
 Fluorouracil, 5-, 3, 155
 Fluotracen 3, 73, 74
 Fluoxetine 3, 32
 Fluoxymestron 1, 175
 Fluperamide 2, 334
 Fluperolone acetate 2, 185
 Fluphenazine 1, 383
 Fluprostenol 2, 6
 Fluquazone 3, 193
 Flurandrenolide 2, 180
 Fluretofen 3, 39
 Flurogestone acetate 2, 183
 Fluspiरणone 2, 292
 Fluspirilene 2, 292
 Flutamide 3, 57
 Flutiazine 2, 431
 Flutroline 3, 242
 Formocortal 2, 189
 Fosazepam 3, 195
 Frentizole 3, 179
 Furaltadone 1, 229
 Furazolidone 1, 229
 Furethidine 1, 301

- Furobufen 2, 416
 Furosemide 1, 134; 2, 87
 Fusaric acid 2, 279
- Gamfexine 2, 56
 Gemcadiol 3, 15
 Gemfibrozil 3, 45
 Gestaclone 2, 169
 Gestodene 3, 85
 Gestonorone 2, 152
 Gestrinone 3, 85
 Glaphenine 1, 342
 Gliamilide 2, 286
 Glibornuride 2, 117
 Glicetanile 3, 61
 Gliflumide 3, 61
 Glipizide 2, 117
 Glutethemide 1, 257
 Glyburide 2, 139
 Glybuthiazole 1, 126
 Glycosulfone 1, 140
 Glyhexamide 1, 138
 Glymidine 1, 125
 Glyoctamide 2, 117
 Glyparamide 2, 117
 Glyprothiazole 1, 125
 Griseofulvin 1, 314
 Guaiaphenesin 1, 118
 Guanabenz 2, 123
 Guanacine 1, 260
 Guanadrel 1, 400
 Guanethidine 1, 282; 2, 100
 Guanfacine 3, 40
 Guanisoquin 2, 375
 Guanoclor 1, 117; 2, 101
 Guanoxabenz 2, 123
 Guanoxan 1, 352
 Guanoxyfen 2, 101
- Halcinonide 2, 187
 Halofantrine 3, 76
 Halofenate 2, 80, 102
 Halopemide 3, 174
 Haloperidol 1, 306
 Haloprednone 3, 99
 Haloprogestosterone 3, 173
- Heptabarbital 1, 269, 272
 Hepzidine 2, 222
 Heroin 1, 288; 2, 315
 Hetacillin 1, 414
 Heteronium bromide 2, 72
 Hexestrol 1, 102
 Hexethal 1, 268
 Hexobarbital 1, 273
 Hexobendine 2, 92
 Hexylcaine 1, 12
 Histapyrrodine 1, 50
 Hoquizil 2, 381
 Hycanthone 1, 398; 2, 413
 Hydracarbazine 2, 305
 Hydralazine 1, 353
 Hydrochlorobenzethylamine 1, 59
 Hydrochlorothiazide 1, 358
 Hydrocodone 1, 288
 Hydrocortisone 1, 190
 Hydrocortisone acetate 1, 190
 Hydroflumethiazide 1, 358
 Hydromorphone 1, 288
 Hydroxyamphetamine 1, 71
 Hydroxychloroquine 1, 342
 Hydroxyphenamate 1, 220
 Hydroxyprocaine 1, 11
 Hydroxyprogesterone 1, 176, 190
 Hydroxyzine 1, 59
- Ibufenac 1, 86
 Ibuprofen 1, 86; 2, 218, 356
 Ifenprodil 2, 39
 Ifosfamide 3, 151
 Imafen 3, 226
 Imidoline 2, 259
 Imipramine 1, 401; 2, 420; 3, 32
 Imolamine 1, 249
 Indacrinone 3, 67
 Indapamide 2, 349
 Indomethacin 1, 318; 2, 345; 3, 165
 Indoprofen 3, 171
 Indoramine 2, 344

- Indorenate 3, 167
 Indoxole 2, 254, 340; 3, 158
 Intrazole 2, 345
 Intriptyline 2, 223
 Ipexidine 3, 157
 Ipratropium bromide 3, 160
 Iodothiouracil 1, 265
 Iprindol 1, 318
 Ipronidazole 2, 244
 Iproniazide 1, 254
 Isamoxole 3, 138
 Isoaminile 1, 82
 Isobucaine 1, 12
 Isobuzole 2, 272
 Isocarboxazide 1, 233; 2, 266
 Isoetharine 2, 9
 Isomethadone 1, 79
 Isomylamine 2, 11
 Isoniazide 1, 254; 2, 266
 Isopentaquine 1, 346
 Isoproterenol 1, 63; 2, 37, 107; 3, 20, 21
 Isopyrine 1, 234
 Isothipendyl 1, 430
 Isotretinoin 3, 12
 Isoxepac 3, 238
 Isoxicam 2, 394
 Isoxsuprine 1, 69

 Ketanserin 3, 193
 Ketamine 1, 57; 2, 16
 Ketasone 1, 237
 Ketazocine 2, 328
 Ketazolam 1, 369
 Ketobemidone 1, 303
 Ketoconazole 3, 132
 Ketoprofen 2, 64
 Ketotifen 3, 239
 Khellin 1, 313, 335

 Labetolol 3, 24
 Legotrile 2, 480
 Leniquinsin 2, 363
 Lenperone 2, 286
 Letimide 2, 393
 Levalorphanol 1, 293

 Levarterenol 1, 63
 Levonantradol 3, 188
 Levonordefrine 1, 68
 Levophenacymorphan 1, 294
 Levopropoxyphene 1, 50
 Levothyroxine 1, 97
 Lidamide 3, 56
 Lidocaine 1, 16; 2, 95, 449; 3, 40
 Lidoflazine 1, 279
 Lifibrate 2, 103
 Liothyronine 1, 97
 Lobendazole 2, 353
 Lodoxamide 3, 57
 Lofentanyl 3, 117
 Lometraline 2, 214
 Lomustine 2, 12, 15
 Loperamide 2, 334
 Lorazepam 1, 368
 Lorbamate 2, 21
 Lorcaïnide 3, 40
 Lormetazepam 3, 196
 Loxapine 2, 427
 Lucanthone 1, 397; 2, 413
 Lynestrol 1, 166, 186

 Mafenide 2, 114
 Maprotiline 2, 220
 Mazindol 2, 462
 Mebendazole 2, 353
 Mebeverine 2, 54
 Mebhydroline 1, 319
 Mebrophenhydramine 1, 44
 Mebutamate 1, 218
 MeCCNU 2, 12
 Mecillinam 3, 208
 Mecloastine 1, 44
 Meclozine 1, 59
 Meclocycline 2, 227
 Meclofenamic acid 1, 110; 2, 88
 Meclorisonone butyrate 3, 95
 Medazepam 1, 368
 Medibazine 2, 30
 Mediquox 2, 390
 Medrogestone 1, 182

- Medroxoalol 3, 25
 Medroxyprogesterone 1, 180,
 186; 2, 165
 Medrylamine 1, 41
 Medrysone 2, 200
 Mefenamic acid 1, 110; 2, 280
 Mefenorex 2, 47
 Mefexamide 2, 103
 Mefruside 1, 134
 Megesterol acetate 1, 180
 Melengesterol acetate 1, 182,
 187
 Melitracen 2, 220
 Melphalan 2, 120
 Memotine 2, 378
 Menoctone 2, 217
 Meobentine 3, 45
 Meparfynol 1, 38
 Meperidine 1, 300; 2, 328; 3,
 116
 Mephenhydramine 1, 44
 Mephenesin 1, 118
 Mephenesin carbamate 1, 118
 Mephenoxalone 1, 119
 Mephentermine 1, 72
 Mephenytoin 1, 246
 Mephobarbital 1, 273
 Mepivacaine 1, 17
 Meprobamate 1, 218; 2, 21
 Mequoqualone 1, 354
 Meralluride 1, 224
 Mercaptomerine 1, 224
 Mesecclazone 3, 245
 Mesoridazine 1, 389
 Mesterolone 1, 174
 Mestranol 1, 162
 Mesuprine 2, 41
 Metabutoxycaine 1, 11
 Metalol 2, 41
 Metampicillin 1, 414
 Metaproterenol 1, 64
 Metaxalone 1, 119
 Meteneprost 3, 9
 Metformin 2, 20
 Methacycline 2, 227
 Methadone 1, 79, 289, 298; 2, 328
 Methallenestril 1, 87
 Methamphetamine 1, 37, 70
 Methandrostenolone 1, 173
 Methantheline bromide 1, 393
 Methaphencycline 1, 53
 Methaphenilene 1, 52
 Methaprylon 1, 259
 Methapyrilene 1, 54
 Methaqualone 1, 353
 Metharbital 1, 273
 Methazolamide 1, 250
 Methdilazine 1, 387
 Methenolone acetate 1, 175
 Methicillin 1, 412
 Methimazole 1, 240
 Methisazone 2, 350
 Methitural 1, 275
 Methixene 1, 400; 2, 413
 Methocarbamol 1, 118
 Methohexital 1, 269
 Methopholine 1, 349
 Methopromazine 1, 374
 Methoxsalen 1, 333
 Methoxypromazine 1, 378
 Methsuximide 1, 228
 Methyclothiazide 1, 360
 Methylchromone 1, 335
 Methyl dihydromorphinone 1, 292
 Methyl dopa 1, 95
 Methylphenidate 1, 88
 Methylprednisolone 1, 193
 Methylprednisolone, 16 β -, 1,
 196
 Methyltestosterone 1, 172
 Methylthiouracil 1, 264
 Methynodiol diacetate 2, 149
 Methyridine 1, 265
 Methysergide 2, 477
 Metiamide 2, 252
 Metiapine 2, 429
 Metioprim 3, 155
 Metizoline 2, 256
 Metolazone 2, 384
 Metopimazine 1, 386
 Metoprine 3, 153
 Metoprolol 2, 109

- Metronidazole 1, 240
 Mexenone 2, 175
 Mexrenoate 2, 175
 Mezlocillin 3, 206
 Mianserin 2, 451
 Mibolerone 2, 144
 Miconazole 2, 249
 Midaflur 2, 259
 Midazolam 3, 197
 Milenperone 3, 172
 Milipertine 2, 341
 Mimbane 2, 347
 Minaxalone 3, 90
 Minocycline 1, 214; 2, 228
 Minoxidil 1, 262
 Misonidazole 3, 132
 Mitoxantrone 3, 75
 Mixidine 2, 54
 Modaline 2, 299
 Mofebutazone 1, 234
 Molinazone 2, 395
 Molindone 2, 455
 Molsidomine 3, 140
 Moprolol 2, 109
 Morantel 1, 266
 Morazone 2, 261
 Morniflumate 3, 146
 Morphazineamide 1, 277
 Morpheridine 1, 300
 Morphine 1, 286; 2, 314; 3, 110
 Motretinide 3, 12
 Moxalactam 3, 218
 Moxazocine 3, 114
 Moxisylyte 1, 116
 Moxnidazole 2, 246
 Muzolimine 3, 137

 Nabilone 3, 189
 Nabitan 3, 190
 Nadolol 2, 110
 Nafacillin 1, 412
 Nafenopin 2, 214
 Nafomine 2, 212
 Nafoxidine 1, 147; 3, 70
 Nafronyl 2, 213
 Naftidine 3, 372

 Nalbuphine 2, 319
 Nalidixic acid 1, 429; 2, 370, 469
 Nalmexone 2, 319
 Nalorphine 1, 288; 2, 318
 Naloxone 1, 289; 2, 318, 323
 Naltrexone 2, 319
 Namoxirate 1, 86
 Nandrolone 1, 164
 Nandrolone decanoate 1, 171
 Nandrolone phenpropionate 1, 171
 Nantradol 3, 186
 Napactidine 3, 71
 Naphazoline 1, 241
 Naproxen 1, 86
 Naranol 2, 454
 Nefopam 2, 447
 Neostigmine 1, 114
 Nequinat 2, 369
 Nexeridine 2, 17
 Nialamide 1, 254
 Nicarpidine 3, 150
 Nicergoline 2, 478
 Niclosamide 2, 94
 Nicotinic acid 1, 253
 Nicordanil 3, 148
 Nicotiny alcohol 1, 253
 Nidroxyzone 1, 228
 Nifedipine 2, 283
 Nifenazone 1, 234
 Nifluminic acid 1, 256
 Nifuratrone 2, 238
 Nifurdazil 2, 239
 Nifurmid 2, 239
 Nifuroxime 2, 238
 Nifurpirinol 2, 240
 Nifurprazine 1, 231
 Nifurquinazol 2, 383
 Nifursemitone 2, 238
 Nifurthiazole 2, 241
 Nikethamide 1, 253
 Nimazone 2, 260
 Nimodipine 3, 149
 Nimorazole 2, 244
 Niridazole 2, 269

- Nisobamate 2, 22
 Nisobutanol 3, 23
 Nisoxetine 3, 32
 Nistremine acetate 3, 88
 Nithiazole 2, 268
 Nitrafudam 3, 130
 Nitrazepam 1, 366
 Nitrimidazine 1, 240
 Nitrofurantoin 1, 230
 Nitrofurantel 1, 229
 Nitrofurazone 1, 229
 Nitrofuraxime 1, 228
 Nitromifene 3, 51
 Nivazol 2, 159
 Nivimedone 3, 67
 Nocodazole 3, 176
 Noracymethadol 2, 58
 Norbolethone 2, 151
 Norephedrine 1, 260
 Norethandrolone 1, 170
 Norethindrone acetate 1, 165
 Norethindrone 1, 164; 2, 145
 Norethynodrel 1, 186
 Norgestatriene 1, 168
 Norgestrel 1, 167; 2, 151; 3, 84
 Norgestatrienone 1, 186
 Normeperidine 1, 300
 Normethandrone 1, 170
 Normethadone 1, 81
 Norpipanone 1, 81
 Nortriptyline 1, 151
 Nufenoxole 3, 42
 Nyldrin 1, 69

 Octazamide 2, 448
 Octriptyline 2, 223
 Orconazole 3, 133
 Ormetoprim 2, 302
 Ornidazole 3, 131
 Orpanoxin 3, 130
 Orphenadrine 1, 42
 Oxacephalothin 1, 420
 Oxacillin 1, 413
 Oxamniquine 2, 372
 Oxanamide 1, 220

 Oxandrolone 1, 174
 Oxantel 2, 303
 Oxaprozin 2, 263
 Oxarbazole 3, 169
 Oxatomide 3, 173
 Oxazepam 1, 366; 2, 402
 Oxazolapam 1, 370
 Oxeladine 1, 90
 Oxethazaine 1, 72
 Oxetorene 3, 247
 Oxfendazole 2, 353
 Oxibendazole 2, 352
 Oxilorphan 2, 325
 Oxiperomide 2, 290
 Oxiramide 3, 40
 Oxisuran 2, 280
 Oxmetidine 3, 134
 Oxolamine 1, 248
 Oxolinic acid 2, 370, 387; 3, 185
 Oxprenolol 1, 117; 2, 109
 Oxybutynin 1, 93
 Oxycodone 1, 290
 Oxyfedrine 2, 40
 Oxyfungin 3, 233
 Oxymestrolone 1, 173
 Oxymetazoline 1, 242
 Oxymetholone 1, 173
 Oxymorphone 1, 290; 2, 319
 Oxypendyl 1, 430
 Oxypertine 2, 343
 Oxyphenbutazone 1, 236
 Oxyphencyclimine 2, 75
 Oxyphenisatin 2, 350
 Oxyprenolol 1, 117
 Oxyipurinol 1, 426
 Oxytetracycline 1, 212; 2, 226
 Ozolinone 3, 140

 Paludrine 1, 115
 Pamaquine 1, 345
 Pamatolol 3, 28
 Pancuronium chloride 2, 163
 Papaverine 1, 347
 Para-aminosalicylic acid 1, 109

- Paraethoxycaine 1, 10
 Paramethadione 1, 232
 Paramethasone 1, 200
 Paranyline 2, 218
 Parapenzolate bromide 2, 75
 Parconazole 3, 133
 Pargyline 1, 54; 2, 27
 Pazoxide 2, 395
 Pacazine 1, 387
 Pemetid 2, 288
 Penfluridol 2, 334
 Pentethylcyclohexanone 1, 38
 Pentapiperium 2, 76
 Pentaquine 1, 346
 Pentazocine 1, 297; 2, 325
 Pentobarbital 1, 268
 Pentomone 3, 248
 Pentoxphylline 2, 466
 Pentylene-tetrazole 1, 281
 Perazine 1, 381
 Pergolide 3, 249
 Perlazine 2, 425
 Perphenazine 1, 383
 Pethidine 1, 300
 Phenacaine 1, 19
 Phenacemide 1, 95
 Phenacetemide 3, 44
 Phenacetin 1, 111
 Phenadoxone 1, 80
 Phenaglycodol 1, 219
 Phenazocine 1, 298
 Phenazopyridine 1, 255
 Phenbenicillin 1, 410
 Phenbenzamine 1, 50
 Phenbutalol 2, 110
 Phencarbamide 2, 97
 Phencyclidine 1, 56
 Phendimetrazine 1, 260; 2, 261
 Phenelazine 1, 74
 Pheneridine 1, 301
 Phenethicillin 1, 410
 Phenformin 1, 75
 Phenindandone 1, 147
 Pheniprane 1, 76
 Pheniprazine 1, 74
 Pheniramine 1, 77
 Phenmetrazine 1, 260
 Phenobarbital 1, 268
 Phenolphthalein 1, 316
 Phenomorphan 1, 294
 Phenoperidine 1, 302
 Phenoxybenzmine 1, 55
 Phenoxy-methylpenicillin 1, 410
 Phensuximide 1, 226
 Phentermine 1, 72
 Phenolamine 1, 242
 Phenyl aminosalicylate 2, 89
 Phenylbutazone 1, 236; 2, 388, 474
 Phenylephrine 1, 63; 2, 265; 3, 20
 Phenylglutarimide 1, 257
 Phenyltoloxamine 1, 115
 Phenylamidol 1, 165
 Pholcodine 1, 287
 Phthaloyl sulfathiazole 1, 132
 Physostigmine 1, 111
 Pimetine 2, 286
 Piminodine 1, 301
 Pimozide 2, 290
 Pindolol 2, 342
 Pinoxepin 2, 419
 Pipamazine 1, 385
 Pipamperone 2, 288
 Pipazethate 1, 390
 Piperacillin 3, 207
 Piperacetazine 1, 386
 Piperidolate 1, 91
 Piperocaine 1, 13
 Piperoxan 1, 352
 Pipobroman 2, 299
 Pipo-sulfan 2, 299
 Pipradol 1, 47
 Piprozolin 2, 270
 Piquizil 2, 381
 Pirandamine 2, 459
 Pirazolac 3, 138
 Pirbencillin 3, 207
 Pirbutorol 2, 280
 Pirenperone 3, 231
 Piretanide 3, 58
 Pirexyl 1, 115

- Piridicillin 3, 207
 Piridocaine 1, 13
 Pirindol 1, 45
 Piritramide 1, 308
 Pirmetol 3, 48
 Piroctone 3, 149
 Pirogliride 3, 57
 Pirolate 3, 245
 Piromidic acid 2, 470
 Pirprofen 2, 69
 Pirquinozol 3, 243
 Pivampicillin 1, 414
 Pizoctyline 2, 420
 Poldine methylsulfate 2, 74
 Polythiazide 1, 360
 Practolol 2, 106, 108
 Pranolium chloride 2, 212
 Pramoxine 1, 18
 Prazepam 2, 405
 Prazocin 2, 382; 3, 194
 Prednimustine 3, 93
 Prednisolone 1, 192; 2, 178
 Prednisolone acetate 1, 192
 Prednisone 1, 192
 Prednival 2, 179
 Prednylene 1, 197
 Prenalterol 3, 30
 Prenylamine 1, 76
 Pridefine 3, 49
 Prilocaine 1, 17
 Primaquine 1, 346
 Primidolol 3, 29
 Primidone 1, 276
 Prizidilol 3, 151
 Probarbital 1, 268
 Probenecid 1, 135
 Probuco 2, 126
 Procainamide 1, 14
 Procaine 1, 9
 Procarbazine 2, 27
 Procaterol 3, 184
 Prochlorperazine 1, 381
 Procinonide 3, 94
 Procyclidine 1, 47
 Prodilidine 1, 305
 Prodolic acid 2, 459
 Progesterone 2, 164
 Proglumide 2, 93
 Proguanil 1, 280
 Prolintane 1, 70
 Promazine 1, 377
 Promethazine 1, 373, 377
 Pronethalol 1, 66
 Propanidid 2, 79
 Propantheline bromide 1, 394
 Proparacaine 1, 11
 Propenzolate 2, 75
 Properidine 1, 299
 Propicillin 1, 410
 Propiomazine 1, 376
 Propionylpromazine 1, 380
 Propizepine 2, 472
 Propoxorphan 3, 113
 Propoxycaine 1, 10
 Propoxyphene 1, 50; 298; 2, 57
 Propranolol 1, 117; 2, 105, 107, 212
 Propylhexedrine 1, 37
 Propylphenazone 1, 234
 Propylthiouracil 1, 265
 Proquazone 2, 386
 Proquinolate 2, 368
 Prorenone 2, 175
 Prostalene 2, 5
 Prothipendyl 1, 430
 Prontosil 1, 121
 Protriptyline 1, 152
 Proxazole 2, 271
 Pyrantel 1, 266; 2, 303
 Pyrathiazine 1, 373
 Pyrazineamide 1, 277
 Pyriline 1, 51
 Pyrimethamine 1, 262
 Pyrindamine 1, 145
 Pyrinoline 2, 34
 Pyrovalerone 2, 124
 Pyroxamine 2, 42
 Pyroxiam 2, 394
 Pyrrobutamine 1, 78
 Pyrrocaine 1, 16
 Pyrroliphen 2, 57
 Pyrroxan 3, 191

- Quazepam 3, 196
 Quazodine 2, 379
 Quinacrine 1, 396
 Quinazosin 2, 382
 Quinbolone 2, 154
 Quindonium Bromide 2, 139
 Quinethazone 1, 354
 Quinfamide 3, 186
 Quinidine 1, 339
 Quinine 1, 337
 Quinterenol 2, 366

 Racemoramide 1, 82
 Racemorphan 1, 293
 Ranitidine 3, 131
 Reproterol 3, 231
 Rescinnamine 1, 319
 Rimantadine 2, 19
 Rimiterol 2, 278
 Ripazepam 3, 234
 Risocaine 2, 91
 Ritodrine 2, 39
 Rodocaine 2, 450
 Roletamide 2, 103
 Rolicyprine 2, 50
 Rolitetracycline 1, 216
 Rolodine 2, 468
 Ronidazole 2, 245
 Ripitoin 3, 139
 Rosoxacin 3, 185
 Rotaxamine 2, 32

 Salicylamide 1, 109
 Salbutamol 2, 280
 Salsalate 2, 90
 Salvarsan 1, 223
 Sarmoxicillin 3, 205
 Sarpicillin 3, 204
 Secobarbital 1, 269
 Semustine 2, 12, 15
 Sermetacin 3, 166
 Solypertine 2, 342
 Sontoquine 1, 344
 Sorbinil 3, 188
 Sotalol 1, 66; 2, 41
 Soterenol 2, 40

 Spirilene 2, 292
 Spironolactone 1, 206; 2, 172;
3, 91
 Spiropiperone 1, 306
 Spirorenone 3, 91, 92
 Spirothiobarbital 1, 276
 Stanazole 1, 174
 Stenbolone acetate 2, 155
 Styramate 1, 219
 Succinyl sulfathiazole 1, 132
 Sudoxicam 2, 394
 Sulazepam 2, 403
 Sulconazole 3, 133
 Sulfabenzamide 2, 112
 Sulfacarbamide 1, 123
 Sulfacetamide 1, 123
 Sulfachloropyridazine 1, 124,
 131
 Sulfacytine 2, 113
 Sulfadiazine 1, 124
 Sulfadimethoxine 1, 125, 129
 Sulfadimidine 1, 125, 128
 Sulfathidole 1, 125
 Sulfaguanidine 1, 123
 Sulfaisodimidine 1, 125, 129
 Sulfalene 1, 125
 Sulfamerazine 1, 124, 128
 Sulfameter 1, 125, 129
 Sulfamethizole 1, 125
 Sulfamethoxypyridazine 1, 124,
 131
 Sulfamoxole 1, 124
 Sulfanilamide 1, 121; 2, 112
 Sulfanitrane 2, 115
 Sulfaphenazole 1, 124
 Sulfaproxyline 1, 123
 Sulfapyridine 1, 124; 2, 114
 Sulfasalazine 2, 114
 Sulfasomizole 1, 124
 Sulfathiazole 1, 124
 Sulfathiourea 1, 123
 Sulfazamet 2, 113
 Sulfentanil 3, 118
 Sulfinalol 3, 25
 Sulfinyprazole 1, 238
 Sulfisoxazole 1, 124

- Sulfonterol 2, 42
 Sulformethoxine 1, 125, 130
 Sulfoxone 1, 140
 Sulindac 2, 210
 Sulnidazole 2, 245
 Suloctidil 3, 26
 Sulpiride 2, 94
 Sulprostene 3, 9
 Sulthiame 2, 306
 Suprofen 2, 65
 Symetine 2, 29
 Syrosingopine 1, 319

 Tacamine 2, 224
 Talampicillin 2, 438
 Talniflumate 3, 146
 Talopram 2, 357
 Tametraline 3, 68
 Tamoxifen 2, 127; 3, 70
 Tandamine 2, 347, 460
 Tazolol 2, 110, 268
 Teclozan 2, 28
 Tegafur 3, 155
 Temazepam 2, 402
 Terazocin 3, 194
 Terconazole 3, 137
 Terolidine 2, 56
 Tesicam 2, 379
 Tesimide 2, 296
 Testolactone 2, 160
 Testosterone cypionate 1, 172
 Testosterone decanoate 1, 172
 Testosterone propionate 1, 172
 Tetrabenazine 1, 350
 Tetracaine 1, 110
 Tetracycline 1, 212
 Tetrahydrocannabinol, Δ -1, 1, 394
 Tetrahydrozoline 1, 242
 Tetramisole 1, 431; 3, 226
 Tetrantoin 1, 246
 Tetroxoprim 3, 154
 Tetrydamine 2, 352
 Thalidomide 1, 257; 2, 296
 Thenium closylate 2, 99
 Theobromine 1, 423; 2, 456
 Theophylline 1, 423; 2, 464; 3, 230
 Thiabendazole 1, 325; 2, 352
 Thiabutazide 1, 358
 Thialbarbital 1, 275
 Thiamphenicol 2, 45
 Thiamprine 2, 464
 Thiamylal 1, 274
 Thiazinium chloride 3, 240
 Thiazolsulfone 1, 141
 Thiethylperazine 1, 382
 Thiobarbital 1, 275
 Thiofuradene 1, 231
 Thioguanine 2, 464
 Thiopental 1, 274
 Thiopropazate 1, 383
 Thioridazine 1, 389
 Thiothixene 1, 400; 2, 412
 Thonzylamine 1, 52
 Thozalinone 2, 265
 Thyromedan 2, 79
 Thyroxine 1, 95; 2, 78
 Tiaconazole 3, 133
 Tiamenidine 3, 137
 Tibolone 2, 147
 Tibric acid 2, 87
 Ticarcillin 2, 437
 Ticlopidine 3, 228
 Ticrynafene 2, 104
 Tigesterol 2, 145
 Tiletamine 2, 15
 Tilorone 2, 219
 Timolol 2, 272
 Tiodazocin 3, 194
 Tioperidone 3, 192
 Tiopinac 3, 238
 Tioxidazole 3, 179
 Tipropidil 3, 28
 Tiquinamide 2, 372
 Tixanox 3, 236
 Tocainide 3, 55
 Tofenacin 2, 32
 Tolamolol 2, 110
 Tolazamide 1, 137
 Tolazoline 1, 241; 2, 106
 Tolbutamide 1, 136; 3, 62

- Tolciclate 3, 69
 Tolimidone 3, 156
 Tolindate 2, 208
 Tolmetin 2, 234
 Tolnaftate 2, 211; 3, 69
 Tolpyrramide 2, 116
 Tolycaine 1, 17
 Tonazocine 3, 115
 Topterone 3, 88
 Tosifen 3, 62
 Tralonide 2, 198
 Tramadol 2, 17
 Tramazoline 1, 243
 Tranexamic acid 2, 9
 Tranylcypromine 1, 73; 2, 7, 50
 Trazodone 2, 472
 Treloxinate 2, 432
 Triafungin 3, 233
 Triamcinolone 1, 201; 2, 185
 Triamcinolone acetonide 1, 201
 Triampyzine 2, 298
 Triamterine 1, 427
 Triazolam 1, 368
 Triazuryl 2, 305
 Tricetamide 2, 94
 Trichlormethiazide 1, 359
 Triclonide 2, 198
 Triflocin 2, 282
 Triflubazam 2, 406
 Triflumidate 2, 98
 Trifluoperidol 1, 306
 Triflupromazine 1, 380
 Trihexyphenidyl 1, 47
 Triiodothyronine 1, 95
 Trilostane 2, 158
 Trimazosin 2, 382
 Trimeprazine 1, 378
 Trimethadone 1, 232
 Trimethobenzamide 1, 110
 Trimethoprim 1, 262; 2, 302
 Trimethoquinol 2, 374
 Trimetozine 2, 94
 Trimoxamine 2, 49
 Trioxifene 3, 70
 Trioxsalen 1, 334
 Tripelennamine 1, 51
 Triprolidine 1, 78
 Trofosfamide 3, 161
 Tropocaine 1, 7
 Tybamate 2, 22
 Verilopam 3, 121
 Verofylline 2, 230
 Viloxazine 2, 306; 3, 32
 Vinbarbital 1, 269
 Volazocine 2, 327
 Warfarin 1, 331
 Xanoxate 3, 235
 Xilobam 3, 56
 Xipamide 2, 93
 Xylamidine 2, 54
 Xylazine 2, 307
 Xylometazoline 1, 242
 Zidomethacin 3, 166
 Zimeldine 3, 49
 Zolamine 1, 52
 Zolterine 2, 301
 Zometapine 3, 234
 Zompirac 3, 128

Index

- acclomethasone dipropionate, 96
- acne, 12, 148
- actodigin, 99
- acyclovir, 229
- addiction, 111
- adinazolam, 197
- adrenergic agonist, 20, 23, 30, 40, 184
- adrenergic blocking agents, 20, 24-30, 151, 191, 196
- D-alanine, 13
- L-alanine, 13
- aldose reductase, 72, 188
- aldosterone, 91
- alfentanyl, 118
- alkylcupate reagent, 111
- allergic mediators, 58
- allergic mediations, 66
- alprazolam, 197
- alprostadiol, 2,4,5
- alrestatin, 72
- Alzheimer's disease, 127
- amacetam, 127
- amantadine, 45
- ametantrone, 75
- amfenac, 38
- amidinocillin, 208
- aminal, 192
- amitriptyline, 32
- amitriptylene, 49, 77
- amoxycillin, 205
- ampicillin, 204
- amrinone, 147
- anagrelide, 244
- angiotensin converting enzyme, 128
- anilopam, 121
- anitrazafen, 158
- antagonists, androgen, 81, 248
- antagonists, histamine, 131, 134
- antifertility activity, 70
- arachidonic acid, 10
- arbaprostil, 8
- arildone, 45
- astemizole, 177
- atherosclerosis, 15
- atropine, 160
- azaclozine, 241
- azarole, 129
- azastene, 89
- azepindole, 242
- azipramine, 246
- azlocillin, 206

- azoconazole, 137
azosemide, 27
- bacampicillin, 204
bamnidazole, 132
Bayer-Villiger cleavage, 103, 105
bentazepam, 235
bentiromide, 60
benzomorphans, 114
benzylpenicillin, 203
benzyne, 69
bepiridil, 46
bevantolol, 28
bezafibrate, 44
bicifadine, 120
biocides, 76
bioisostere, see isosteric replacement
biotransformation, 167
bitolterol, 22
 α -blockers, see adrenergic antagonists
 β -blockers, see adrenergic antagonists
 β -blockers, synthesis, 27
blood-brain barrier, 46
blood lipids, 15
breast tumors, 70
brofloxine, 191
broperamole, 139
bucindolol, 28
budesonide, 95
butamisolol, 226
buterizine, 175
butoconazole, 134
butopamine, 23
- calcifediol, 101, 102
calcitriol, 101, 103
calcium channel blockers, 46, 149
calcium metabolism, 101
cannabinoid synthesis, 187, 189
captopril, 128
- carbantel, 57
carbaprost, 7
carbazeran, 195
cardiac glycosides, 99
carfentanyl, 117
caries, dental, 157
caroxazone, 191
carprofen, 169
carteolol, 183
cataracts, 72, 188
cefaparole, 212
cefatrizine, 211
cefazafur, 213
cefonicid, 213
ceforanide, 214
cefotaxime, 216
cefotiam, 215
cefroxadine, 210
cefsulodin, 214
ceftazidime, 216
ceftizoxime, 218
cefuroxime, 216
cell walls, bacterial, 13
cephachlor, 209
cetaben, 60
cetiedil, 42
chlorpromazine, 72
chymotrypsin, 60
cimetidine, 134
cinepazet, 157
cinromide, 44
ciprefadol, 119
ciprocinonide, 94
ciprofibrate, 44
ciramadol, 122, 123
clodanolene, 130
clofibrate, 15, 44
clofilium phosphate, 46
clomacran, 73
clonidine, 40
cloperone, 150
clonipazam, 237
closantel, 43
codorphone, 112
cognition enhancers, 127

- colterol, 21
Corey lactone, 2, 7
cromoglycate, 66, 235
cyclindole, 168
cyclobenzaprine, 77
cyclooxygenase, 39, 110, 165
cyclophosphamide, 161
cycloserine, 14
- declenperone, 172
diacetolol, 28
diazepam, 195
diethylstilbestrol, 50
diflaltone, 246
Digitalis, 99, 147
digitoxin, 99
dihydropyridine synthesis, 185
dihydropyridines, 149
diltiazem, 198
dipirefrin, 22
disobutamide, 41
disopyramide, 41
DNA, disruption of, 83
docanazole, 133
domperidone, 174
dothiepin, 239
doxipicomine, 122, 123
drindene, 65
drobuline, 47
droprenylamine, 47
droxacin, 185
drug absorption, 22
drugs, site directed, 83
- elfazepam, 195
emilium tosylate, 47
encainide, 56
endralazine, 232
enviroxime, 177
epinephrine, 19
epirazole, 152
epoprostenol, 10
ergolines, 249
esterases, 94, 146
estramustine, 83
- ethacrynic acid, 87
etinfidine, 135
etomidate, 135
etoprine, 153
exapralol, 28
- felsantel, 57
fenbendazole, 176
fenclofenac, 37
fendosal, 170
fenobam, 136
fenquizone, 192
fentanyl, 116
flecainide, 59
floctafenine, 184
flucindole, 168
flucinolone, 94
fludalanine, 14
flumequine, 186
flumizole, 158
flumoxonide, 95
fluproquazone, 193
fluquazone, 193
fluretofen, 39
flutamide, 57
5-fluorouracil, 155
fluotracer, 73, 74
fluoxetine, 32
flutroline, 242
fosazepam, 195
foxglove, 99
frentizole, 179
- gemcadiol, 15
gemfibrozil, 45
gestodene, 85
gestrinone, 85
glicetanile, 61
gliflumide, 61
glycosidation, 100
Gould-Jacobs reaction, 184
guanfacine, 40
- halofantrine, 76
halopemide, 174

- haloprednone, 99
helminths, 57
 hormone antagonists, 50
 histamine, antagonists, 131
 homologation, 138

ifosfamide, 151
imafen, 226
imipramine, 32
indacrinone, 67
indomethacin, 165
indoprofen, 171
indorenate, 167
indoxole, 158
 inhibitors, mechanism based, 14
 insulin, 61
ipexidine, 157
ipratropium bromide, 160
isamoxole, 138
isoproterenol, 20, 21
 isosteric replacement, 9, 42, 45, 58, 59, 61, 118, 128, 137, 145, 169, 178, 183, 185, 192, 234
isotretinoin, 12
isoxepac, 238

 Janssen laboratories, 116, 172

ketanserine, 193
ketoconazole, 132
ketotifen, 239

labetolol, 24
levonandrodo, 188
leucotrienes, 1
lidamide, 56
lidocaine, 40
liver flukes, 43
lodoxamide, 57
lofentanyl, 117
lorcainide, 40
lormetazepam, 196

mecillinam, 208

meclorisonone butyrate, 95
medroxoalol, 25
 Meerwein reagent, 71, 119
meobentine, 45
meperidine, 116
mesecclazone, 245
 metabolic inactivation, 9, 21
 metabolic protection, 8, 9
 metabolic transformation, 39, 55, 101
meteneprost, 9
metioprim, 155
metoprine, 153
mezlocillin, 206
midazolam, 197
migrane, 247
milenperone, 172
minaxalone, 90
misonidazole, 132
mitoxantrone, 75
moisidomine, 140
morniflumate, 146
morphinans, 115
morphine, 110
motretinide, 12
moxalactam, 218
moxazocine, 114
muzolimine, 137

nabilone, 189
nabitan, 190
nafoxidine, 70
naftidine, 72
nantradol, 186
napactidine, 71
 nervous system, sympathetic, 20
 Newman thiophenol synthesis, 236
nicarbidine, 150
nicordanil, 148
nifedipine, 149
nimodipine, 149
nisobutanol, 23
nisoxetine, 32

- nistremine acetate, 88
nitrafudam, 130
nitrogen mustards, 82
nitromifene, 51
nivimdone, 67
nocodazole, 176
norepinephrine, 19, 24
norgestrel, 84
nonsteroidal antiinflammatory agents, 37
nufenoxole, 42

oral contraceptives, 84
orconazole, 133
ornidazole, 131
orpanoxin, 130
oxarbazole, 169
oxatomide, 173
oxetorene, 247
oxidation, microbiological, 93
oxiramide, 40
oxmetidine, 134
oxolinic acid, 185
oxyfungin, 233
ozolinone, 140

pain, 109
pamatolol, 28
parconazole, 133
pattern recognition, 42
penicillin V, 205
pentomone, 248
pergolide, 249
pharmacokinetics, 99
phenacetamide, 44
phenoxymethylpenicillin, 205
phenylephrine, 20
piperacillin, 207
pirazolac, 138
pirbencilin, 207
pirenperone, 231
piretanide, 58
piridicillin, 207
pirmentol, 48
piroctone, 149

pirogilide, 57
pirolate, 245
pirquinozol, 243
Polonovski rearrangement, 196
prazosin, 194
prednimustine, 83, 93
premature senility, 127
prenalterol, 30
prenylamine, 47
pridefine, 49
primidolol, 29
prizidilol, 151
procaterol, 184
procinonide, 94
prodrug, 39, 94, 146, 155, 166, 187, 204, 209
propoxorphan, 113
prostacyclin, 10
prostaglandin F_{2α}, 11
pyroxan, 191

quazepam, 196
quinfamide, 186

ranitidine, 131
reduction, microbiological, 6
reproterol, 231
rickets, 101
ripazepam, 234
Robinson annulation, 85
Robinson, tropine synthesis, 160
ropitoin, 139
rosoxacin, 185
Rule, Beckett and Casey, 121

Sarmoxicillin, 205
sarpicillin, 204
screening, random, 75
sermetacin, 166
serotonin, 167
sorbinil, 188
spironolactone, 91
spirorenone, 91, 92
SRS-A, 1

- stereospecificity of drug
 action, 30
 stereospecific synthesis, 3,
 6, 9, 30, 103, 219
sulconazole, 133
sulfentany, 118
sulfinalol, 25
suloctidil, 26
sulprostene, 9
sydnone, 140
 sympathomimetic agents, see
 adrenergic agents
 synthesis, Grewe, 114
 synthesis, stereospecific, 103,
 219
- talniflumate, 146
tametriline, 68
tamoxifen, 51, 70
tegafur, 155
terazocin, 194
terconazole, 137
testosterone, 88
tetramisole, 226
tetroxoprim, 154
thebaine, 111
theophylline, 230
thiazinium chloride, 240
thrombosis, 10
thromboxane A₂, 10
tiaconazole, 133
tiamenidine, 137
ticlopidine, 228
tiodazocin, 194
tioperidone, 192
tiopinac, 238
tioxidazole, 179
tipropidil, 28
tixanox, 236
tocainide, 55
tolbutamide, 62
tolciclate, 69
tolimidone, 156
tolnaftate, 69
tonazocine, 115
- topterone, 88
tosifen, 62
triafungin, 233
trioxifene, 70
trofosfamide, 161
- umpohlung, 6
uric acid, retention, 67
- vasodilator, peripheral, 26
verilopam, 121
verofylline, 230
Villsmeyer reaction, 156
viloxazine, 32
vitamin D, 101
- xanoxate, 235
xilobam, 56
- zidomethacin, 166
zimeldine, 49
zometapine, 234
zompirac, 128