

ED-381-FRM
REV 1 05.10.21

Conway Medical Center
Conway, South Carolina

Center Patient Admission
Label Here

PHYSICIAN'S ORDERS
For IV Nitroglycerin (Tridil)
ER/ICU/PCU and CCL Only

Used in treatment of angina, CHF with acute MI, hypertension.

Another brand of drug identical in form
and content may be dispensed unless checked. ☐

Drug Allergies:

Concentration: 25mg Nitroglycerin in 250ml bottle of D5W (100 microgram/ml)

1. Discontinue any topical Nitroglycerin at time of initiating IV infusion
2. VS upon initiation and with all titrations.
3. VS after titration may be Q15 minutes X 2 hrs, then Q 30 minutes X 2 hrs, then Q 1 hr if stable.
4. Initial dose begins 5 microgram/min.
5. Titration instructions: 5 microgram/min Q 3 minutes; Max dose 200mcg/min.
6. Goal:
 - ☐ Resolved chest pain
 - ☐ SBP< _____ and/or DBP<_____

*****Notify MD of unrelieved chest pain and further titration at 200 micrograms/min*****

CONTRAINDICATED Known hypersensitivity to nitroglycerin, nitrates, increased ICP, constrictive pericarditis, pericardial tamponade, severe hypotension, severe anemia. Avoid with use of phosphodiesterase-5 inhibitors.

MD Signature

_____/_____/_____
Date

Time

Nurse # 1 verifying rate

Nurse # 2 verifying rate