

ED-364-FRM
REV 0 03.12.08

Conway Medical Center

Conway, South Carolina

Physician's Orders

ER/ICU/PCU/Cath Lab ONLY

For IV Corlopadol (FENOLDOPAM)

Center Patient Admission
Label Here

Administration in Treatment of patients with normal or borderline renal function at risk of developing acute renal failure after receiving Radiocontrast Media (RCM)

Another brand of drug identical in form and content may be dispensed unless checked ☐

Allergies:

Contraindicated in patients with Sulfites allergy

Patient Weight _____ Creat _____

Indications:

1. Patients with diabetes and a creatinine of greater than 1.5/mg/dl
2. Patients without diabetes who have a creatinine of greater than 1.7mg/dl

Not compatible with Lasix and sodium bicarbonate.

☐ Initiate Corlopadol 10mg/250ml of normal saline (40mcg/ml) at 0.1 mcg/kg/min

Rate _____ Started _____

☐ Continue post procedure at 0.1 mcg/kg/min times _____ hrs.
D/C at _____ a.m./p.m. (May be cut off, does not have to be weaned)

Obtain BP and heart rate every 15 min X 4 after Corlopadol started then every 1 hr until dc'd.

Stop infusion and notify MD for greater than 25% decrease in BP from baseline.

Decrease infusion rate by half and notify MD for systolic BP less than 110.

Contraindications: High intraocular pressure, glaucoma, allergy to sulfites. BP systolic <90

MD Signature

_____/_____/_____
Date

Time

Nurse # 1 verifying rate

Nurse # 2 verifying rate