

CCU-369-FRM
REV 0 04.15.08

Conway Medical Center
Conway, South Carolina
Physician's Orders

Center Patient Admission
Label Here

**For IV Octreotide Acetate Injection
(SANDOSTATIN)**

Treatment for Acute GI Bleed. Actions: Long acting octapeptide. Mimics actions of the natural hormone somatostatin, suppressing secretions of serotonin, gastroenteropancreatic peptides, and growth hormone.

Patient Data:

Patient Weight _____

The following orders will be initiated:

Lab ☐ BMP
☐ FSBS q _____

☐ Initiate Bolus of 50 mcg IV over 3 – 5 minutes

☐ Initiate Bolus of _____ IV over 3 – 5 minutes

Continuous Drip:

Diluent:	Quantity of Octreotide Acetate (SANDOSTATIN)	Final Concentration
<ul style="list-style-type: none">• 238 ml of NSS or D5W• Remove 12 ml from 250ml bag	600 mcg (use 1 vial 500 mcg and 1 vial 100 mcg for mix)	2.5 mcg/ml

Administration Dose

☐ 25 mcg/hr 10 ml/hr
☐ 50 mcg/hr 20 ml/hr

Precautions: Use caution in patients with diabetes; **requirements for insulin, sulfonylureas, and diazoxide may be altered.** Use caution in patients receiving concomitant beta-blockers, propranolol, calcium channel blockers, verapamil, or any agents used for fluid and electrolyte imbalance. Markedly increases adsorption of insulin and reduces availability. Concurrent use with oral antidiabetic agents, glucagons, growth hormone, or insulin may cause hypoglycemia or hyperglycemia.

MD Signature

_____/_____/_____
Date

Time

Nurse # 1 verifying rate

Nurse # 2 verifying rate