ED-353-FRM REV 1 01.07.14

Conway Medical Center Conway, South Carolina

PHYSICIAN'S ORDERS ER/ICU/PACU/Cath Lab ONLY Center Patient Admission Label Here

For IV Propofol (Diprivan) Administration in Ventilator Support Sedation Only

Another brand of drug identical in form and content may be dispensed unless checked.		Drug Allergies:	
RN can never administer Diprivan boluses		(* Contraindicated in patients with soybean oil, egg, lecithin, glycerol allergies)	
Patient Weight:			
Additional order sets:			
Activity	☐ Specialty Bed		
Treatment/ Nursing	Titrate by 5 mcg/kg/min q 5-10 minutes until desired effects. (Onset peak effects = 5 min.) a. Controlled Respiratory efforts by rate and tidal volume b. Controlled stress response for continuous sedation for intubated, mechanically ventilated adults c. Slow titration with BP monitoring with every change and q 5 min. until stable to minimize hypotensive response d. Documentation of response for desired effects using the Modified Ramsey Scale: Modified Ramsey Scale 1. Patient anxious, agitated, or restless 2. Patient cooperative, oriented and tranquil 3. Patient responds to commands only 4. Patient responds to gentle shaking 5. Patient responds to noxious stimulus 6. Patient has no response to firm nailbed pressure or other noxious stimuli M.D. modified Ramsey Scale orders for Scale Maintenance dose 5 - 55 mcg/kg/min. Notify MD if dose required increase above 55 mcg/kg/min. Triglyceride levels on morning of 3 rd day of infusion Scheduled Date Perform q 24 hr. Wake-Up between 0800-1000 q AM utilizing the Modified Ramsey Scale and Neuro Assessment, document in the Nursing Notes assessment.		
Consults	☐ Dietary Consult		
Medication	☐ Initiate Propofol drip at 5 mcg/kg/min. ☑ VTE Prophylaxis – Physician must complete Form # 395		
Physician Signature Nurse #1 Signature		// Date /	Time
Nurse #2 Signature		// Date	Time