ED-379-FRM REV 2 05.10.21

Conway Medical Center Conway, South Carolina

PHYSICIAN'S ORDERS For IV Nitroprusside (NIPRIDE) ER/ICU Only

Center Patient Admission Label Here

quickly	fusion of Nitroprusside, the hemod . Uses include immediate relief of yy be used for patients with severe n	hypertensive crisis, used f		
Another brand of drug identical in form and content may be dispensed unless checked.			Drug Allergies:	
Initial	weight at infusion start	then daily weights		
	Concentration: 50	0mg Nitroprusside/500ml	of D5W = 100mcg/ml (0.3	Img/ml)
1.	Central line preferred or u MEDICATIONS – Line is to be			VITH OTHER
2.	VS upon initiation of infusion and with all titrations.			
3.	VS after titration may be Q15 minutes X 2 hrs, then Q30 min. X 2 hrs, then Q1 hr if stable			
4.	Initiate at 0.3 microgram/kg/min.			
5.	Titrate instructions: 0.1 microgram/kg/min Q 3min; Max dose 10 mcg/kg/min (At max dose, patient is at risk for cyanide toxicity. Do not infuse at max dose for greater than 10 minutes.)			
6.	Goal: SBP or MAP			
	PRUSSIDE decomposes on exposu			
	ution is stable for 24 hours when p			_
	AINDICATED Sudden, severe di EMOGLOBINEMIA. Renal Impa			
			e consecue 1 2021 are crossing con a	
			//	
	MD Signature		Date	Time
Nurse # 1	verifying rate		Nurse # 2 verifying rate	