Fields, Forms and Flows 3/34

Problem Sheet 7

Due: Wednesday 28 November

To hand in: FFF3: 1, 2, 3(a) FFF34: 1, 2, 3(a)

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only. Problems marked * are not examinable (but are still worth trying!)

1. Determine whether s should be +1 or -1 in the formula

$$f^{(3)} \wedge f^{(8)} \wedge f^{(6)} \wedge f^{(4)} \wedge f^{(1)} = sf^{(1)} \wedge f^{(3)} \wedge f^{(4)} \wedge f^{(6)} \wedge f^{(8)}.$$

(Suggestion: effect the required permutation by a sequence of transpositions between adjacent factors $f^{(i_j)}$ and $f^{(i_{j+1})}$. That is, start by moving $f^{(1)}$ on the left-hand side to the "front", as follows:

$$f^{(3)} \wedge f^{(8)} \wedge f^{(6)} \wedge f^{(4)} \wedge f^{(1)} = -f^{(3)} \wedge f^{(8)} \wedge f^{(6)} \wedge f^{(1)} \wedge f^{(4)} = \\ = +f^{(3)} \wedge f^{(8)} \wedge f^{(1)} \wedge f^{(6)} \wedge f^{(4)} = \cdots$$

Once $f^{(1)}$ is "where it belongs", move $f^{(3)}$ to the left in a similar fashion to where it belongs. Keep track of sign changes.)

2. Let $a=a_if^{(i)}$ be an algebraic 1-form on \mathbb{R}^n and $b=\frac{1}{2}b_{jk}f^{(j)}\wedge f^{(k)}$ be an algebraic 2-form on \mathbb{R}^n with

$$a_i = \begin{cases} \alpha, & i = 1, \\ 0, & \text{otherwise} \end{cases}, \quad b_{jk} = \begin{cases} \beta, & j = 2, k = 3, \\ -\beta, & j = 3, k = 2, \\ 0, & \text{otherwise.} \end{cases}$$

Let $c = a \wedge b$. From the component formula for the wedge product,

$$c = \frac{1}{2!} a_i b_{jk} f^{(i)} \wedge f^{(j)} \wedge f^{(k)}. \tag{1}$$

We also have that

$$c = \frac{1}{3!} c_{ijk} f^{(i)} \wedge f^{(j)} \wedge f^{(k)}, \tag{2}$$

where c_{ijk} is defined in the usual way, ie

$$c_{ijk} = c(e_{(i)}, e_{(j)}, e_{(k)}).$$

Show that

$$c_{123} = a_1 b_{23}$$

but that

$$c_{231} \neq a_2 b_{31}$$

so that, in general, $c_{ijk} \neq a_i b_{jk}$. Explain why Eqs. (1) and (2) are nevertheless compatible.

3. (a) Let a be a nonvanishing (n-1)-form on \mathbb{R}^n . Find (n-1) linearly independent algebraic 1-forms $g^{(1)}, g^{(2)}, \ldots, g^{(n-1)}$ on \mathbb{R}^n such that $g^{(j)} \wedge a = 0$. (Hint: show that $f^{(j)} \wedge a$ cannot vanish for at least one j. WLOG, assume j = n. $g^{(j)}$ can be taken to be an appropriate linear combination of $f^{(j)}$ and $f^{(n)}$, whose coefficients will depend on the components of a.) Hence show that

$$a = \alpha q^{(1)} \wedge q^{(2)} \wedge \dots \wedge q^{(n-1)}$$

for some $\alpha \in \mathbb{R}$, and give a formula for α in terms of the components of a. (For n=3, this is equivalent to showing that an arbitrary vector \mathbf{u} can be expressed as the cross-product $\mathbf{v} \times \mathbf{w}$ of two vectors \mathbf{v} and \mathbf{w} .)

(b) Suppose that a, b, c and d are linearly independent algebraic 1-forms on \mathbb{R}^n (hence $n \geq 4$). Let

$$r = a \wedge b + c \wedge d$$
.

Show that $r \wedge r \neq 0$. Hence show that one cannot find one-forms $g^{(1)}$ and $g^{(2)}$ for which $r = g^{(1)} \wedge g^{(2)}$.

- 4. Suppose that $v = v^1 e_{(1)} + v^2 e_{(2)} + v^3 e_{(3)} \in \mathbb{R}^3$.
 - (a) Let $a = a_1 f^{(1)} + a_2 f^{(2)} + a_3 f^{(3)}$ be a one-form on \mathbb{R}^3 . Compute $i_v a$. Associating v and a to conventional vectors on \mathbb{R}^3 and $i_v a$ to a scalar, what operation in vector algebra does $i_v a$ correspond to?
 - (b) Let $b = b_{12}f^{(1)} \wedge f^{(2)} + b_{23}f^{(2)} \wedge f^{(3)} + b_{31}f^{(3)} \wedge f^{(1)}$ be a two-form on \mathbb{R}^3 . Associating v, b and $i_v b$ to conventional vectors on \mathbb{R}^3 , what operation in vector algebra does $i_v b$ correspond to?
 - (c) Let $\rho = \rho_{123} f^{(1)} \wedge f^{(2)} \wedge f^{(3)}$ be a three-form on \mathbb{R}^3 . Associating v and $i_v \rho$ to conventional vectors on \mathbb{R}^3 , show that these vectors are related by scalar multiplication.