

## Fields, Forms and Flows 3/34

### Solution Sheet 1

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only.

1. i) For  $F(x) = A \cdot x$  to be 1-1, we require that  $x_1 = x_2$  whenever  $A \cdot x_1 = A \cdot x_2$ . Since the map is linear, is equivalent to requiring that  $x = 0$  whenever  $A \cdot x = 0$ . This condition holds if and only if the  $m$  columns of  $A$  are linearly independent. For this to be true, it is necessary (but not sufficient) that  $m \leq n$  (ie,  $A$  can't have more columns than rows). As example,  $F$  is 1-1 for

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 2 \end{pmatrix} \quad \text{but not for} \quad A = \begin{pmatrix} 1 & 2 \\ 1 & 2 \\ 1 & 2 \end{pmatrix}.$$

One can express the condition also by saying that  $F(x)$  is 1-1 if the *rank* of the matrix  $A$  is  $m$ .

- ii) For  $F(x) = A \cdot x$  to be onto, we require that, for any  $y \in \mathbb{R}^n$ , the equation  $y = A \cdot x$  always has at least one solution  $x$  ( $x$  needn't be unique). This condition holds if and only if the  $m$  columns of  $A$  span  $\mathbb{R}^n$ . For this to be true, it is necessary (but not sufficient) that  $m \geq n$ . (ie,  $A$  must have at least as many columns as rows). As example,  $F$  is onto for

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \end{pmatrix} \quad \text{but not for} \quad A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{pmatrix}.$$

One can express the condition also by saying that  $F(x)$  is onto if the *rank* of the matrix  $A$  is  $n$ .

2. We'll construct the map from  $\mathbb{R}^2$  to  $\mathbb{R}$  in 3 stages. First, we construct an invertible map  $f$  from the interval  $(0, 1)$  to  $\mathbb{R}$ . Next, using  $f$ , we'll construct an invertible map  $G$  from  $\mathbb{R}^2 = \{(u, v)\}$  to  $U = \{(u, v) | 0 < u, v < 1\}$  ( $U$  is the open unit square). Then we'll construct a map  $H$  from  $U$  to  $(0, 1)$ . Our required map  $F : \mathbb{R}^2 \rightarrow \mathbb{R}$  may then be taken to be

$$F = f \circ H \circ G.$$

If  $H$  is 1-1, then  $F$  is 1-1, and if  $H$  is invertible, then  $F$  is invertible with inverse

$$F^{-1} = G^{-1} \circ H^{-1} \circ f^{-1}.$$

First, we'll construct  $H$ , and therefore  $F$ , which is 1-1.

- (a) Step 1. Take  $f : (0, 1) \rightarrow \mathbb{R}$  to be

$$f(x) = \ln \frac{x}{1-x},$$

for example. We can show  $f$  is invertible by writing down the inverse explicitly,  $f^{-1} : \mathbb{R} \rightarrow (0, 1)$ , which is given by

$$f^{-1}(y) = \frac{e^y}{1 + e^y}.$$

- (b) Step 2. Define  $G : \mathbb{R}^2 \rightarrow U$  by

$$G(u, v) = (f^{-1}(u), f^{-1}(v)).$$

Then  $G^{-1} : U \rightarrow \mathbb{R}^2$  is given by  $G^{-1}(r, s) = (f(r), f(s))$ .

(c) Step 3. Given  $(u, v) \in U$ , so that  $0 < u, v < 1$ , we can write  $u$  and  $v$  as decimals,

$$u = 0.u_1u_2\cdots, \quad v = 0.v_1v_2\cdots,$$

where  $u_j$  and  $v_j$  are sequences of decimal digits between 0 and 9. Decimal expansions aren't always unique; for example,  $.200\cdots = .1999\cdots$ . However, if we specify that neither  $u_j$  nor  $v_j$  can end with an infinite sequence of consecutive 9's, then  $u_j$  and  $v_j$  are uniquely determined by  $u$  and  $v$ .

We define  $H : U \rightarrow (0, 1); (u, v) \mapsto H(u, v)$  by interleaving the decimal expansions of  $u$  and  $v$  to obtain a single decimal, as follows:

$$H(u, v) = 0.u_1v_1u_2v_2\cdots$$

It is clear that  $H$  is 1-1, ie  $H(u, v) = H(u', v')$  implies that  $u = u'$  and  $v = v'$ . However,  $H$  is not onto, since, for example,  $x = .x_19x_39x_59\cdots$  is not contained in its image.

An invertible map from  $\mathbb{R}^2$  to  $\mathbb{R}$  can also be constructed. We'll just give a sketch of one possible construction. From the preceding, it suffices to construct a mapping from the open unit square  $U$  to the open unit interval  $I = (0, 1)$ .

- (a) Let  $U^*$  denote the set of points  $(u, v) \in U$  with  $u$  and  $v$  both irrational, and similarly let  $I^*$  denote the irrational points in  $I$ . An irrational number  $u$  between 0 and 1 can be uniquely expressed as a continued fraction,

$$u = \frac{1}{a_1 + \frac{1}{a_2 + \cdots}},$$

where  $a_1, a_2, \dots$  is an infinite sequence of positive integers. We write  $u = [a_1, a_2, \dots]$ , and similarly  $v = [b_1, b_2, \dots]$ . Then define  $Q : U^* \rightarrow I^*$  by  $P(u, v) = x = [a_1, b_1, a_2, b_2, \dots]$ .  $Q$  is invertible.

- (b) We also need to construct invertible maps  $P : U \rightarrow U^*$  and  $R : I \rightarrow I^*$ . For  $R$ , note that the interval  $I$  can be divided into the algebraic numbers  $\mathcal{A}$ , ie numbers which are roots of a polynomial with integer coefficients, and the transcendental numbers  $\mathcal{T}$ , which are not the roots of such polynomials. Algebraic numbers include the rational numbers  $\mathbb{Q}$ , and both  $\mathbb{Q}$  and  $\mathcal{A}$  are countable. Hence there is an invertible map – let's call it  $r$  – from  $\mathcal{A}$  to  $\mathcal{A} - \mathbb{Q}$  (ie, the irrational algebraic numbers). Define  $R$  by  $R(x) = x$  if  $x \in \mathcal{T}$  and  $R(x) = r(x)$  if  $x \notin \mathcal{T}$ .  $R$  is then an invertible map from  $I$  to  $I^*$ , and  $P$  can be taken to be  $P(u, v) = (R(u), R(v))$ .

3. Take  $x \in U$ . Then  $F(x) \in F(U)$ , so that  $x \in F^{-1}(F(U))$ , as required. Next, take  $y \in F(F^{-1}(V))$ . Then  $y = F(x)$  for some  $x \in F^{-1}(V)$ . But  $x \in F^{-1}(V)$  implies that  $F(x) \in V$ . Since  $y = F(x)$ , it follows that  $y \in V$ , as required. Let  $F(x) = x^2$ . Let  $U = (0, 1)$ . Then  $F(U) = (0, 1)$ , while  $F^{-1}(F(U)) = (-1, 0) \cup (0, 1)$ . Let  $V = (-1, 0]$ . Then  $F^{-1}(V) = \{0\}$ , so that  $F(F^{-1}(V)) = \{0\}$ .
4. Given  $y \in B_\epsilon(x)$ . We must show that there exists a  $\delta > 0$  such that  $B_\delta(y)$  is contained in  $B_\epsilon(x)$  (in other words, any  $z$  within a distance  $\delta$  of  $y$  is within a distance  $\epsilon$  of  $x$ ). In fact, any positive  $\delta$  with

$$\delta < \epsilon - \|y - x\|$$

will work. For suppose  $z \in B_\delta(y)$ . Then

$$\|z - x\| = \|(z - y) + (y - x)\| \leq \|z - y\| + \|y - x\|,$$

using the triangle inequality. But  $\|z - y\| < \delta$  and  $\|y - x\| < \epsilon - \delta$ , so  $\|z - x\| < \epsilon$ , as required.

5. (a) Let  $U_\alpha$  be a family of open sets in  $\mathbb{R}^m$ . The index  $\alpha$  could range over a finite set, a countable infinite set (eg, the natural numbers) or an uncountable set (eg, the real numbers). Let

$$V = \bigcup_{\alpha} U_{\alpha},$$

where the union is taken over all  $\alpha$ . We must show that  $V$  is open, ie, for all  $x \in V$  there exists an  $\epsilon > 0$  such that  $B_\epsilon(x)$  is contained in  $V$ .

Let  $x \in V$ . Then  $x \in U_\alpha$  for some  $\alpha$ . Since  $U_\alpha$  is open,  $\exists \epsilon > 0$  such that  $B_\epsilon(x) \subset U_\alpha$ . Since  $U_\alpha$  is contained in  $V$ , it follows that  $B_\epsilon(x) \subset V$ , as required.

- (b) Let  $U_1$  and  $U_2$  be open sets in  $\mathbb{R}^m$ . Let

$$V = U_1 \cap U_2.$$

We must show that  $V$  is open.

If  $V$  is empty, there is nothing to show; the empty set is open. If  $V$  is not empty, let  $x \in V$ . Then  $x \in U_1$  and  $x \in U_2$ . Since  $U_1$  is open, there  $\exists \epsilon_1 > 0$  such that  $B_{\epsilon_1}(x) \subset U_1$ . Similarly, since  $U_2$  is open,  $\exists \epsilon_2 > 0$  such that  $B_{\epsilon_2}(x) \subset U_2$ . Let  $\epsilon$  be the smaller of  $\epsilon_1$  and  $\epsilon_2$ . Then  $B_\epsilon(x)$  is contained in both  $U_1$  and  $U_2$ , so  $B_\epsilon(x) \subset V$ , as required.

Note that this argument extends to the intersection of a finite collection of open sets. However, the intersection of an infinite collection of open sets need not be open. For example, the intersection of the open intervals  $I_n = (-1/n, 1/n)$  over all positive integers  $n$  contains only 0, and therefore is not open.

- (c) Let  $U \subset \mathbb{R}$  be open. Given  $x \in U$ , choose  $\delta(x) > 0$  such that  $B_{\delta(x)}(x) \subset U$ . Then

$$U = \bigcup_x B_{\delta(x)}(x),$$

for it is clear that every point in  $U$  is contained in at least one of the intervals on the right-hand side (namely, the interval around that point), while the intervals on the right-hand side are contained in  $U$  by assumption.

6. First, suppose that  $X$  is closed. This means that  $X$  contains all its boundary points. Let  $x \in \tilde{X}$ . Then  $x \notin X$ , so that  $x$  is not a boundary point of  $X$ . This means there exists an  $\epsilon > 0$  such that  $B_\epsilon(x) \cap X$  is empty. Then  $B_\epsilon(x) \subset \tilde{X}$ . But this shows that  $\tilde{X}$  contains a neighbourhood of each of its points, so that  $\tilde{X}$  is open.

Next, suppose that  $\tilde{X}$  is open. We want to show that  $X$  is closed, i.e. that  $X$  contains each of its boundary points. Let  $x$  be a boundary point of  $X$ . Then for all  $\epsilon > 0$ ,  $B_\epsilon(x) \cap X$  is not empty. This means that, for all  $\epsilon > 0$ ,  $B_\epsilon(x)$  is not wholly contained in  $\tilde{X}$ . Thus, since  $\tilde{X}$  is open, we may conclude that  $x \notin \tilde{X}$ , which means that  $x \in X$ .

7. First, suppose that  $F$  is continuous. Let  $Y \subset V$  be open. We wish to show that  $X := F^{-1}(Y)$  is open. Therefore, given  $x \in X$ , we must find  $\delta > 0$  such that  $B_\delta(x) \subset X$ . Here is how: Let  $y = F(x)$ . Since  $x \in X$ , it follows that  $y \in Y$  (this is just what it means for  $x$  to belong to the inverse image of  $Y$ ). Since  $Y$  is open,  $\exists \epsilon > 0$  such that  $B_\epsilon(y) \subset Y$ . Now we use the fact the  $F$  is continuous: Since  $F$  is continuous, it follows that  $\exists \delta > 0$  such that  $\|x' - x\| < \delta$  implies that  $\|F(x') - y\| < \epsilon$ . Another way to say this is that if  $x' \in B_\delta(x)$ , then  $F(x') \in B_\epsilon(y)$ . But this is just saying that  $B_\delta(x)$  is contained in  $F^{-1}(B_\epsilon(y))$ , which in turn is contained in  $F^{-1}(Y) = X$ . This is what we wanted to show.

Next, suppose that the inverse image of every open set  $Y$  in  $V$  is an open set  $X$  in  $U$ . We wish to show that for all  $x \in X$  and for all  $\epsilon > 0$ , there exists  $\delta > 0$  such that  $\|x' - x\| < \delta$  implies that  $\|F(x') - F(x)\| < \epsilon$ , or, in other words,  $F(B_\delta(x)) \subset B_\epsilon(F(x))$ . Here is how. Let  $y = F(x)$ . Since  $Y$  is open,  $\exists \epsilon > 0$  such that  $B_\epsilon(y) \subset Y$ . By assumption, since  $B_\epsilon(y)$  is open, then  $F^{-1}(B_\epsilon)$  is open. Clearly  $x \in F^{-1}(B_\epsilon)$ . Therefore,  $\exists \delta > 0$  such that  $B_\delta(x) \subset F^{-1}(B_\epsilon)$ . But this implies that  $F(B_\delta(x)) \subset B_\epsilon(F(x))$ , which is what we wanted to show.

8. i) Let  $A = (\frac{1}{2}, \frac{3}{2})$ , for example. Then  $f^{-1}(A) = [0, \infty)$ , which is not open. ii) Let  $B = (-\frac{1}{2}, \frac{1}{2})$ . Then  $g^{-1}(B)$  is not open. In particular, while  $0 \in g^{-1}(B)$  (since  $g(0) = 0$ , and  $0 \in (-\frac{1}{2}, \frac{1}{2})$ ), no  $\delta$ -neighbourhood of 0 is contained in  $g^{-1}(B)$ . This is because, arbitrarily close to 0, there are points - let's call them  $x_n$  - for which, say,  $|\sin 1/x_n| = 1$ . Just let  $x_n = 1/((n + \frac{1}{2})\pi)$ .
9. Let  $f(x) = 1$  for all  $x$ . Then the image of any set, open or otherwise, is  $\{1\}$ , which is closed.