

# Fields, Forms and Flows 3/34

## Problem Sheet 5

Due: Wednesday 14 November (2 weeks)

To hand in: FFF3: 1, 2    FFF34: 1, 2

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only. Problems marked \* are not examinable (but are still worth trying!)

1. An  $n$ th order ODE can be reduced to a system of  $n$  first-order ODE's (we had examples in Problem Sheet 2). In a similar way, an  $n$ th order PDE can be reduced to a system of first-order PDE's. You might wonder whether such a system can be solved using the Frobenius theorem (provided the necessary condition is satisfied), but, as the following examples show, this is generally not possible.

- (a) Consider the one-dimensional wave equation,

$$u_{tt} - u_{xx} = 0,$$

where  $u_{tt} = \partial^2 u / \partial t^2$ , etc. Write the wave equation as a system of first-order partial differential equations for  $\phi(x, t) = (\phi^1(x, t), \phi^2(x, t), \phi^3(x, t))$ , where  $\phi^1 = u$ ,  $\phi^2 = u_x$  and  $\phi^3 = u_t$ . Explain why this system cannot be expressed in the form  $\phi_x^\alpha = f^\alpha(x, t, \phi)$ ,  $\phi_t^\alpha = g^\alpha(x, t, \phi)$ .

- (b) Same as the preceding question but for the one-dimensional heat equation,

$$u_t - u_{xx} = 0.$$

2. Consider the system of coupled first-order nonlinear partial differential equations,

$$\begin{aligned} \frac{\partial u}{\partial x} &= f^1(x, y, u, v), & \frac{\partial v}{\partial x} &= f^2(x, y, u, v), \\ \frac{\partial u}{\partial y} &= g^1(x, y, u, v), & \frac{\partial v}{\partial y} &= g^2(x, y, u, v), \end{aligned}$$

with initial data

$$u(0, 0) = r_0, \quad v(0, 0) = s_0.$$

- (a) Derive necessary and sufficient conditions on  $f$  and  $g$  in order for this system to have a solution, and give an expression for  $u$  and  $v$  in terms of the flows of two vector fields  $X$  and  $Y$  on  $\mathbb{R}^4$ .

(b) For the case

$$f(x, y, r, s) = (a, b),$$

where  $a$  and  $b$  are constants, show that the condition is satisfied if and only if  $g$  is of the form

$$g(x, y, r, s) = (G^1(y, r - ax, s - bx), G^2(y, r - ax, s - bx)),$$

where  $G^1$  and  $G^2$  are arbitrary functions of three variables. (Remark: If you find the “only if” part difficult just show the “if” part.)

(c) For

$$G^1 = -G^2 = (r - ax)(s - bx)$$

and  $r_0 = s_0 = 1$ , find  $u(x, y)$  and  $v(x, y)$  explicitly in a neighbourhood of  $(0, 0)$ .

3. Consider the following system of coupled first-order nonlinear partial differential equations for the scalar-valued function  $u = u(x, y)$ , which is a generalization of the system we considered in Section 1.11.2 of the Notes:

$$\begin{aligned} a \frac{\partial u}{\partial x} + b \frac{\partial u}{\partial y} &= f(x, y, u), \\ c \frac{\partial u}{\partial x} + d \frac{\partial u}{\partial y} &= g(x, y, u). \end{aligned} \quad (*)$$

Here  $a, b, c$  and  $d$  are given smooth functions of  $x$  and  $y$  and we assume that

$$ad - bc \neq 0 \text{ for all } (x, y) \in \mathbb{R}^2.$$

Show that the system has a unique solution for all initial data  $u(x_0, y_0) = u_0$  if and only if

$$[\mathbb{V}, \mathbb{W}] = r\mathbb{V} + s\mathbb{W}, \quad (**)$$

where  $\mathbb{V}$  and  $\mathbb{W}$  are the vector fields on  $\mathbb{R}^3$  given by

$$\mathbb{V}(x, y, z) = (a(x, y), b(x, y), f(x, y, z)), \quad \mathbb{W}(x, y, z) = (c(x, y), d(x, y), g(x, y, z)),$$

for some functions  $r(x, y, z)$  and  $s(x, y, z)$ . ((\*\*) is equivalent to saying the Jacobi bracket of  $\mathbb{W}$  and  $\mathbb{V}$  lies in the plane spanned by  $\mathbb{W}$  and  $\mathbb{V}$ .) (Suggestion: show that the problem can be reduced to one to which the Frobenius theorem applies. A more general version of this result is discussed in Section 1.12 of the notes, which you can consult, but you can also answer this question without referring to Section 1.12.)