

Fields, Forms and Flows 3/34

Solution Sheet 10

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only.

1. Consider the singular two-cube in \mathbb{R}^2 ,

$$c : [0, 1]^2 \rightarrow \mathbb{R}^2; \quad c(x^1, x^2) = (x^1 x^2, x^2).$$

- (a) The image of c is the right triangle with vertices $(0, 0)$, $(0, 1)$ and $(1, 1)$.
 (b) First, let's compute the faces of c . We have that

$$\begin{aligned} c_{(1,0)}(x) &= c(0, x) = (0, x), \\ c_{(1,1)}(x) &= c(1, x) = (x, x), \\ c_{(2,0)}(x) &= c(x, 0) = (0, 0), \\ c_{(2,1)}(x) &= c(x, 1) = (x, 1). \end{aligned}$$

Then

$$\partial c = -c_{(1,0)} + c_{(1,1)} + c_{(2,0)} - c_{(2,1)}.$$

To evaluate $\partial(\partial c)$, we first compute the boundaries of the faces. We have that

$$\begin{aligned} \partial c_{(1,0)}(x) &= c_{(1,0)}(1) - c_{(1,0)}(0) = (0, 1) - (0, 0), \\ \partial c_{(1,1)}(x) &= c_{(1,1)}(1) - c_{(1,1)}(0) = (1, 1) - (0, 0), \\ \partial c_{(2,0)}(x) &= c_{(2,0)}(1) - c_{(2,0)}(0) = (0, 0) - (0, 0) = 0, \\ \partial c_{(2,1)}(x) &= c_{(2,1)}(1) - c_{(2,1)}(0) = (1, 1) - (0, 1). \end{aligned}$$

Add up the terms on the right-hand side with the appropriate signs to get

$$\partial(\partial c) = -\partial c_{(1,0)} + \partial c_{(1,1)} + \partial c_{(2,0)} - \partial c_{(2,1)} = +((0, 0) - (0, 1)) - (0, 0) + (1, 1) + ((0, 1) - (1, 1)) = 0.$$

- (c) Let

$$\omega = \frac{1}{2}(y^1 dy^2 - y^2 dy^1).$$

First, let's compute

$$\int_c d\omega = \int_{[0,1]^2} c^* d\omega.$$

We have that

$$d\omega = dy^1 \wedge dy^2.$$

Then

$$c^* d\omega = \left(\frac{\partial y^1}{\partial x^1} \frac{\partial y^2}{\partial x^2} - \frac{\partial y^1}{\partial x^2} \frac{\partial y^2}{\partial x^1} \right) dx^1 \wedge dx^2 = x^2 dx^1 \wedge dx^2.$$

This is equivalent to noting that $c^* y^1 = x^1 x^2$ and $c^* y^2 = x^2$, so that $c^* dy^1 = x^1 dx^2 + x^2 dx^1$ and $c^* dy^2 = dx^2$, and therefore

$$c^* \omega = (x^1 dx^2 + x^2 dx^1) \wedge dx^2 = x^2 dx^1 \wedge dx^2.$$

It follows that

$$\int_c d\omega = \int_0^1 \int_0^1 x^2 dx^1 dx^2 = \frac{1}{2}$$

Next we compute

$$\int_{\partial c} \omega = \sum_{j=1}^2 \sum_{\alpha=0,1} (-1)^{j+\alpha} \int_{[0,1]} c_{(j,\alpha)}^* \omega.$$

From above,

$$(c_{(2,1)}^* \omega)(x) = \frac{1}{2} \left(y^1(x) \frac{\partial y^2}{\partial x}(x) - y^2(x) \frac{\partial y^1}{\partial x}(x) \right) dx,$$

where

$$(y^1(x), y^2(x)) = c_{(2,1)}(x) = (x, 1).$$

Therefore,

$$(c_{(2,1)}^* \omega)(x) = -\frac{1}{2} dx.$$

Similar calculations give

$$(c_{(1,0)}^* \omega)(x) = (c_{(1,1)}^* \omega)(x) = (c_{(2,0)}^* \omega)(x) = 0.$$

Therefore,

$$\int_{\partial c} \omega = - \int_{[0,1]} c_{(2,1)}^* \omega = \int_0^1 \frac{1}{2} dx = \frac{1}{2}.$$

Thus, $\int_c d\omega = \int_{\partial c} \omega$, in accord with Stokes' theorem.

2. (a) Let $\mathbf{r}(t)$, $0 \leq t \leq 1$, denote a smooth parameterised curve and $F(\mathbf{r}) = F_x dx + F_y dy + F_z dz$ a one-form on \mathbb{R}^3 . Then

$$(\mathbf{r}^* F)(t) = \left(F_x(\mathbf{r}(t)) \frac{dx}{dt}(t) + F_y(\mathbf{r}(t)) \frac{dy}{dt}(t) + F_z(\mathbf{r}(t)) \frac{dz}{dt}(t) \right) dt = \mathbf{F}(\mathbf{r}(t)) \cdot \dot{\mathbf{r}}(t) dt,$$

where $\mathbf{F} = (F_x, F_y, F_z)$.

- (b) Let $\mathbf{S}(u, v)$, $0 \leq u, v \leq 1$, denote a smooth parameterised surface and $B(\mathbf{r}) = B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy$ a two-form on \mathbb{R}^3 . Then

$$\begin{aligned} (\mathbf{S}^* B)(u, v) &= B_x(\mathbf{S}(u, v)) \left(\frac{\partial y}{\partial u} \frac{\partial z}{\partial v} - \frac{\partial y}{\partial v} \frac{\partial z}{\partial u} \right) du \wedge dv \\ &\quad + B_y(\mathbf{S}(u, v)) \left(\frac{\partial z}{\partial u} \frac{\partial x}{\partial v} - \frac{\partial z}{\partial v} \frac{\partial x}{\partial u} \right) du \wedge dv \\ &\quad + B_z(\mathbf{S}(u, v)) \left(\frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u} \right) du \wedge dv. \end{aligned}$$

Let

$$\mathbf{N} = \frac{\partial \mathbf{S}}{\partial u} \times \frac{\partial \mathbf{S}}{\partial v}.$$

Then

$$\begin{aligned} N_x &= \frac{\partial y}{\partial u} \frac{\partial z}{\partial v} - \frac{\partial y}{\partial v} \frac{\partial z}{\partial u}, \\ N_y &= \frac{\partial z}{\partial u} \frac{\partial x}{\partial v} - \frac{\partial z}{\partial v} \frac{\partial x}{\partial u}, \\ N_z &= \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}. \end{aligned}$$

Substituting this in the expression above, we can write

$$(\mathbf{S}^* B)(u, v) = \mathbf{B}(\mathbf{S}(u, v)) \cdot \mathbf{N}(u, v) du \wedge dv.$$

- (c) Let $S(u, v)$ be a smooth parameterised surface, as above. Let

$$\begin{aligned} \mathbf{r}_1(t) &= \mathbf{S}_{(2,0)}(t) = \mathbf{S}(t, 0), \\ \mathbf{r}_2(t) &= \mathbf{S}_{(1,1)}(t) = \mathbf{S}(1, t), \\ \mathbf{r}_3(t) &= \mathbf{S}_{(2,1)}(t) = \mathbf{S}(t, 1), \\ \mathbf{r}_4(t) &= \mathbf{S}_{(1,0)}(t) = \mathbf{S}(0, t). \end{aligned}$$

Then

$$\partial S = \mathbf{r}_1 + \mathbf{r}_2 - \mathbf{r}_3 - \mathbf{r}_4.$$

- (d) According to Stokes' theorem,

$$\int_S dF = \int_{\partial S} F.$$

We want to write this in vector notation. The two-form dF corresponds to the vector $\nabla \times \mathbf{F}$, so, using the preceding results, the left-hand side can be written as

$$\int_0^1 \int_0^1 (\nabla \times F)(\mathbf{S}(u, v)) \cdot \mathbf{N}(u, v) du dv.$$

The right-hand side is just

$$\int_0^1 (\mathbf{F}(\mathbf{r}_1(t)) \cdot \dot{\mathbf{r}}_1(t) + \mathbf{F}(\mathbf{r}_2(t)) \cdot \dot{\mathbf{r}}_2(t) - \mathbf{F}(\mathbf{r}_3(t)) \cdot \dot{\mathbf{r}}_3(t) - \mathbf{F}(\mathbf{r}_4(t)) \cdot \dot{\mathbf{r}}_4(t)) dt.$$

3. (a) If ω is a k -form on \mathbb{R}^n with $d\omega = 0$, then, for any singular $(k+1)$ -cube c on \mathbb{R}^n , Stokes' theorem implies that

$$\int_{\partial c} \omega = \int_c d\omega = 0 \quad (\text{since } d\omega = 0).$$

- (b) If ω is a k -form on \mathbb{R}^n and $\omega = d\alpha$ for some $(k-1)$ -form α , then for any singular k -cube c on \mathbb{R}^n with $\partial c = 0$, Stokes' theorem implies that

$$\int_c \omega = \int_c d\alpha = \int_{\partial c} \alpha = 0 \quad (\text{since } \partial c = 0).$$

- (c) Consider the one-form

$$\omega = \frac{ydx - xdy}{x^2 + y^2}$$

on the punctured $\mathbb{R}^2 - \{0\}$. We have that

$$d\omega = \frac{\partial}{\partial y} \left(\frac{y}{x^2 + y^2} \right) dy \wedge dx - \frac{\partial}{\partial x} \left(\frac{x}{x^2 + y^2} \right) dx \wedge dy.$$

Substituting the formulas

$$\frac{\partial}{\partial y} \left(\frac{y}{x^2 + y^2} \right) = \frac{(x^2 + y^2) - 2y^2}{(x^2 + y^2)^2}, \quad \frac{\partial}{\partial x} \left(\frac{x}{x^2 + y^2} \right) = \frac{(x^2 + y^2) - 2x^2}{(x^2 + y^2)^2},$$

into the preceding, we get that

$$d\omega = \left(\frac{(x^2 + y^2) - 2y^2 + (x^2 + y^2) - 2x^2}{(x^2 + y^2)^2} \right) dy \wedge dx = 0.$$

- (d) Let c be the singular one-cube on $\mathbb{R}^2 - \{0\}$ given by

$$c(t) = (\cos(2\pi t), \sin(2\pi t)).$$

Since $c(0) = c(1)$, we have that

$$\partial c = -c(0) + c(1) = 0.$$

We have that

$$\int_c \omega = \int_{[0,1]} c^* \omega.$$

Also,

$$c^* \omega = \left(\frac{ydx/dt - xdy/dt}{x^2 + y^2} dt \right) \Big|_{x=\cos 2\pi t, y=\sin 2\pi t} = -2\pi(\sin^2(2\pi t) + \cos^2(2\pi t)) dt = -2\pi dt.$$

Alternative calculation: $c^*x = \cos 2\pi t$, so that $c^*dx = dc^*x = -2\pi \sin 2\pi t dt$. Similarly, $c^*y = \sin 2\pi t$, so that $c^*dy = 2\pi \cos 2\pi t dt$. Therefore,

$$c^* \omega = c^* \left(\frac{y}{x^2 + y^2} \right) c^* dx - c^* \left(\frac{x}{x^2 + y^2} \right) c^* dy = -2\pi dt.$$

Either way, we get that

$$\int_c \omega = - \int_0^1 2\pi dt = -2\pi. \tag{1}$$

We cannot have $\omega = df$ for any function f on $\mathbb{R}^2 - \{0\}$, for this would imply that

$$\int_c \omega = \int_c df = \int_{\partial c} f = 0 \quad (\text{since } \partial c = 0),$$

in contradiction to the preceding. Similarly, we cannot have $c = \partial c_2$ for a singular two-cube c_2 on $\mathbb{R}^2 - \{0\}$, for this would imply that

$$\int_c \omega = \int_{\partial c_2} \omega = \int_{c_2} d\omega = 0 \quad (\text{since } d\omega = 0),$$

in contradiction to the preceding.