

# Fields, Forms and Flows 3/34

## Problem Sheet 6

Due: Wednesday 21 November

To hand in: FFF3: 1, 3(b), 6, 7    FFF34: 1, 3(b), 6, 7

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only. Problems marked \* are not examinable (but are still worth trying!)

1. (a) Let  $e_{(1)}$  and  $e_{(2)}$  be a basis for a two-dimensional vector space  $V$ , and let  $f^{(1)}$  and  $f^{(2)}$  be the associated dual basis on  $V^*$ . Let

$$\bar{e}_{(1)} = e_{(1)}, \quad \bar{e}_{(2)} = e_{(1)} + e_{(2)}.$$

Let  $\bar{f}^{(1)}$  and  $\bar{f}^{(2)}$  denote the dual basis to the  $\bar{e}_{(j)}$ 's. Express  $\bar{f}^{(1)}$  and  $\bar{f}^{(2)}$  as linear combinations of  $f^{(1)}$  and  $f^{(2)}$ .

- (b) Let  $e_{(1)}, \dots, e_{(n)}$  and  $\bar{e}_{(1)}, \dots, \bar{e}_{(n)}$  be two bases for  $V$ , and suppose

$$\bar{e}_{(i)} = \sum_{j=1}^n M_{ij} e_{(j)},$$

where  $M$  is an  $n \times n$  matrix. Let  $f^{(j)}$  and  $\bar{f}^{(j)}$  denote the dual bases of  $e_{(i)}$  and  $\bar{e}_{(i)}$  respectively, and let

$$\bar{f}^{(i)} = \sum_{j=1}^n N_{ij} f^{(j)}.$$

Show that

$$N = M^{T^{-1}}.$$

2. Let  $V$  and  $W$  denote vector spaces of dimensions  $n$  and  $m$ , respectively, and let  $L(V, W)$  denote the set of linear maps from  $V$  to  $W$ .
- (a) Show that  $L(V, W)$  is a vector space.
- (b) Show that  $L(V, W)$  may be identified with the space of functions on  $V \times W^*$  which are linear in each argument.
3. (a) Show that every permutation  $\sigma \in S_n$  can be expressed as a product of transpositions.
- (b) Write

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 6 & 4 & 5 & 1 & 2 \end{pmatrix}$$

as a product of transpositions.

4. Show that

$$\tau_{ij} = \sigma \tau_{12} \sigma^{-1},$$

where  $\sigma$  is any permutation for which  $\sigma(1) = i$  and  $\sigma(2) = j$ .

5. Let  $(i_1, \dots, i_k)$  denote an (ordered)  $k$ -tuple of distinct integers in  $\{1, \dots, N\}$ . Define a permutation  $\sigma \in S_n$  by

$$\begin{aligned} \sigma(i_1) &= i_2, & \sigma(i_2) &= i_3, & \dots, & & \sigma(i_{k-1}) &= i_k, & \sigma(i_k) &= i_1, \\ \sigma(j) &= j & \text{if } j &\neq i_1, \dots, i_k. \end{aligned}$$

$\sigma$  is called a  $k$ -cycle. Show that

$$\operatorname{sgn} \sigma = (-1)^{k-1}.$$

(Note that a transposition is a 2-cycle.)

6. Show that the permutation matrix  $P(\sigma)$  is orthogonal, ie

$$P(\sigma)^{-1} = P(\sigma)^T,$$

where  $P(\sigma)^T$  denotes the transpose of  $P(\sigma)$ . Hence show that  $(\det P(\sigma))^2 = 1$ , so that  $\det P(\sigma) = \pm 1$ . (It then follows that  $\operatorname{sgn} \det P(\sigma) = \det P(\sigma)$ .)

7. Given  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w} \in \mathbb{R}^3$ , let

$$a(\mathbf{u}, \mathbf{v}, \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}.$$

Show that  $a$  is an algebraic 3-form on  $\mathbb{R}^3$ .