

Fields, Forms and Flows 3/34

Problem Sheet 2

Due: Wednesday 17 October

To hand in: DM3: 3, 4, 8, 9 DM34: 6(a), 8, 9

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only. Problems marked * are not examinable (but are still worth trying!)

1. Let $f(x) = ax^2 + bx + c$ be a function on \mathbb{R} , where a, b, c are constants and $a \neq 0$. According to the Inverse Function Theorem, which values of x have a neighbourhood $I_\delta(x)$ on which f is invertible?
2. Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by $(x, y) \mapsto (x^2 - y^2, 2xy)$. Compute F' , and show that $\det F' = 0$ if and only if $x = y = 0$. Compute the inverse of F explicitly, and discuss how the result is consistent with the Inverse Function Theorem.
3. (a) If $f : \mathbb{R} \rightarrow \mathbb{R}$ (say f is smooth) satisfies $f'(a) \neq 0$ for all $a \in \mathbb{R}$, show that f is 1-1 on \mathbb{R} . (You may use Rolle's theorem – if f has a continuous derivative and $f(a) = f(b)$ for some $a \neq b$, then f' must vanish somewhere between a and b). (b) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by $f(x, y) = (e^x \cos y, e^x \sin y)$. Show that $\det f'(x, y) \neq 0$ for all $(x, y) \in \mathbb{R}^2$, but that f is not 1-1.
4. * Let $f(x) = \frac{1}{2}x + x^2 \sin(1/x)$ for $x \neq 0$ and $f(0) = 0$. Show that $f'(0) = \frac{1}{2}$, but that $f'(x)$ is not continuous at $x = 0$. Show that f is not 1-1 in any neighbourhood of 0. (Thus, the Inverse Function Theorem need not hold if f' is not continuous.)
5. * Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth function on \mathbb{R}^2 . Use the Inverse Function Theorem to show that f is not 1-1.
6. (a) Let

$$Y_0 = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$$

and let K be an arbitrary 2×2 matrix. Show that, for ϵ sufficiently small, the matrix $Y_0 + \epsilon K$ has a square root that is close to the matrix

$$X_0 = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}.$$

That is, for sufficiently small ϵ , there exists a matrix $X(\epsilon)$ depending smoothly on ϵ such that $X^2(\epsilon) = Y_0 + \epsilon K$ and $X(0) = X_0$. (Suggestion: identify the space of 2×2 matrices with \mathbb{R}^4 , and apply the inverse function theorem to an appropriately defined map $F : \mathbb{R}^4 \rightarrow \mathbb{R}^4$.)

- (b) Show that, for ϵ small and K an arbitrary 2×2 matrix, the matrix $I + \epsilon K$ does not necessarily have a real square root that is close to the matrix

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

(Suggestion: Let $K = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.) This is in contrast to what happens for 1×1 matrices, ie real numbers. Let $y_0 > 0$, and let $x_0^2 = y_0$ (x_0 could be the positive or negative square root of y_0). Then for ϵ small, $y_0 + \epsilon$ always has a square root that is close to x_0 .

7. Write the third-order nonautonomous ODE

$$\frac{d^3 x}{dt^3} = \sin t \frac{x \ddot{x}}{\dot{x}^2}$$

as a first-order autonomous system on \mathbb{R}^4 .

8. For $q \in \mathbb{R}$, write $\ddot{q} = 1$ as a first-order system, find the flow $\Phi_t : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, and verify that $\Phi_t \circ \Phi_s = \Phi_{t+s}$.
9. Let $\dot{x} = x^2$. Solve this (autonomous) differential equation, and show that the flow $\Phi_t(x_0)$ has a singularity at $t = T$, where T depends on x_0 . Over what interval is the flow defined? Next, let $\dot{x} = x^2/(1+x^2)$. Solve the differential equation, and show that the flow $\Phi_t(x_0)$ is defined for all t .

10. * Let

$$V(q) = \begin{cases} -\frac{3}{4}q^{4/3}, & q > 0, \\ 0, & q \leq 0. \end{cases}$$

Write Newton's second law, $\ddot{q} = -V'(q)$, as a first-order system in $x = (q, \dot{q})$. Show that $q(t) = 0$ is a solution to the equation with initial conditions $q(0) = 0$, $\dot{q}(0) = 0$. Find another solution which satisfies these initial conditions of the form $q(t) = at^b$. Consider more generally the differential equation

$$\ddot{q} = \begin{cases} 0, & q \leq 0, \\ q^\alpha, & q > 0. \end{cases}$$

For which values of $\alpha > 0$ are there two solutions, $q(0)$ and $q = at^b$ (for appropriate a and b), which satisfy the initial conditions $q(0) = \dot{q}(0) = 0$?