

Fields, Forms and Flows 3/34

Problem Sheet 4

Due: Wednesday 31 October

To hand in: FFF3: 1, 3 FFF34: 1, 3

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only. Problems marked * are not examinable (but are still worth trying!)

1. We define the Lie derivative of a vector field \mathbb{Y} with respect to a vector field \mathbb{X} as follows:

$$L_{\mathbb{X}}\mathbb{Y} := [\mathbb{X}, \mathbb{Y}],$$

where \mathbb{X} and \mathbb{Y} are vector fields. (The connection with the Lie derivative on functions is as follows: We have that

$$L_{\mathbb{X}}f = \mathbb{X} \cdot \nabla f = \left. \frac{\partial}{\partial t} \Phi_t^* f \right|_{t=0},$$

where Φ_t is the flow of \mathbb{X} . We therefore define

$$L_{\mathbb{X}}\mathbb{Y} := - \left. \frac{\partial}{\partial t} \Phi_{t*} \mathbb{Y} \right|_{t=0} = -[\mathbb{Y}, \mathbb{X}] = [\mathbb{X}, \mathbb{Y}].$$

The reason for the minus sign in the definition is that the push-forward goes in the opposite direction to the pull-back.) Then

$$L_{\mathbb{X}}^2 \mathbb{Y} = L_{\mathbb{X}}(L_{\mathbb{X}}\mathbb{Y}) = [\mathbb{X}, [\mathbb{X}, \mathbb{Y}]], \quad L_{\mathbb{X}}^3 \mathbb{Y} = [\mathbb{X}, [\mathbb{X}, [\mathbb{X}, \mathbb{Y}]]], \quad \text{etc}$$

(it's the ability to write the nested brackets in a compact way that makes this notation useful here).

- (a) Let Ψ_s be the flow of \mathbb{Y} . Show that


$$\frac{\partial^n}{\partial s^n} \Psi_{s*} \mathbb{X} = (-1)^n L_{\mathbb{Y}}^n \Psi_{s*} \mathbb{X}.$$

(Suggestion: recall the derivation of equation (10) following Definition 1.9.3 in the notes. Use induction.)

- (b) Derive the (formal) power series

$$\Psi_{s*} \mathbb{X} = e^{-sL_{\mathbb{Y}}} \mathbb{X}.$$

(Suggestion: follow the model of Proposition 1.9.4.) Use this expression to show that $\Psi_{s*} \mathbb{Y} = \mathbb{Y}$.



(c) Let \mathbb{Z} be a vector field. Show that the Jacobi identity can be written in the form

$$(L_{\mathbb{X}}L_{\mathbb{Y}} - L_{\mathbb{Y}}L_{\mathbb{X}})\mathbb{Z} = L_{[\mathbb{X}, \mathbb{Y}]}\mathbb{Z}.$$

2. Parallel parking. Consider the model car shown above. The centre of the back axle is at P , with coordinates (x, y) . The direction of the back wheels makes an angle θ with the vertical. The length of the body is L . The centre of the front axle is at Q , and the direction of the front wheels makes an angle $\theta + \alpha$ with the vertical. The configuration of the car is determined by x , y , θ and α , which collectively we regard as a point (x, y, θ, α) in \mathbb{R}^4 (of course, changing θ or α by 2π does not change the configuration of the car).

(a) Let \mathbb{S} be the smooth vector field on \mathbb{R}^4 given by

$$\mathbb{S}(x, y, \theta, \alpha) = (0, 0, 0, \omega),$$

where ω is a constant (\mathbb{S} stands for ‘steer’). Determine the flow Φ_t of \mathbb{S} .

- (b) Determine the changes in x , y and θ if the back wheels roll forward a small distance $v\delta t$ in the direction θ (the front wheels roll forward, too, in the direction $\theta + \alpha$, by a distance you should determine). Show that this infinitesimal change in the configuration is described by the vector field \mathbb{D} given by

$$\mathbb{D}(x, y, \theta, \alpha) = (v \sin \theta, v \cos \theta, (v/L) \tan \alpha, 0),$$

where v is a constant (\mathbb{D} stands for ‘drive’). (Hint: the car can’t stretch! Its length is fixed.) Compute the flow Ψ_s of \mathbb{D} . (If you find the derivation of \mathbb{D} difficult just continue with the computation of its flow.)

- (c) Compute $\Psi_\epsilon(x, y, \theta, \epsilon\Omega)$ up to and including terms of order ϵ^2 . If ϵ is regarded as a small parameter, this represents driving for a short time ϵ with the front wheels only slightly turned by angle $\epsilon\Omega$.
- (d) Compute $\mathbb{A} = [\mathbb{S}, \mathbb{D}]$ and $\mathbb{B} = [[\mathbb{S}, \mathbb{D}], \mathbb{D}]$. Show that for $\cos \alpha \neq 0$, \mathbb{A} , \mathbb{B} , \mathbb{S} and \mathbb{D} are linearly independent. Compute the flows, Γ_a and Δ_b , of \mathbb{A} and \mathbb{B} respectively. (It turns out that Γ_a describes a pure rotation in θ , while Δ_b describes a pure translation of the car perpendicular to its direction.)
- (e) Verify Theorem 1.10.1 by showing explicitly that

$$\begin{aligned} (\Psi_{-\epsilon} \circ \Phi_{-\epsilon} \circ \Psi_\epsilon \circ \Phi_\epsilon)(x, y, \theta, 0) &= \Gamma_{\epsilon^2}(x, y, \theta, 0) + O(\epsilon^3) \\ &= (x, y, \theta, 0) + \epsilon^2 \mathbb{A}(x, y, \theta, 0) + O(\epsilon^3). \end{aligned}$$

Explain how, by a sequence of small steer and drive maneuvers, you can change the orientation θ of your car by as much as you like while changing x , y and α as little as you like.

- (f) Verify Theorem 1.10.1 by showing explicitly that

$$\begin{aligned} (\Psi_{-\delta} \circ \Gamma_{-\delta T} \circ \Psi_\delta \circ \Gamma_{\delta T})(x, y, \theta, 0) &= \Delta_{\delta^2 T}(x, y, \theta, 0) + O(\delta^3) \\ &= (x, y, \theta, 0) + \delta^2 T \mathbb{B}(x, y, \theta, 0) + O(\delta^3). \end{aligned}$$

Here T may be regarded as a fixed constant with dimensions of time (if you are keeping track of dimensions, this makes the expression dimensionally sensible), or else may be set equal to 1 (if you prefer to ignore dimensional considerations). Explain how, by a sequence of small steer and drive maneuvers, you can translate your car in a direction perpendicular to its length by as much as you like while changing θ and α as little as you like. (This is parallel parking into a tight space.)

3. Noncommutativity of rotations in \mathbb{R}^3 .

- (a) Let

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

and let

$$\mathbb{X}_A(r) = A \cdot r, \quad \mathbb{X}_B(r) = B \cdot r, \quad \mathbb{X}_C(r) = C \cdot r$$

be linear vector fields on \mathbb{R}^3 , with $r = (x, y, z)$. Show that

$$[\mathbb{X}_A, \mathbb{X}_B] = -\mathbb{X}_C, \quad [\mathbb{X}_B, \mathbb{X}_C] = -\mathbb{X}_A, \quad [\mathbb{X}_C, \mathbb{X}_A] = -\mathbb{X}_B.$$

(It may be helpful to refer to Problem 3.6.)

- (b) Show that Φ_{Ct} , the flow of \mathbb{X}_C , is given by

$$\Phi_{Ct}(r) = \begin{pmatrix} \cos t & -\sin t & 0 \\ \sin t & \cos t & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

- (c) Derive similar formulas for the flows Φ_{At} and Φ_{Bt} of \mathbb{X}_A and \mathbb{X}_B .
(d) Show explicitly that, for small θ ,

$$(\Phi_{B\theta} \circ \Phi_{A\theta} \circ \Phi_{B(-\theta)} \circ \Phi_{A(-\theta)})(r) = \Phi_{C(-\theta^2)}(r) + O(\theta^3).$$

(A related fact: Hold a bank note in front of you so that the Queen is upright on your right looking at you. Take the z -direction to be pointing up and y to be pointing in front of you. Rotate the bank note about y by -90° (that is, anticlockwise from your perspective), then about z by 180° , then about y by 90° (that is, clockwise), then about z by 180° . The Queen should be looking at you again, but upside-down on your left.)