

Differentiation and Integration

*Please hand in questions 1 and 2(a)-(c) by Thursday 7 March.*

1. Consider calculating  $f'(1)$  for  $f(x) = e^x$  using the forward difference formula

$$f'(x) \approx \frac{f(x+h) - f(x)}{h}.$$

Derive the truncation error incurred by using this formula. Now pretend you can only work to 4 decimal figures and use various  $h = 10^{-n}$  with  $n = 0, 1, \dots, 5$  to generate estimates of  $f'(1)$ . What is the optimal choice of  $n$  which gives greatest accuracy and explain how you can predict this from the sizes of the truncation and round-off errors.

2. You are told that a function  $f(x)$  has the following values

$$\begin{aligned} f(1.0) &= 2.287355, \\ f(1.1) &= 2.677335, \\ f(1.2) &= 3.094479, \\ f(1.3) &= 3.535581, \\ f(1.4) &= 3.996196. \end{aligned}$$

- (a) Using forward difference and central difference formulae numerically approximate the derivative  $f'(1.2)$  with  $h = 0.2$  and  $h = 0.1$ .
- (b) Derive a forward 3-point formula to approximate  $f'(x_0)$  using  $f(x_0)$ ,  $f(x_0 + h)$  and  $f(x_0 + 2h)$  which is accurate to  $O(h^2)$ . Use this to find  $f'(1.2)$ .
- (c) Produce a better estimate of  $f'(1.2)$  by using Richardson's extrapolation for the central difference formula. (The actual value of  $f'(1.2)$  is 4.297549.)
- (d) Use the composite Trapezoidal Rule to estimate the integral

$$\int_1^{1.4} f(x) dx.$$

3. Given  $f(x_0)$ ,  $f'(x_0 + h)$  and  $f(x_0 - \lambda h)$  where  $\lambda = O(1)$  and  $h \ll 1$ , build a formula for estimating  $f''(x_0)$ . What is the error incurred by using this formula. Show that for  $\lambda^2 = 3$ , the error is  $O(h^2)$ .