

## Summary notes for Fields, Forms and Flows 3 and 34

©University of Bristol 2018. This material is copyright of the University unless explicitly stated otherwise. It is provided exclusively for educational purposes at the University and is to be downloaded or copied for your private study only.

These summary notes are intended to present the main results in outline form. They may be helpful for revision, but they are not a substitute for going through the lecture notes. In particular, you can't assume that all you need to know is in these summary notes.

1. **Diffeomorphisms (Section 1.5).** Let  $U$  and  $V$  be open subsets of  $\mathbb{R}^n$ . A smooth map  $F \in C^\infty(U, V)$  is a *diffeomorphism* if  $F$  is invertible with  $F^{-1} \in C^\infty(V, U)$ . Let  $\text{Diff}(U, V)$  denote the set of diffeomorphisms between  $U$  and  $V$ . If  $U$  and  $V$  are the same, we write  $\text{Diff}(U)$  for the set of diffeomorphisms of  $U$  to itself. The set of diffeomorphisms  $\text{Diff}(U)$  on  $U$  forms a group under composition (Proposition 1.5.3).
2. **ODEs, vector fields and flows (Section 1.6).** Let  $\mathbb{X} : U \rightarrow \mathbb{R}^n$  be a smooth vector field on an open set  $U \subset \mathbb{R}^n$ . Then the first-order system

$$\dot{x}(t) = \mathbb{X}(x(t)), \quad x(0) = x_0,$$

has a unique solution  $x(t, x_0)$  for  $-T < t < T$  for some  $T > 0$  (which may depend on  $x_0$ ) (Theorem 1.6.3). As a function of initial conditions,  $x(t, x_0)$  is smooth (Theorem 1.6.7). A vector field  $\mathbb{X}$  is *complete* if  $x(t, x_0)$  is defined for all  $t$  and  $x_0 \in U$ . Suppose  $\mathbb{X}$  is smooth and complete. We define a map

$$\Phi : \mathbb{R} \times U \rightarrow U; (t, x_0) \mapsto \Phi_t(x_0) = x(t, x_0).$$

$\Phi$  is called the *flow* of the vector field  $\mathbb{X}$ .  $\Phi$  has the following properties (Proposition 1.6.9):

- (a)  $\Phi_0 = \text{Id}_U$ .
- (b)  $\Phi_t \circ \Phi_s = \Phi_{t+s}$ .
- (c)  $\Phi_t : U \rightarrow \Phi_t(U)$  is a diffeomorphism.
- (d)  $\Phi \in C^\infty(\mathbb{R} \times U, \mathbb{R}^n)$ .

A map with properties (a)–(d) is called a *one-parameter subgroup of diffeomorphisms*.

Conversely, if  $\Phi$  is a one-parameter subgroup of diffeomorphisms on  $U \subset \mathbb{R}^n$ , we define a vector field  $\mathbb{X}$  on  $U$  by

$$\mathbb{X}(x) = \left. \frac{\partial}{\partial t} \right|_{t=0} \Phi_t(x).$$

Then  $\Phi$  is the flow of  $\mathbb{X}$  (Proposition 1.6.12).

**Matrix exponential (Examples 1.6.10, 1.6.13.)** Let  $A \in \mathbb{R}^{n \times n}$  be an  $n \times n$  matrix. Define

$$e^{tA} = (I + tA + \tfrac{1}{2}t^2A^2 + \cdots) = \sum_{j=0}^{\infty} \frac{t^j A^j}{j!}.$$

**Proposition 1.6.9** says that

$$e^{tA} e^{sA} = e^{(s+t)A}.$$

**Proposition 1.6.12** says that

$$\left. \frac{d}{dt} \right| e^{tA} = A.$$

3. **Pushforward of vector fields (Section 1.7)** Let  $U \subset \mathbb{R}^n$  be open, and let  $\mathcal{X}(U)$  denote the set of smooth vector fields on  $U$ . Furthermore, let  $V \subset \mathbb{R}^n$  be open, and let  $F \in \text{Diff}(U, V)$ . We define a map  $F_* : \mathcal{X}(U) \rightarrow \mathcal{X}(V) : \mathbb{X} \mapsto F_*\mathbb{X}$ , by either of the following equivalent formulas:

$$\begin{aligned} F_*\mathbb{X}(y) &= F'(F^{-1}(y)) \cdot \mathbb{X}(F^{-1}(y)), \\ F_*\mathbb{X}(F(x)) &= F'(x) \cdot \mathbb{X}(x), \end{aligned} \tag{1}$$

or without arguments

$$F_*\mathbb{X} = (F' \cdot \mathbb{X}) \circ F^{-1} \quad \text{or} \quad F_*\mathbb{X} \circ F = F' \cdot \mathbb{X}.$$

$F_*\mathbb{X}$ , a smooth vector field on  $V$ , is called the *pushforward* of  $\mathbb{X}$  by  $F$ . The definition (1) is motivated by changing variables in the system of ODE's described by  $\mathbb{X}$ . That is, if

$$\dot{x} = \mathbb{X}(x)$$

and we define  $y(t) = F(x(t))$  and  $\mathbb{Y} = F_*\mathbb{X}$ , then  $y(t)$  satisfies the system

$$\dot{y} = \mathbb{Y}(y).$$

At the level of flows, we have the following (**Proposition 1.7.3**): Let  $\Phi_t$  be the flow of  $\mathbb{X}$ . Then  $\Psi_t$ , the flow of  $F_*\mathbb{X}$ , is given by

$$\Psi_t = F \circ \Phi_t \circ F^{-1}$$

For a linear vector field  $\mathbb{X}(x) = Ax$ , where  $A \in \mathbb{R}^{n \times n}$ , and a linear diffeomorphism  $F(x) = Sx$ , where  $S \in \mathbb{R}^{n \times n}$  and  $S$  is invertible,

$$(F_*\mathbb{X})(y) = SAS^{-1}y.$$

(See Example 1.7.4).

The pushforward by a composition of two maps is given by **Proposition 1.7.5**: Let  $F \in \text{Diff}(U, V)$ ,  $G \in \text{Diff}(V, W)$ , and  $\mathbb{X} \in \mathcal{X}(U)$  be a smooth vector field. Then

$$(G \circ F)_*\mathbb{X} = G_*F_*\mathbb{X}.$$

4. **Jacobi bracket. Commuting flows (Section 1.8).** Let  $U \subset \mathbb{R}^n$  be open and let  $\mathbb{X}, \mathbb{Y} \in \mathcal{X}(U)$ . The *Jacobi bracket* of  $\mathbb{X}$  and  $\mathbb{Y}$ , denoted  $[\mathbb{X}, \mathbb{Y}]$ , is the vector field in  $\mathcal{X}(U)$  given by

$$[\mathbb{X}, \mathbb{Y}] = (\mathbb{X} \cdot \nabla)\mathbb{Y} - (\mathbb{Y} \cdot \nabla)\mathbb{X}.$$

Here,  $\nabla = (\partial/\partial x_1, \dots, \partial/\partial x_n)$ . Let  $\Psi_s$  be the flow of  $\mathbb{Y}$ . Then (**Proposition 1.8.2**)

$$[\mathbb{X}, \mathbb{Y}] = \left. \frac{\partial}{\partial s} \right|_{s=0} \Psi_{s*}\mathbb{X}.$$

The Jacobi bracket has the following properties (**Proposition 1.8.5**):

- (a) Linearity.  $[a\mathbb{X} + b\mathbb{Y}, \mathbb{Z}] = a[\mathbb{X}, \mathbb{Z}] + b[\mathbb{Y}, \mathbb{Z}]$ , where  $a, b \in \mathbb{R}$ .
- (b) Antisymmetry.  $[\mathbb{X}, \mathbb{Y}] = -[\mathbb{Y}, \mathbb{X}]$ .
- (c) Product rule.  $[\mathbb{X}, f\mathbb{Y}] = f[\mathbb{X}, \mathbb{Y}] + (\mathbb{X} \cdot \nabla f)\mathbb{Y}$ , where  $f : U \rightarrow \mathbb{R}$  is a smooth function.

**Proposition 1.8.6:** The pushforward of the Jacobi bracket is given by

$$F_*[\mathbb{X}, \mathbb{Y}] = [F_*\mathbb{X}, F_*\mathbb{Y}].$$

**Proposition 1.8.8:** Let  $\mathbb{Z} \in \mathcal{X}(U)$  be a third vector field. The Jacobi identity states that

$$[[\mathbb{X}, \mathbb{Y}], \mathbb{Z}] + [[\mathbb{Y}, \mathbb{Z}], \mathbb{X}] + [[\mathbb{Z}, \mathbb{X}], \mathbb{Y}] = 0.$$

**Proposition 1.8.10:** The derivative of the pushforward by a flow at arbitrary time. Let  $U \subset \mathbb{R}^n$  be open, let  $\mathbb{X}, \mathbb{Y} \in \mathcal{X}(U)$  and let  $\Psi_s$  denote the flow of  $\mathbb{Y}$ . Then

$$\frac{\partial}{\partial s} \Psi_{s*} \mathbb{X} = [\Psi_{s*} \mathbb{X}, \mathbb{Y}].$$

**Proposition 1.8.11:** A vector field is invariant under pushforward by its flow. Let  $\mathbb{X} \in \mathcal{X}(U)$  be a vector field with flow  $\Phi_t$ . Then

$$\Phi_{t*} \mathbb{X} = \mathbb{X}.$$

**Theorem 1.8.12:** Commutativity of flows and Jacobi bracket. Let  $\mathbb{X}$  and  $\mathbb{Y}$  be smooth vector fields on an open set  $U \subset \mathbb{R}^n$  with flows  $\Phi_t$  and  $\Psi_s$ , respectively. Then

$$\Phi_t \circ \Psi_s = \Phi_s \circ \Phi_t \iff [\mathbb{X}, \mathbb{Y}] = 0.$$

## 5. Pullback and Lie derivative on smooth functions. Noncommuting flows. (Sections 1.9 and 1.10.)

Let  $U, V \subset \mathbb{R}^n$  be open, and let  $F \in \text{Diff}(U, V)$ . We define a map  $F^* : C^\infty(V) \rightarrow C^\infty(U)$ , called the *pullback by  $F$* , which maps smooth functions on  $V$  into smooth functions on  $U$ . (Later we'll extend the definition to differential forms.) Given  $f \in C^\infty(V)$ , the pullback of  $f$  by  $F$  is defined by

$$F^*f = f \circ F, \quad \text{i.e.} \quad F^*f(x) = f(F(x)).$$

The pullback is a linear map; that is, if  $f, g \in C^\infty(V)$  and  $a, b \in \mathbb{R}$ , then

$$F^*(af + bg) = aF^*f + bF^*g.$$

**Proposition 1.9.2:** The pullback by a composition. Let  $U, V, W \subset \mathbb{R}^n$  be open, and  $F \in \text{Diff}(U, V)$  and  $G \in \text{Diff}(V, W)$ . Then

$$(G \circ F)^* = F^*G^*.$$

Let  $\mathbb{X} \in \mathcal{X}(U)$ . We define a map  $L_{\mathbb{X}} : C^\infty(U) \rightarrow C^\infty(U)$ , called the *Lie derivative with respect to  $\mathbb{X}$* , which maps smooth functions into smooth functions. (Later we will extend the definition to differential forms.) Given  $f \in C^\infty(U)$ , we define the Lie derivative of  $f$  by  $\mathbb{X}$  as

$$L_{\mathbb{X}}f = (\mathbb{X} \cdot \nabla f) = X^i \frac{\partial f}{\partial x^i}.$$

That is,  $L_{\mathbb{X}}f$  is the directional derivative of  $f$  along  $\mathbb{X}$ .

**Proposition 1.9.4:** Let  $\mathbb{X} \in \mathcal{X}(U)$  and let  $\Phi$  be the flow of  $\mathbb{X}$ . Then

$$\Phi_t^*f = \sum_{j=0}^{\infty} \frac{t^j}{j!} L_{\mathbb{X}}^j f = e^{tL_{\mathbb{X}}} f.$$

Here it is assumed that  $\Phi_t^* f(x)$  is analytic in  $t$ . Then the series for  $\Phi_t^* f(x)$  converges in some neighbourhood.

**Proposition 1.9.6** Differential-operator form of Jacobi bracket. Let  $U \subset \mathbb{R}^n$  be open and let  $\mathbb{X}, \mathbb{Y} \in \mathcal{X}(U)$ , and  $f \in C^\infty(U)$ . Then

$$L_{\mathbb{X}}L_{\mathbb{Y}}f - L_{\mathbb{Y}}L_{\mathbb{X}}f = L_{[\mathbb{X}, \mathbb{Y}]}f.$$

**Theorem 1.10.1** Noncommuting flows. Let  $\mathbb{X}$  and  $\mathbb{Y}$  be vector fields with flows  $\Phi_t$  and  $\Psi_s$  respectively, and let

$$\Gamma_{s,t} = \Psi_{-s} \circ \Phi_{-t} \circ \Psi_s \circ \Phi_t.$$

Let  $f$  be a smooth function. Then

$$\Gamma_{s,t}^* f = f + stL_{[\mathbb{X}, \mathbb{Y}]}f + O(3),$$

where  $O(3)$  denotes terms of third and higher order in  $s$  and  $t$ . Equivalently,

$$\Gamma_{s,t}(x) = x + st[\mathbb{X}, \mathbb{Y}](x) + O(3).$$

6. **Frobenius theorem. (Section 1.11.)** Let  $x$  denote coordinates on  $\mathbb{R}^p$  and  $z$  coordinates on  $\mathbb{R}^q$ . Let  $U \subset \mathbb{R}^p$  and  $V \subset \mathbb{R}^q$  be open. Let  $f_i^\alpha$  denote smooth functions on  $U \times V$ ,

$$f_i^\alpha : U \times V \rightarrow \mathbb{R}; \quad (x, z) \mapsto f_i^\alpha(x, z),$$

where  $1 \leq i \leq p$  and  $1 \leq \alpha \leq q$ . Consider the system of first-order partial differential equations for  $u : U \rightarrow V$  given by

$$\begin{aligned} \frac{\partial u^\alpha}{\partial x^i}(x) &= f_i^\alpha(x, u(x)), \\ u(x_0) &= u_0, \quad x_0 \in U, u_0 \in V. \end{aligned} \tag{2}$$

Define  $p$  vector fields  $\mathbb{X}_{(i)}$ ,  $1 \leq i \leq p$ , on  $U \times V$  as follows:

$$\mathbb{X}_{(i)}^j(x, z) = \delta_i^j, 1 \leq j \leq p, \quad \mathbb{X}_{(i)}^{p+\alpha}(x, z) = f_i^\alpha(x, z), 1 \leq \alpha \leq q.$$

That is, among the first  $p$  components of  $\mathbb{X}_{(i)}$ , there is a single nonzero component, namely the  $i$ th, which is equal to one, while the last  $q$  components of  $\mathbb{X}_{(i)}$  are given by  $f_i^1, \dots, f_i^q$ . Suppose the vector fields  $\mathbb{X}_{(i)}$  are complete.

**Frobenius theorem (Theorem 1.11.2.)** For all  $(x_0, u_0) \in U \times V$ , the system (2) has a solution  $u(x)$  if and only if

$$[\mathbb{X}_{(i)}, \mathbb{X}_{(j)}] = 0, \quad 1 \leq i, j \leq p.$$

Moreover, if a solution exists, then it is unique.

**Explicit construction of solutions:** Let  $\Phi_{(i)t}$  denote the flow of  $\mathbb{X}_{(i)}$ . The first  $p$  components of  $\Phi_{(i)t}$  are given by

$$\Phi_{(i)t}^j(x_0) = x_0^j + \delta_i^j t, \quad 1 \leq j \leq p.$$

For simplicity, suppose  $x_0 = 0$  (it is easy to generalise to  $x_0 \neq 0$ ). Then  $u(x)$  is obtained from

$$(x, u(x)) = (\Phi_{(1)x^1} \circ \dots \circ \Phi_{(p)x^p})(0, u_0).$$

## 7. Dual space

**Definition 2.1.1.** Let  $V$  denote an  $n$ -dimensional vector space. The *dual space* of  $V$ , denoted  $V^*$ , is the vector space consisting of *linear functions* on  $V$ .

Let  $e_{(1)}, \dots, e_{(n)}$  denote a basis for  $V$ . We define a set of  $n$  elements of  $V^*$ , denoted  $f^{(1)}, \dots, f^{(n)}$ , by

$$f^{(j)}(e_{(i)}) = \delta_i^j.$$

**Proposition 2.1.3.**  $f^{(1)}, \dots, f^{(n)}$  constitute a basis for  $V^*$ , called the *dual basis*.

Under a linear transformation, vectors in  $V$  and  $V^*$  transform differently. Let  $e_{(1)}, \dots, e_{(n)}$  and  $\bar{e}_{(1)}, \dots, \bar{e}_{(n)}$  be two bases for  $V$ . Then one set of basis vectors can be expressed as linear combinations of the others, e.g.

$$\bar{e}_{(i)} = \sum_{j=1}^n M_{ij} e_{(j)},$$

where  $M$  is an  $n \times n$  matrix. Let  $f^{(j)}$  and  $\bar{f}^{(j)}$  denote the dual bases of  $e_{(i)}$  and  $\bar{e}_{(i)}$  respectively. Here, too, one set of basis vectors can be expressed as linear combinations of the others,

$$\bar{f}^{(i)} = \sum_{j=1}^n N_{ij} f^{(j)}.$$

**Proposition 2.1.4.**

$$N = (M^T)^{-1}.$$

## 8. Permutations

**Proposition 2.2.1.** Every permutation can be written as a product (composition) of transpositions.

**Proposition 2.2.2.** For all  $\sigma, \tau \in S_n$ ,

$$P(\sigma\tau) = P(\sigma)P(\tau),$$

where  $P(\sigma)$  is the permutation matrix with elements  $P_{ij}(\sigma) = \delta_{i, \sigma(j)}$ .

**Definition 2.2.3.** The *sign* of a permutation, denoted  $\text{sgn } \sigma$ , is defined by

$$\text{sgn } \sigma = \text{sgn } \det P(\sigma).$$

**Proposition 2.2.4.** For all  $\sigma, \tau \in S_n$ ,

$$\text{sgn } (\sigma\tau) = \text{sgn } (\sigma) \text{sgn } (\tau).$$

**Proposition 2.2.5.**

$$\text{sgn } (\sigma^{-1}) = \text{sgn } (\sigma).$$

**Proposition 2.2.6.** If  $\tau_{rs}$  is a transposition, then  $\text{sgn } \tau_{rs} = -1$ .

**Proposition 2.2.7.** If  $\sigma$  is a product of  $k$  transpositions, then  $\text{sgn } \sigma = (-1)^k$ .

**Proposition 2.2.8.** Let  $f : S_n \rightarrow \mathbb{R}$  be a function on  $S_n$ . Then for all  $\alpha, \beta \in S_n$ ,

$$\sum_{\sigma \in S_n} f(\sigma) = \sum_{\sigma \in S_n} f(\alpha\sigma\beta).$$

Also,

$$\sum_{\sigma \in S_n} f(\sigma) = \sum_{\sigma \in S_n} f(\sigma^{-1}).$$

## 9. Algebraic $k$ -forms

An *algebraic  $k$ -form on  $V$*  is a function on  $V^k$  that is linear in each argument and which changes sign if two arguments are interchanged. That is, letting  $a$  denote an algebraic  $k$ -form, we have that

$$a : V^k \rightarrow \mathbb{R}; (\mathbf{v}_{(1)}, \dots, \mathbf{v}_{(k)}) \mapsto a(\mathbf{v}_{(1)}, \dots, \mathbf{v}_{(k)}).$$

Linearity with respect to each argument means that, for  $u, w \in V$  and  $\alpha, \beta \in \mathbb{R}$ ,

$$a(\mathbf{v}_{(1)}, \dots, \alpha u + \beta w, \dots, \mathbf{v}_{(k)}) = \alpha a(\mathbf{v}_{(1)}, \dots, u, \dots, \mathbf{v}_{(k)}) + \beta a(\mathbf{v}_{(1)}, \dots, w, \dots, \mathbf{v}_{(k)}).$$

Changing sign under the interchange of two arguments means that, for any  $j, l$  with  $1 \leq j < l \leq k$ ,

$$a(\mathbf{v}_{(1)}, \dots, \mathbf{v}_{(j)}, \dots, \mathbf{v}_{(l)}, \dots, \mathbf{v}_{(k)}) = -a(\mathbf{v}_{(1)}, \dots, \mathbf{v}_{(l)}, \dots, \mathbf{v}_{(j)}, \dots, \mathbf{v}_{(k)}).$$

Denote the set of algebraic  $k$ -forms by  $\Lambda^k(V)$ . By convention,  $\Lambda^0(V)$  is given by  $\mathbb{R}$ . Also,  $\Lambda^1(V)$  is identified with the dual space  $V^*$  (the antisymmetry requirement is empty for  $k = 1$ ).

**Proposition 2.3.1.**  $\Lambda^k(V)$  is a vector space.

**Proposition 2.3.2.** For  $a \in \Lambda^k(V)$  and for  $\sigma \in S_k$ ,

$$a(v_{(\sigma(1))}, \dots, v_{(\sigma(k))}) = \text{sgn } \sigma a(v_{(1)}, \dots, v_{(k)}).$$

Let  $e_{(1)}, \dots, e_{(n)}$  denote a basis for  $V$ . Given  $v \in V$ , we write

$$v = v^i e_{(i)},$$

where we use the summation convention: if an index appears twice on one side of an equation, once as an upper index and once as a lower index, then we sum over that index. We introduce some notation. Let  $I = (i_1, \dots, i_k)$  denote an ordered  $k$ -tuple of indices, where  $1 \leq i_r \leq n$  ( $I$  is also called a multi-index). We introduce a Kronecker delta for pairs of  $k$ -tuples of indices, defined by

$$\delta(I, J) = \begin{cases} 1, & i_1 = j_1, \dots, i_k = j_k, \\ 0, & \text{otherwise.} \end{cases}$$

Given  $\sigma \in S_k$ , define

$$\sigma(I) = (i_{\sigma^{-1}(1)}, \dots, i_{\sigma^{-1}(k)}).$$

That, is  $\sigma(I)$  is a permutation of the indices comprising  $I$ .

**Proposition 2.4.1.** Let  $\sigma, \tau \in S_k$ . Then

$$\sigma(\tau(I)) = (\sigma\tau)(I).$$

Let  $E_{(I)} \in V^k$  denote the  $k$ -tuple of basis vectors given by

$$E_{(I)} = (e_{(i_1)}, \dots, e_{(i_k)}).$$

Given  $a \in \Lambda^k(V)$ , we write

$$a_I = a(E_{(I)}).$$

We will call the  $a_I$ 's the *coefficients* of  $a$  with respect to the basis  $e_{(i)}$ . An alternative notation for the coefficients is

$$a_{i_1 \dots i_k} = a(e_{(i_1)}, \dots, e_{(i_k)}).$$

**Definition 2.4.2.** Let  $J = (j_1, \dots, j_k)$ . The *basis  $k$ -form*  $F^{(J)}$  is the algebraic  $k$ -form on  $V$  defined by

$$F_I^{(J)} := F^{(J)}(E_{(I)}) = \begin{cases} 0, & \text{if } j_r = j_s \text{ for some } r \neq s, \\ \text{sgn } \sigma, & J = \sigma(I), \\ 0, & \text{otherwise.} \end{cases}$$

**Proposition 2.4.3.** Suppose  $J$  consists of distinct indices. Then

$$F^{(J)}(E_{(I)}) = \sum_{\sigma \in S_k} \text{sgn } \sigma \delta(\sigma(I), J).$$

**Proposition 2.4.5.**

$$F^{(J)}(v_{(1)}, \dots, v_{(k)}) = \det \begin{pmatrix} v_{(1)}^{j_1} & v_{(2)}^{j_1} & \dots & v_{(k)}^{j_1} \\ v_{(1)}^{j_2} & v_{(2)}^{j_2} & \dots & v_{(k)}^{j_2} \\ \vdots & \vdots & \dots & \vdots \\ v_{(1)}^{j_k} & v_{(2)}^{j_k} & \dots & v_{(k)}^{j_k} \end{pmatrix}.$$

**Proposition 2.4.7.** For  $\alpha \in S_k$ ,

$$F^{(\alpha(J))} = \text{sgn } \alpha F^{(J)}.$$

**Proposition 2.4.9.** Let  $a \in \Lambda^k(V)$ . Then

$$a = \frac{1}{k!} a_J F^{(J)},$$

where we use the summation convention for  $J$  (that is, there is a sum over each index  $j_r$  in  $J = (j_1, \dots, j_k)$ ).

Let  $J_*$  denote a  $k$ -tuple of indices which are distinct and in ascending order. That is,  $J_* = (j_1, \dots, j_k)$  with  $j_1 < j_2 < \dots < j_k$ .

**Proposition 2.4.10.** The  $F^{(J_*)}$ 's form a basis for  $\Lambda^k(V)$ , and for all  $a \in \Lambda^k(V)$ ,

$$a = \sum_{J_*} a_{J_*} F^{(J_*)}.$$

## 10. Wedge product

**Proposition 2.5.1.** Let  $a \in \Lambda^k(V)$  be an algebraic  $k$ -form and  $b \in \Lambda^l(V)$  be an algebraic  $l$ -form. Their *wedge product*, denoted  $a \wedge b$ , is the algebraic  $(k + l)$ -form defined by

$$a \wedge b(v_{(1)}, \dots, v_{(k+l)}) = \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} \text{sgn } \sigma a(v_{(\sigma(1))}, \dots, v_{(\sigma(k))}) b(v_{(\sigma(k+1))}, \dots, v_{(\sigma(k+l))}). \quad (3)$$

**Proposition 2.5.3.**

i)  $a \wedge b$  is an algebraic  $(k + l)$ -form. That is,  $a \wedge b$ , as defined by (3), is linear in each argument and changes sign under interchange of any pair of arguments.

ii) Linearity. If  $a$  is an algebraic  $k$ -form and  $b$  and  $c$  are algebraic  $l$ -forms, then

$$a \wedge (b + c) = a \wedge b + a \wedge c.$$

iii) (Anti)commutativity. If  $a$  is an algebraic  $k$ -form and  $b$  is an algebraic  $l$ -form, then

$$a \wedge b = (-1)^{kl} b \wedge a.$$

In other words, if either  $k$  or  $l$  is even, then  $a \wedge b = b \wedge a$ . If both  $k$  and  $l$  are odd, then  $a \wedge b = -b \wedge a$ .

iv) Associativity. If  $a$  is an algebraic  $k$ -form,  $b$  an algebraic  $l$ -form, and  $c$  an algebraic  $m$ -form, then

$$a \wedge (b \wedge c) = (a \wedge b) \wedge c.$$

v) Basis  $k$ -forms. Let  $J = (j_1, \dots, j_k)$ . Then

$$F^{(J)} = f^{(j_1)} \wedge \dots \wedge f^{(j_k)}.$$

## 11. Contraction mapping

**Definition 2.7.1.** Let  $v \in V$ . We define  $i_v : \Lambda^k(V) \rightarrow \Lambda^{k-1}(V)$ , the contraction with  $v$ , by

$$i_v c = 0, \text{ for } c \in \Lambda^0(V),$$

$$(i_v a)(w_{(1)}, \dots, w_{(k-1)}) = a(v, w_{(1)}, \dots, w_{(k-1)}), \text{ for } a \in \Lambda^k(V), k > 0,$$

where  $w_{(1)}, \dots, w_{(k-1)} \in V$ . Thus,  $i_v$  map a  $k$ -form  $a$  to a  $(k - 1)$ -form  $i_v a$  by fixing the first argument of the  $k$ -form to be  $v$ . The contraction on any zero-form is defined to be zero.

**Proposition 2.7.2.**

$$i_v(a + b) = i_v a + i_v b, \quad a, b \in \Lambda^k(V),$$

$$i_v(a \wedge b) = (i_v a) \wedge b + (-1)^k a \wedge (i_v b), \quad a \in \Lambda^k(V), b \in \Lambda^l(V).$$

**Proposition 2.7.5.**

Let

$$a = \frac{1}{k!} a_{i_1 \dots i_k} f^{(i_1)} \wedge \dots \wedge f^{(i_k)} \in \Lambda^k(V).$$

Then

$$i_v a = \frac{1}{(k-1)!} v^j a_{j i_2 \dots i_k} f^{(i_2)} \wedge \dots \wedge f^{(i_k)}.$$



## 12. Differential forms.

**Definition 3.1.1.** Let  $U$  be an open subset of  $\mathbb{R}^n$ . A **differential  $k$ -form** on  $U$ , or  **$k$ -form** for short, is a smooth map

$$\alpha : U \rightarrow \Lambda^k(\mathbb{R}^n); \quad x \mapsto \alpha(x) = \frac{1}{k!} \alpha_J(x) F^{(J)}.$$

Here, “smooth” means that the coefficient functions are smooth, i.e.  $\alpha_J(x) \in \mathbb{C}^\infty(U)$ . Zero-forms are just smooth functions on  $U$ .

**Definition 3.1.2.** Given  $\alpha, \beta \in \Omega^k(U)$ , then  $\alpha(x)$  and  $\beta(x)$  are both algebraic  $k$ -forms, and it makes sense to add them. Thus, we define  $\alpha + \beta \in \Omega^k(U)$  by

$$(\alpha + \beta)(x) := \alpha(x) + \beta(x).$$

**Definition 3.1.3.** If  $\alpha$  is a  $k$ -form on  $U$  and  $\beta$  is an  $l$ -form, then we define the  $(k+l)$ -form  $\alpha \wedge \beta$  by

$$(\alpha \wedge \beta)(x) := \alpha(x) \wedge \beta(x).$$

**Definition 3.1.4.** Let  $U \subset \mathbb{R}^n$  be open. Given a differential  $k$ -form  $\omega \in \Omega^k$  and a vector field  $\mathbb{X} \in \mathcal{X}(U)$ , the **contraction** of  $\omega$  with  $\mathbb{X}$ , denoted  $i_{\mathbb{X}}\omega$ , is the differential  $(k-1)$ -form defined by

$$(i_{\mathbb{X}}\omega)(x) := i_{\mathbb{X}(x)}\omega(x).$$

## 13. Exterior derivative.

**Definition 3.2.1.** Let  $U \subset \mathbb{R}^n$  be open. The **exterior derivative**, denoted  $d : \Omega^k(U) \rightarrow \Omega^{k+1}(U)$ , is a map from differential  $k$ -forms to differential  $(k+1)$ -forms defined as follows:

- $k = 0$ . For  $g \in \Omega^0(U) = C^\infty(U)$ ,

$$dg := \frac{\partial g}{\partial x^i} f^{(i)}.$$

- $k > 0$ . For  $\omega \in \Omega^k(U)$ , we may write that

$$\omega = \frac{1}{k!} \omega_{i_1 \dots i_k} f^{(i_1)} \wedge \dots \wedge f^{(i_k)},$$

where  $\omega_{i_1 \dots i_k}(x) \in C^\infty(U)$  (cf Proposition 2.4.10). Then

$$d\omega(x) := \frac{1}{k!} d\omega_{i_1 \dots i_k} \wedge f^{(i_1)} \wedge \dots \wedge f^{(i_k)}.$$

Equivalently,

$$d\omega(x) := \frac{1}{k!} \frac{\partial \omega_{i_1 \dots i_k}}{\partial x^j} f^{(j)} \wedge f^{(i_1)} \wedge \dots \wedge f^{(i_k)}.$$

**Notation.** It is conventional to write  $dx^i$  instead of  $f^{(i)}$ .

**Proposition 3.2.2.** Let  $\alpha, \beta \in \Omega^k(U)$ . Then

$$d(\alpha + \beta) = d\alpha + d\beta.$$

**Proposition 3.2.3.** Let  $\alpha \in \Omega^k(U)$  and  $\beta \in \Omega^l(U)$ . Then

$$d(\alpha \wedge \beta) = (d\alpha) \wedge \beta + (-1)^k \alpha \wedge d\beta.$$

**Proposition 3.2.4.** For all  $\omega \in \Omega^k(U)$ ,

$$d^2\omega = 0.$$

#### 14. Pullback.

**Definition 3.3.1.** Let  $U \subset \mathbb{R}^m$  and  $V \subset \mathbb{R}^n$  be open, and let  $F : U \rightarrow V$  be a smooth map. The **pullback**, denoted  $F^*$ , is a map  $F^* : \Omega^k(V) \rightarrow \Omega^k(U)$  – that is, the pullback maps differential forms on  $V$  back to differential forms on  $U$ . Given  $\beta \in \Omega^k(V)$ ,  $F^*\beta$  is defined as follows. We note that as  $F^*\beta$  is a differential  $k$ -form on  $U$ ,  $F^*\beta(x)$  is an algebraic  $k$ -form on  $\mathbb{R}^m$ , which may be defined by specifying its value when applied to  $k$  arbitrary vectors in  $\mathbb{R}^m$ . Denoting these vectors by  $u_{(1)}, \dots, u_{(k)}$ , the definition is given by

$$(F^*\beta)(x; \mathbf{u}_{(1)}, \dots, \mathbf{u}_{(k)}) = \beta(F(x); F'(x)\mathbf{u}_{(1)}, \dots, F'(x)\mathbf{u}_{(k)}).$$

For 0-forms, i.e. functions,  $F^*f = f \circ F$ , in accord with Definition 1.9.1

**Proposition 3.3.4.**

$$F^*(\beta \wedge \gamma) = F^*\beta \wedge F^*\gamma.$$

**Proposition 3.3.7.** Let  $\beta \in \Omega^k(V)$ . Then

$$F^*d\beta = dF^*\beta.$$

**Proposition 3.3.9.** Let  $U \subset \mathbb{R}^m$ ,  $V \subset \mathbb{R}^n$  and  $W \subset \mathbb{R}^p$  be open. Let  $F : U \rightarrow V$  and  $G : V \rightarrow W$  be smooth maps. Then

$$(G \circ F)^* = F^*G^*.$$

#### 15. Lie derivative and Poincaré Lemma

**Definition 3.4.1.** Let  $U \subset \mathbb{R}^n$  be open, and let  $\mathbb{X} \in \mathcal{X}(U)$  be a smooth vector field on  $U$  with flow  $\Phi_t$ . Let  $\omega \in \Omega^k(U)$  be a differential  $k$ -form. The **Lie derivative** of  $\omega$  with respect to  $\mathbb{X}$ , denoted  $L_{\mathbb{X}}\omega$ , is given by

$$L_{\mathbb{X}}\omega = \left. \frac{\partial}{\partial t} \right|_{t=0} \Phi_t^*\omega.$$

**Proposition 3.4.2.**

- i)  $L_{\mathbb{X}}(\alpha + \beta) = L_{\mathbb{X}}\alpha + L_{\mathbb{X}}\beta$
- ii)  $L_{\mathbb{X}}(\alpha \wedge \beta) = (L_{\mathbb{X}}\alpha) \wedge \beta + \alpha \wedge L_{\mathbb{X}}\beta$
- iii)  $L_{\mathbb{X}}d = dL_{\mathbb{X}}$ .

**Proposition 3.4.3.** For  $\omega \in \Omega^k(U)$ ,

$$L_{\mathbb{X}}\omega = i_{\mathbb{X}}d\omega + di_{\mathbb{X}}\omega.$$

**Definition 3.5.1.** A differential  $k$ -form  $\omega$  is **closed** if  $d\omega = 0$ .  $\omega$  is **exact** if  $\omega = d\alpha$  for some  $(k-1)$ -form  $\alpha$ .

**Definition 3.5.2.** Let  $U, V \subset \mathbb{R}^n$  be open, and let  $I$  be an open interval in  $\mathbb{R}$ . A **one-parameter family of diffeomorphisms** is a smooth family of maps

$$\hat{\Phi} : I \times U \rightarrow V; \quad (t, x) \mapsto \hat{\Phi}_t(x)$$

such that  $\hat{\Phi}_t$  is a diffeomorphism onto its image. That is, letting  $U_t = \hat{\Phi}_t(U) \subset V$ , then  $\hat{\Phi}_t : U \rightarrow U_t$  is a diffeomorphism.

**Proposition 3.5.4.**

$$\left. \frac{\partial}{\partial t} \hat{\Phi}_t^* \right|_{t=t_0} \omega = \hat{\Phi}_{t_0}^* L_{\hat{\mathbb{X}}_{t_0}} \omega.$$

**Theorem 3.5.5. (Poincaré Lemma.)** Let  $\hat{\Phi}_t : U \rightarrow U$  be a one-parameter family of diffeomorphisms defined for  $0 < t \leq 1$ . Let  $\beta \in \Omega^k(U)$  be a closed  $k$ -form. Suppose that

$$\hat{\Phi}_1^* \beta = \beta, \quad \lim_{t \rightarrow 0} \hat{\Phi}_t^* \beta = 0.$$

Then

$$\beta = d\alpha,$$

where

$$\alpha = \int_0^1 \hat{\Phi}_t^* (i_{\hat{\mathbb{X}}_t} \beta) dt,$$

and  $\hat{\mathbb{X}}_t$  is defined as above by

$$\frac{\partial}{\partial t} \hat{\Phi}_t(x) = \hat{\mathbb{X}}_t(\hat{\Phi}_t(x)).$$

## 16. Singular $k$ -cubes and integration of differential forms.

**Definition 4.1.1.** Let  $U \subset \mathbb{R}^n$  be open. A singular  $k$ -cube on  $U$  is a smooth map

$$c : I^k \rightarrow U.$$

**Definition 4.1.2.** The integral of a  $k$ -form  $\omega$  over a singular  $k$ -cube  $c$ , denoted  $\int_c \omega$ , is defined by

$$\int_c \omega := \int_{I^k} c^* \omega := \int_{I^k} f(t) dt^1 \cdots dt^k.$$

**Definition 4.1.4.** Let  $U, V \subset \mathbb{R}^n$  be open. Let  $G : U \rightarrow V$  be a diffeomorphism. Then  $\det G'(x) \neq 0$  for all  $x \in U$ . We say that  $G$  is orientation-preserving if  $\det G' > 0$  on  $U$ .

**Proposition 4.1.6.** Let  $B \subset \mathbb{R}^k$  be an open set which contains the  $k$ -cube  $I^k$ . Let  $G : B \rightarrow B$  be an orientation-preserving diffeomorphism, and suppose that  $G(I^k) = I^k$ . Let  $c : I^k \rightarrow U$  be a singular  $k$ -cube, and  $\omega \in \Omega^k(U)$ . Then

$$\int_c \omega = \int_{c \circ G} \omega.$$

## 17. Boundaries.

**Definition 4.2.1.** A singular  $k$ -chain on  $U$ , denoted  $\mathcal{C}$ , is a formal sum of a finite number of singular  $k$ -cubes  $c_r : I^k \rightarrow U$  with integer coefficients, i.e.

$$\mathcal{C} = a_1 c_1 + \cdots + a_s c_s, \quad a_r \in \mathbb{Z}.$$

**Definition 4.2.2.** Let  $c : I^k \rightarrow U$  be a singular  $k$ -cube on  $U$ . Take  $j$  such that  $1 \leq j \leq k$  and  $\alpha = 0$  or  $1$ . The  $(j, \alpha)$ -th face of  $c$ , denoted  $c_{(j, \alpha)}$ , is the singular  $(k-1)$ -cube given by

$$c_{(j, \alpha)} : I^{k-1} \rightarrow U,$$

where

$$c_{(j,\alpha)}(t^1, \dots, t^{k-1}) = c(t^1, \dots, t^{j-1}, \alpha, t^j, \dots, t^{k-1}).$$

**Definition 4.2.3.** Let  $c : I^k \rightarrow U$  be a singular  $k$ -cube. The boundary of a singular  $k$ -cube  $c$ , denoted  $\partial c$ , is the singular  $(k-1)$ -chain given by

$$\partial c = \sum_{j=1}^k \sum_{\alpha=0,1} (-1)^{j+\alpha} c_{(j,\alpha)}.$$

Let  $\omega \in \Omega^{k-1}(U)$  be a  $(k-1)$ -form, and  $c : I^k \rightarrow U$  a singular  $k$ -cube. Then

$$\int_{\partial c} \omega = \sum_{j=1}^k \sum_{\alpha=0,1} (-1)^{j+\alpha} \int_{c_{(j,\alpha)}} \omega.$$