

Crop Impacts from Stratospheric Aerosol Climate Intervention: A Multi-Scenario Overview





Brendan Clark

Outline





- Introduce model
- Scenarios overview
- Global crop impacts
- Regional impacts
- Individual climate variable impacts
- Future directions


Model

- Results here are from fully coupled CESM2-CLM5crop simulations
- $0.95^{\circ} \times 1.25^{\circ}$ latitude-longitude resolution
- CLM5crop simulates diffuse radiation fertilization and includes a surface ozone parameterization that is the same for all crops
- Yield and production are calculated using year 2000 constant cropping area




(Tilmes et al., 2020)







Maize SSP2-4.5-1.5°C minus SSP2-4.5 (2060-2070 avg.)




Maize SSP5-8.5-1.5°C minus SSP5-8.5 (2060-2070 avg.)




Maize SSP5-3.4-OS-1.5°C minus SSP5-3.4-OS (2060-2070 avg.)





Maize G6Solar minus SSP5-8.5 (2060-2070 avg.)




Maize SSP5-3.4-OS-2.0°C minus SSP5-3.4-OS (2060-2070 avg.)




Maize G6Sulfur minus SSP5-8.5 (2060-2070 avg.)




Change in **Maize** Production for Top 10 Producers Under SAI (2060-2070 avg.)




Change in **Rice** Production for Top 10 Producers Under SAI (2060-2070 avg.)




Change in **Soybean** Production for Top 10 Producers Under SAI (2060-2070 avg.)



Change in **Spring Wheat** Production for Top 10 Producers Under SAI (2060-2070 avg.)



- Offline (land only) CLM5crop simulations forced with atmospheric coupler data from SSP2-4.5-1.5°C and SSP2-4.5 fully coupled CESM2 simulations
- More simulations currently being run with UKESM1
- Hope to extend simulations to the end of the century for a larger signal
- Can use atmospheric coupler data to force other crop models



Conclusion

- Lowering temperature while maintaining elevated CO₂ benefits maize, rice, soybean, and spring wheat on a global average under all stratospheric aerosol injection scenarios
- All scenarios show a yield reduction for multiple top producing countries under stratospheric aerosol injection
 - Who gets to decide to increase production in some countries while decreasing production in others?
- Need to incorporate UV radiation impacts and surface ozone impacts for each crop
- Atmospheric coupler data output from SSP2-4.5-1.5°C and SSP2-4.5 simulations (CESM2 and UKESM1) can be used to force a multi-model assessment of crop impacts from stratospheric aerosol injection
- Currently planning to use CLM5crop, pDSSAT, and LPJmL, but other models are encouraged to participate