



Designing Microbiome Trials

Unique Challenges & Considerations

Brendan J. Kelly, MD, MS
Infectious Diseases, Microbiology & Epidemiology
University of Pennsylvania

GMFH World Summit
12 March 2022

Disclosures

- No conflicts of interest.
- Research supported by:
 - NIAID K23 AI121485
 - CDC BAA 200-2016-91964
 - CDC BAA 200-2016-91937
 - CDC BAA 200-2018-02919
 - CDC BAA 200-2021-10986

How Randomized Trials Go Wrong

How Randomized
Trials Go Wrong

RCTs of FMT

How Randomized
Trials Go Wrong

RCTs of FMT

Strategies for
the LBP Era

How Randomized
Trials Go Wrong

RCTs of FMT

Strategies for
the LBP Era

"Understanding and Misunderstanding RCTs"

- Randomized clinical trials = gold standard for clinical research:
 - balance known & unknown confounders
 - powerful tool to measure causal effects
- RCTs can still mislead:
 - selection bias (subjects \neq population)
 - "trial interventions are interactions" resulting in heterogenous effects (subject features modify effect of trial intervention)
 - "subjects aren't even fully representative of themselves" (time-varying features also interact with trial intervention)

Time-Varying Gut Microbiota

Table 1. Effect sizes observed from various exposures/interventions in studies of various microbiome sampling sites are shown as measured by omega-squared (ω^2) statistics, together with the P -values from PERMANOVA test

Site	Comparison groups		ω^2/P -value				Reference
	Control	Exposure	Weighted UniFrac	Unweighted UniFrac	Weighted Jaccard	Unweighted Jaccard	
Nares	Non-smoker (33)	Smoker (29)	0.042/0.001	0.009/0.001	0.023/0.001	0.007/0.001	Charlson <i>et al.</i> (2010)
Oral	Non-smoker (33)	Smoker (29)	0.032/0.001	0.008/0.001	0.024/0.001	0.007/0.001	Charlson <i>et al.</i> (2010)
Gut	Before feeding (10)	After feeding (10)	0.056/0.138	0.013/0.986	0/0.989	0.014/0.985	Wu <i>et al.</i> (2011)
Oral	No azithromycin (42)	Azithromycin (6)	0.063/0.01	0.039/0.001	0.099/0.004	0.032/0.001	Charlson <i>et al.</i> (2012)
Lung	No azithromycin (34)	Azithromycin (6)	0.065/0.005	0.038/0.001	0.019/0.089	0.033/0.001	Charlson <i>et al.</i> (2012)
Skin	Left retroauricular (186)	Right retroauricular (187)	0.000/0.828	0.0001/0.327	0.000/0.986	0.000/1.000	HMP Consortium (2012b)
Human	Anterior nares (161)	Stool (187)	0.567/0.001	0.201/0.001	0.230/0.001	0.117/0.001	HMP Consortium (2012b)

The range of observed effect sizes differs according to the metric of pairwise distance chosen for analysis. HMP data are shown to demonstrate a large effect (the degree of difference between two different human microbiome sampling sites) and a negligible effect (the degree of difference between skin sampling in the left versus right retroauricular crease)

Time-Varying Gut Microbiota

- Dietary changes and antibiotics exert large effects...
 - ... larger effects when followed by colonization from healthcare environment
- Gut microbiota at time of intervention \neq microbiota at time of enrollment:
 - misclassification of inclusion criteria kills randomized trials
 - "the medicine doesn't work if the patient isn't sick"
 - microbiome trials run a **high risk of eligibility misclassification**

How Microbiota Trials Go Right

- RCTs can overcome effect modification by changing gut microbiota if:
 - very precise disease phenotype (less likely time-varying gut microbiota)
 - precision medicine: tailor treatment to near-real-time measures of gut microbiota
 - interventions with huge effects (antibiotic conditioning prior to FMT)

How Randomized
Trials Go Wrong

RCTs of FMT

Strategies for
the LBP Era

Fecal Microbiota Transplant (FMT) Trials


- FMT in Infectious Diseases @ the University of Pennsylvania:
 - FMT dose finding for recurrent *C. difficile* infection (NCT03973697)
 - serial FMT for severe *C. difficile* infection (NCT03970200)
 - FMT for MDRO colonization (CDC sponsored)
- Successful applications of FMT:
 - recurrent & severe *C. difficile*, inflammatory bowel diseases, potentiation of anti-PD1 immunotherapy, MDRO colonization, food allergy mitigation
 - heterogeneity of gut microbiota community composition?

Microbial Heterogeneity in *C. difficile*

- **Aim:** microbiome features that discriminate *C. difficile* colonization / infection
- **Population:** 384 consecutive positive *C. difficile* tests (in- & outpatient)


<i>C. difficile</i> Category	Subject Count	Proportion
GDH+ Toxin EIA- NAAT-	94	24.5%
GDH+ Toxin EIA- NAAT+	213	55.5%
GDH+ Toxin EIA+	77	20.1%

- **Sampling:** stool 16S rRNA gene sequencing & 16S rRNA gene qPCR
- **Comparison:** toxin EIA+ (infection) versus NAAT+ only (colonization)
- **Outcome:** EIA+, with fecal lactoferrin as sensitivity analysis



Univariable toxin EIA model: *Clostridioides* OR 1.07 (95%CrI 1.02 - 1.12)

Multivariable toxin EIA model: *Clostridioides* OR 1.05 (95%CrI 1.01 - 1.16) & *Lachnospiraceae* OR 0.32 (95%CrI 0.14 - 0.96)



Phenotype Definitions & FMT Outcomes

- The success of FMT depends on homogeneity of gut microbiota phenotype:
 - moderate heterogeneity of *C. difficile* infection
 - large heterogeneity across IBD
- Pre-FMT antibiotic conditioning as a method to enforce homogeneity:
 - which antibiotics are used?
 - effects of other medications active on gut microbiota (e.g., PPIs)?
 - FMT as part of a bundled intervention?

How Randomized
Trials Go Wrong

RCTs of FMT

**Strategies for
the LBP Era**

Into the LBP Era

- Live biotherapeutic products (LBPs) replacing FMT:
 - Ferring, Seres, Vedanta, Finch...
 - ensure greater homogeneity of intervention
 - larger effects?
- Are bundled interventions still necessary?
 - role for pre-LBP antibiotic conditioning?
 - restrictions on medications that reshape gut microbial communities?
 - role for precision medicine & pre-treatment microbiota profiles?

How Randomized
Trials Go Wrong

RCTs of FMT

Strategies for
the LBP Era

Topics for Discussion

- Every RCT intervention is an interaction:
 - gut microbiota community composition varies over time
 - account for changes in diet & medications, environment
- RCTs can overcome effect modification if:
 - very precise disease phenotype (less likely time-varying gut microbiota)
 - precision medicine: tailor treatment to near-real-time measures of gut microbiota
 - interventions with huge effects (antibiotic conditioning prior to FMT)
- How to translate lessons from FMT trials to the LBP era?



Questions?

slides ↓

bjklab.org