

Table 1: Treatment Effects on Fertilizer Use

	All crops		Maize only	
	(1)	(2)	(3)	(4)
Treatment one	150.57 (147.71)	52.77 (145.49)	3.37 (6.87)	2.99 (6.56)
Treatment two	249.05 (217.73)	147.35 (156.70)	28.60*** (6.01)	35.07*** (5.61)
Constant	78.46*** (5.84)	132.30* (72.77)	83.36*** (5.59)	22.29** (10.66)
Control variables	No	Yes	No	Yes
Number of Observations	1953	1869	1626	1626

Notes: This table reports the effects of two interventions on total fertilizer use (in kilograms): (i) providing site-specific fertilizer recommendations (Treatment 1), and (ii) providing site-specific fertilizer recommendations together with a voucher to purchase the recommended amount (Treatment 2). Columns (1)–(2) present the effects on fertilizer use across all crops, while Columns (3)–(4) focus specifically on maize. Columns (1) and (3) report estimates without controlling for household- and plot-level characteristics, whereas Columns (2) and (4) include these controls: household size, household head age, household head education, distance to agro-dealer, plot size, visited agro-dealer, access to extension services, plot slope, soil structure, and seed type used. Asterisks indicate statistical significance levels: *** at 1%, ** at 5%, and * at 10%. Standard errors are clustered at the cluster level.

Table 2: Effect of the Program on Fertilizer Use (AIP)

	All crops		Maize only	
	(1)	(2)	(3)	(4)
Treatment one	-1.44 (1.60)	-1.28 (1.76)	-1.14 (2.01)	-0.88 (2.04)
Treatment two	-7.49*** (1.38)	-7.97*** (1.45)	-9.00*** (1.64)	-8.44*** (1.66)
Constant	10.23*** (1.17)	11.69*** (2.95)	12.00*** (1.44)	9.39*** (3.09)
Control variables	No	Yes	No	Yes
Number of Observations	1953	1869	1626	1626

Notes: This table reports the effects of two interventions on fertilizer use (AIP, in kilograms): (i) providing site-specific fertilizer recommendations (Treatment 1), and (ii) providing site-specific fertilizer recommendations together with a voucher to purchase the recommended amount (Treatment 2). Columns (1) and (3) report estimates without controlling for household- and plot-level characteristics; columns (2) and (4) include these controls: household size, household head age, household head education, distance to agro-dealer, plot size, visited agro-dealer, access to extension services, plot slope, soil structure, and seed type used. Asterisks indicate statistical significance: *** $p < 0.01$, ** $p < 0.05$, * $p < 0.10$. Standard errors are clustered at the cluster level.

Table 3: Effect of the Program on Fertilizer Use by Nutrient

		Dependent variable: Total fertilizer use in nutrient amount (kg)					
Total Nitrogen (kg)		Total Phosphorus (kg)		Total Potassium (kg)		Total Nutrients (N+P+K, kg)	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Treatment one	0.54 (2.21)	0.43 (2.11)	0.36 (0.34)	0.20 (0.18)	0.18 (0.17)	1.12 (2.66)	0.97 (2.55)
Treatment two	9.51*** (1.90)	11.58*** (1.80)	1.52*** (0.30)	1.85*** (0.28)	0.76*** (0.16)	11.79*** (2.33)	14.37*** (2.19)
Control variables	No	Yes	No	Yes	No	Yes	Yes
Observations	1626	1626	1626	1626	1626	1626	1626

Notes: This table reports the effects of two interventions on total fertilizer use (in kilograms): (i) providing site-specific fertilizer recommendations (Treatment 1), and (ii) providing site-specific fertilizer recommendations together with a voucher to purchase the recommended amount (Treatment 2). Columns (1)–(2) present effects on nitrogen use, (3)–(4) on phosphorus, (5)–(6) on potassium, and (7)–(8) on total nutrients (N+P+K). Odd-numbered columns report estimates without controlling for household- and plot-level characteristics; even-numbered columns include these controls: household size, household head age, household head education, distance to agro-dealer, plot size, visited agro-dealer, access to extension services, plot slope, soil structure, and seed type used. Asterisks denote significance at the 1% (***), 5% (**), and 10% (*) levels. Standard errors are clustered at the cluster level.

Table 4: Impact of the Program on Maize Yield

	(1) Log of maize yield (kg/acre)	(2) Log of maize yield (kg/acre)
Treatment one	-0.13 (0.11)	-0.14 (0.10)
Treatment two	0.36*** (0.10)	0.31*** (0.10)
Constant	6.01*** (0.06)	6.09*** (0.16)
Control variables	No	Yes
Number of Observations	1610	1610

Notes: This table reports the impact of two interventions on maize yield (kg/acre): (i) providing site-specific fertilizer recommendations (T1), and (ii) providing site-specific fertilizer recommendations together with a voucher to purchase the recommended amount (T2). Column (1) reports estimates without controlling for household- and plot-level characteristics; Column (2) includes these controls: household size, household head age, household head education, distance to agro-dealer, plot size, visited agro-dealer, access to extension services, plot slope, soil structure, and seed type used. Asterisks indicate statistical significance: *** $p < 0.01$, ** $p < 0.05$, * $p < 0.10$. Standard errors are clustered at the cluster level.