

Table 1: Impact of of IVR and SMS treatments on knowledge (conditional on access to mobile phone)

	Mean	+IVR	p-value	N	+SMS	p-value	N
Knows optimal spacing (yes=1)	0.174 (0.380)	-0.015 (0.020)	0.392	3,002	0.002 (0.022)	0.941	2,736
Knows inputs best combined (yes=1)	0.905 (0.293)	-0.013 (0.011)	0.173	3,002	0.005 (0.012)	0.699	2,736
Knows optimal time for weeding (yes=1)	0.945 (0.228)	0.003 (0.011)	0.742	3,002	0.007 (0.012)	0.618	2,736
Knows how to fight armyworm (yes=1)	0.338 (0.474)	-0.009 (0.021)	0.690	3,002	0.008 (0.023)	0.720	2,736
Knowledge index	-0.109 (0.610)	-0.023 (0.027)	0.317	3,002	0.018 (0.030)	0.613	2,736

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3 and number of observations in column 4; column 5 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 6; sample size is reported in column 7. Reported p-values are based on randomization inference (10,000 permutations). All specifications control for the other orthogonal factors in the factorial design.

Table 2: 2SLS estimates of impact of IVR and SMS treatments on household level knowledge

	Mean	+IVR	p-value	+SMS	p-value	N
Knows optimal spacing (yes=1)	0.160 (0.367)	-0.360 (0.351)	0.306	0.014 (0.026)	0.596	3,619
Knows inputs best combined (yes=1)	0.908 (0.290)	-0.331 (0.210)	0.115	0.014 (0.015)	0.379	3,619
Knows optimal time for weeding (yes=1)	0.954 (0.210)	0.178 (0.191)	0.350	0.000 (0.015)	0.979	3,619
Knows how to fight armyworm (yes=1)	0.336 (0.473)	-0.286 (0.363)	0.431	0.023 (0.028)	0.401	3,619
Knowledge index	-0.077 (0.562)	-0.396 (0.473)	0.376	0.026 (0.036)	0.458	3,619

Note: In the first column, means (and standard deviations) in the comparison group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3; column 4 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 5; sample size is reported in column 6. All specifications control for the other orthogonal factors in the factorial design.

Table 3: Impact of IVR and SMS treatments on agronomic practices (conditional on phone access)

	Mean	+IVR	p-value	N	+SMS	p-value	N
Planted immediately after start of rains (yes=1)	0.365 (0.483)	0.013 (0.022)	0.559	2,924	-0.020 (0.024)	0.433	2,673
Used spacing of 75cm x 30cm with a reduced seed rate (yes=1)	0.030 (0.171)	-0.011 (0.013)	0.360	2,962	0.014 (0.014)	0.331	2,707
Removed striga early on (yes=1)	0.690 (0.464)	-0.003 (0.020)	0.867	2,962	0.021 (0.022)	0.376	2,707
First weeding after 18-20 days (yes=1)	0.435 (0.497)	0.019 (0.023)	0.331	2,962	0.002 (0.025)	0.950	2,707
Recommended practices index	-0.088 (0.482)	0.008 (0.024)	0.685	2,924	0.014 (0.026)	0.584	2,673

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3 and number of observations in column 4; column 5 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 6; sample size is reported in column 7. Reported p-values are based on randomization inference (10,000 permutations). All specifications control for the other orthogonal factors in the factorial design.

Table 4: 2SLS estimates of impact of IVR and SMS treatments on agronomic practices

	Mean	+IVR	p-value	+SMS	p-value	N
Planted immediately after start of rains (yes=1)	0.370 (0.484)	0.163 (0.391)	0.677	-0.002 (0.029)	0.940	3,500
Used spacing of 75cm x 30cm with a reduced seed rate (yes=1)	0.026 (0.158)	-0.089 (0.210)	0.670	0.020 (0.017)	0.231	3,560
Removed striga early on (yes=1)	0.685 (0.465)	-0.170 (0.339)	0.617	0.033 (0.027)	0.215	3,560
First weeding after 18-20 days (yes=1)	0.426 (0.495)	0.228 (0.383)	0.552	-0.012 (0.030)	0.693	3,560
Recommended practices index	-0.086 (0.478)	0.102 (0.411)	0.804	0.028 (0.031)	0.364	3,500

Note: In the first column, means (and standard deviations) in the comparison group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3; column 4 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 5; sample size is reported in column 6. All specifications control for the other orthogonal factors in the factorial design.

Table 5: Impact of IVR and SMS treatments on fertilizer and improved seed use (conditional on access to mobile phone)

	Mean	+IVR	p-value	N	+SMS	p-value	N
				<i>fertilizer use</i>			
Used DAP/NPK on at least one plot? (yes=1)	0.285 (0.453)	0.029 (0.020)	0.095	2,962	-0.022 (0.022)	0.408	2,707
Used urea on at least one plot? (yes=1)	0.050 (0.218)	0.013 (0.014)	0.288	2,962	-0.031 (0.015)	0.050	2,707
Used organic fertilizer on at least one plot? (yes=1)	0.150 (0.358)	-0.044** (0.019)	0.011	2,962	0.037 (0.021)	0.083	2,707
Fertilizer index	-0.074 (0.531)	-0.002 (0.027)	0.908	2,962	-0.015 (0.029)	0.675	2,707
				<i>seed use</i>			
Used hybrid maize seed on at least one plot? (yes=1)	0.330 (0.471)	0.023 (0.021)	0.234	2,962	-0.053 (0.023)	0.028*	2,707
Used Open Pollinated Varieties on at least one plot? (yes=1)	0.275 (0.448)	0.011 (0.021)	0.543	2,962	0.026 (0.023)	0.386	2,707
Seed index	0.010 (0.701)	0.038 (0.030)	0.161	2,962	-0.026 (0.033)	0.475	2,707

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3 and number of observations in column 4; column 5 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 6; sample size is reported in column 7. Reported p-values are based on randomization inference (10,000 permutations); ** and * denote that the difference is significant at the 5 and 10 percent level, respectively, after correcting for multiple hypothesis testing using a family-wise sharp null (10,000 permutations). All specifications control for the other orthogonal factors in the factorial design.

Table 6: 2SLS estimates of impact of IVR and SMS treatments on fertilizer and improved seed use

	Mean	+IVR	p-value	+SMS	p-value	N
				<i>fertilizer use</i>		
Used DAP/NPK on at least one plot? (yes=1)	0.264 (0.442)	0.393 (0.325)	0.228 (0.025)	-0.011 (0.025)	0.661	3,560
Used urea on at least one plot? (yes=1)	0.051 (0.221)	0.225 (0.226)	0.320 (0.018)	-0.030 (0.044)	0.092	3,560
Used organic fertilizer on at least one plot? (yes=1)	0.157 (0.365)	-0.704 (0.337)	0.037 (0.025)	0.044 (0.025)	0.077	3,560
Fertilizer index	-0.057 (0.547)	-0.078 (0.443)	0.846	-0.003 (0.035)	0.994	3,560
				<i>seed use</i>		
Used hybrid maize seed on at least one plot? (yes=1)	0.289 (0.454)	0.732 (0.367)	0.046 (0.027)	-0.063 (0.027)	0.022	3,560
Used Open Polinated Varieties on at least one plot? (yes=1)	0.302 (0.460)	0.164 (0.345)	0.635 (0.027)	0.033 (0.027)	0.220	3,560
Seed index	0.032 (0.700)	0.242 (0.447)	0.605	0.004 (0.036)	0.876	3,560

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3 and number of observations in column 4; column 5 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 6; sample size is reported in column 7. Reported p-values are based on randomization inference (10,000 permutations); ***, ** and * denote that the difference is significant at the 1, 5 and 10 percent level, respectively, after correcting for multiple hypothesis testing using a family-wise sharp null (10,000 permutations). All specifications control for the other orthogonal factors in the factorial design.

Table 7: Impact of IVR and SMS treatments on household level production (conditional on access to mobile phone)

	Mean	+IVR	p-value	N	+SMS	p-value	N
Maize production (log(kg))	5.921 (0.706)	0.057 (0.036)	0.146	2,742	-0.001 (0.039)	0.894	2,511
Maize area (log(acre))	0.054 (0.554)	-0.018 (0.029)	0.492	2,782	0.004 (0.031)	0.997	2,545
Maize yield (log(kg/acre))	5.879 (0.654)	0.042 (0.031)	0.203	2,756	0.011 (0.034)	0.737	2,518
Yield better than normal (yes=1)	0.405 (0.492)	-0.005 (0.023)	0.812	2,962	0.036 (0.024)	0.166	2,707
Labour(log(mandays))	4.157 (0.571)	-0.009 (0.027)	0.729	2,802	0.021 (0.028)	0.515	2,549
Labour productivity (log(kg/mandays))	1.673 (0.708)	0.060 (0.036)	0.082	2,780	-0.012 (0.039)	0.672	2,543
Production index	-0.052 (0.378)	0.016 (0.017)	0.306	2,756	0.009 (0.018)	0.572	2,518

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3 and number of observations in column 4; column 5 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 6; sample size is reported in column 7. Reported p-values are based on randomization inference (10,000 permutations). All specifications control for the other orthogonal factors in the factorial design.

Table 8: 2SLS estimates of impact of IVR and SMS treatments on production

	Mean	+IVR	p-value	+SMS	p-value	N
Maize production (log(kg))	5.814 (0.765)	0.912 (0.670)	0.173	0.054 (0.050)	0.276	3,344
Maize area (log(acre))	0.018 (0.580)	-0.240 (0.476)	0.614	0.031 (0.038)	0.415	3,341
Maize yield (log(kg/acre))	5.850 (0.658)	0.690 (0.568)	0.224	0.005 (0.041)	0.898	3,302
Yield better than normal (yes=1)	0.387 (0.488)	0.008 (0.374)	0.983	0.048 (0.030)	0.103	3,560
Labour(log(mandays))	4.132 (0.577)	-0.056 (0.421)	0.895	0.037 (0.036)	0.296	3,370
Labour productivity (log(kg/mandays))	1.650 (0.720)	1.185 (0.667)	0.076	0.000 (0.048)	0.996	3,341
Production index	-0.053 (0.365)	0.159 (0.297)	0.625	0.007 (0.022)	0.702	3,302

Note: In the first column, means (and standard deviations) in the control group are presented for each variable. Column 2 reports differences between video only and video+ivr (and standard error) with its corresponding p-value in column 3; column 4 reports differences between video+ivr and video+ivr+SMS (and standard error) with its corresponding p-value in column 5; sample size is reported in column 6. All specifications control for the other orthogonal factors in the factorial design.

ICT-mediated agricultural knowledge transfer: A pre-analysis plan

January 9, 2019

Abstract

In agricultural information dissemination campaigns through agricultural extension services, seemingly small attributes, such as the way the information message is delivered, who delivers the information messages or who is targeted by the information message, can result in significant differences in outcomes, such as knowledge transferred, practices adopted, agricultural yields, and ultimately well-being within the household. In the context of Information & Communication Technology (ICT) mediated knowledge transfer, this study investigates the role of the gender composition of the person(s) who provide(s) the information and the gender composition of the person(s) who receive(s) the information in making the information transfer more effective. Furthermore, we quantify the additional impact of augmenting video as a way to deliver extension information with a more demand-driven approach such as Interactive Voice Response (IVR) technology. Effectiveness is assessed in terms of knowledge gain, increased adoption, yield increase, and poverty reduction, and outcomes will be disaggregated by gender to study changes in intra-household equity and women empowerment. The research questions will be answered using field experiments, where farmers are randomly assigned to particular combinations of interventions. This pre-analysis plan outlines hypotheses which will be tested, tools that will be used in the field, power calculations and sample size projections on which sampling is based, outcome variables that will be used to assess impact, and specifications that will be estimated. As such, it will provide a useful reference in evaluating the final results of the study (Humphreys, Sanchez de la Sierra, and van der Windt, 2013).

Motivation

In a world characterized by incomplete and asymmetric information, targeted efforts to fill knowledge gaps can make a big difference. However, not every information campaign is equally effective, and often, seemingly small attributes, such as the way a message is framed, how it is delivered, who is targeted by the message and who delivers the message, can result in significant differences in impact (Jack, 2013). Understanding and quantifying the importance of each of these attributes through rigorous evaluation research should therefore be an essential part of any knowledge exchange model.

This research will investigate some attributes that are thought to be key in effective agricultural extension information delivery. In particular, we will zoom in on the role of (i) the gender composition of the receiver(s) of the agricultural extension information messages; (ii) the gender composition of the messenger(s) of the information message; and (iii) the way it is delivered. The questions will be answered through field experiments, where farmers are randomly assigned to a group that receives particular interventions, and their outcomes will be compared to outcomes of a group of farmers that did not receive these particular interventions (?). As we want to test more than one research hypothesis and are particularly interested in comparing the relative effectiveness of various attributes of an ICT-mediated extension approach (instead of simply finding out what works and compare treatment to control groups), we opt for a factorial design. In such a design, farmers get a particular combination of interventions and, in general, a smaller sample size is needed to answer a fixed number of research questions.

We will work with maize farmers in eastern Uganda. The interventions will take the form of short videos that explain simple yet effective ways to increase maize productivity, such as best practices in row spacing and soil fertility management. To test hypotheses related to the gender composition of the receivers, we will show the video to either the man, the woman, or man and women as a couple within the household. To test hypotheses related to the gender composition of the messenger(s), we will produce different versions of the video. In one version of the video, the information will be provided by a male farmer. In a second version of the video, the same information will be provided by a woman farmer. Finally, in a third version of the video, the information will be provided by a couple. To test hypotheses related to the way extension information is delivered, the videos will be augmented with an alternative technology to deliver the same information, such

as Interactive Voice Response (IVR). This intervention will be implemented using an encouragement design, where part of the sample is encouraged to use IVR through an “IVR starter kit”. In addition, part of the farmers that get an IVR starter kit will also be reminded to use the IVR through SMS messages. Power calculations suggest we would need a sample of about 3,600 farmers, which will be drawn from five districts. The videos will be shown twice to farmers, once before the beginning of the main season, in July 2017, and once in the beginning of the season, in August 2017. An endline survey will collect information on outcomes after harvesting, which ends in January 2018.

The results of this study are expected to inform agricultural extension system reform currently underway in Uganda. Uganda’s new director of extension services signaled a role for digital extension within the future extension strategy. While the subject of this research study is maize intensification, we believe the findings can be readily extended to include similar interventions on different crops and subjects, including extension on post-harvest storage and handling, agricultural value addition, or agricultural commodity marketing.

Research Questions

This research aims to answer which combinations of attributes of an agricultural extension information campaign are most important for its effectiveness. In particular, we will investigate attributes related to the gender composition of both messenger and recipient of the messages. We will also look at the model of delivery of the information.

Gender Composition of Messenger and Receiver

Even though research suggests that the gender composition of both messenger and receiver are important in making information effective, agricultural extension information services are generally biased toward men. Most often, extension officers are male who target the main decision maker with respect to agriculture within households, which is also often assumed to be the male farmer. The assumption that extension messages targeting one household member will trickle down to the rest of the household, including women and younger household members, may be false. Men do not necessarily discuss

production decisions or transfer extension knowledge to women household members, especially if extension messages focus on men’s priorities and crops (Fletschner and Mesbah, 2011). Gender-matching effects, where men learn more from other men and women learn more from other women, have also been documented in the context of agricultural extension services (Doss and Morris, 2001). At the same time, farms are essentially run and managed at the household level, and it may therefore be more effective if information is both provided and targeted at this level. For instance, Lambrecht, Vanlauwe, and Maertens (2016) found joint male and female program participation leads to higher adoption rates of fertilizer in Eastern DR Congo. The analysis of a Digital Green project in Ethiopia also concluded that there is much to be learned from observing the interactions between men and women who learn about the same technologies and practices (Bernard et al., 2016).

We therefore will investigate the relative importance of (i) the gender composition of the messenger, and (ii) the gender composition of the audience for effective agricultural extension information delivery to encourage sustainable crop intensification in smallholder household farms and for improving gender equity in household farming. In particular, our research is designed to answer the following research questions:

- Does promoting a household approach to farming improve outcomes at the household level? This research question will be answered by comparing outcomes of households that are shown a video with agricultural extension information where the messenger is an individual farmer (ie. a man or a women) to outcomes of households that are shown a video providing the same information by a couple of farmers (ie. man and woman who are shown to participate as equals in the family farm and deliver the message as a couple);
- Does reducing agricultural extension information asymmetries within the household improves outcomes at the household level? This research question will be answered by comparing outcomes of households in which one adult individual (husband or wife) is shown a video with agricultural extension information to outcomes of households in which the couple (husband and wife together) is shown a video with the same information;
- Is there evidence of a gender-matching or homophily effect? This research question will be answered by comparing outcomes of households

where the gender of the messenger and the recipient of the message is matched (male-male and female-female) to outcomes of households where the gender of the messenger and recipient of the message differs (male-female and female-male).

We expect gender composition of receiver and messenger will affect aggregate household productivity. However, we also expect that gender composition of receiver and messenger will have an impact at outcomes within the household. To assess the effect the gender composition of messenger and receiver on equity within the household, we will therefor also look at the following research questions:

- Does reducing information asymmetries related to agricultural extension information within the household increase equity? To answer this question, we will compare the gender productivity gap within households where the video was shown to a man to the gender productivity gap in households where the video was shown to the woman.
- Does projecting a cooperative approach to agriculture increases equity. To answer this question, we will compare the gender productivity gap within households that were shown the video that promotes a household approach featuring a couple versus to the gender productivity gap in households that were shown a video where a single farmer provides the messages.
- How effective is a the interaction of both projecting a cooperative approach and reducing information asymmetries? To answer this question, we will compare the gender productivity gap within households that are both shown the video that also projects a household approach to the gender productivity gap in all the other households.

Finally, in addition to the questions above, we check if the variation in gender or recipient or messenger yields better equity outcomes than the status quo, which is defined as a man receiving agricultural extension information from another man.

Different ICT Channels

The tools and technologies through which the information is transferred are also likely to influence effectiveness. There are many different ways of deliver-

ing agricultural extension messages through ICT. Broadly, one can differentiate between two different approaches. In one approach, extension resembles the traditional teacher-pupil model, where the farmer is assumed to absorb knowledge from experts. Showing videos to farmers or pushing information over a mobile phone would fall into this category. A second approach relies more on a consultative model, where the farmer is assumed to know his or her information needs and requests this information from a service provider. Call centers and Interactive Voice Response (IVR) technology are examples of this second approach. Furthermore, recent research suggests that reminders through SMS is also effective to increase agricultural technology adoption (Casaburi et al., 2014). We will therefore also augment the IVR with SMS messages that remind the farmers of the existence of the IVR service.

We will assess the effectiveness in terms of changing knowledge, practices and outcome such as yields and poverty of different modes of information delivery. In particular, we will:

- Establish if video is a cost-effective medium to transfer agricultural extension information at the farm household level. To test this, we will compare outcomes of farmers that were shown an agricultural extension video to outcomes of households that are shown a random video (placebo).
- Is it cost effective to add a demand driven service to extension using video? To test this, we will compare outcomes of farmers that were shown an agricultural extension video to outcomes of households that were shown an agricultural extension video and were also encouraged to use IVR.
- What is the effect of adding reminders at set time intervals to the video and IVR system? To answer this questions, we will compare farmers that were encouraged to use IVR to farmers that were encouraged to use IVR and also got weekly SMS messages reminding them of the existence of the IVR option.

Outcomes

The effectiveness of the different interventions will be judged by their effect on a range of outcomes at the farm household level. Therefore, a first set of

outcomes will investigate changes in knowledge due to the interventions. We will include changes in knowledge about the existence and use of inputs or practices that were featured in the video (such as proper spacing of maize, application rates of fertilizer, ect). We will also add some practices that are not directly shown in the video, as farmers that were shown the video may be motivated to search for additional information. It is further assumed that knowledge translates into increased adoption of the technologies. We will investigate both adoption as a binary outcome (yes/no), but also look at adoption intensity (adopted on share of total cultivated area, application rates, etc.). Again here, we will not only include practices shown in the video, but also record adoption of other practices or inputs (such as pesticide use), as our interventions may crowd in other intensification investments (Emerick et al., 2016). This, in turn, is expected to benefit agricultural production and yields. Finally, we expect this will affect household well-being, through increased consumption and income derived from marketing of more, better quality crops. We will therefore include consumption expenditure as a proxy for well-being.

Defining the outcomes will enable us to compare the cost-effectiveness of each intervention (e.g., is showing a video to a woman more cost effective than showing it to a couple in terms of its effect on aggregate productivity?). We will also collect information that we think may be affected by focusing on gender aspects in agricultural extension. For instance, showing extension information to the wife instead of the husband may result in substantially different labor allocation effects: women may prefer labor saving innovations, while men may focus more on yield increasing investments. Most of the outcomes we will collect information on will also be disaggregated by gender. For instance, we will estimate maize yield for male managed and female managed plots separately. We will also include IFPRI's Abbreviated Women Empowerment in Agriculture Index (A-WEAI) (Alkire et al., 2013).

Research Strategy

The questions will be answered through field experiments, where farmers are randomly assigned to a group that receives particular interventions (?). An identification strategy that is based on randomization allows us to quantify the causal linkage between an intervention and the outcomes.

The research design itself will take the form of a mixed level factorial

design. To answer the first set of research question related to gender, we define two different factors, each with three levels. The first factor relates to the messenger and has three levels (male, female and couple). Similarly, the second factor relates to the recipient of the message and also has these three levels (male, female and couple). The second set of research questions related to the way the information is brought and where video is now augmented with a demand-driven technology such as IVR, can be answered by adding an extra factor with three levels (no IVR, IVR, IVR + SMS reminders). In practice, we therefore add the IVR treatment to two thirds of the participants who get to see a video, of which half also gets weekly sms reminders to use the service. Finally, we will add a pure control group to the design, such that we do not only investigate relative effectiveness of the different attributes, but also relative to households who receive no information at all. As we want to make sure the impact is related to the actual information given in the video, and not merely a result of being shown a video, households in the control group will also be shown a “placebo” video. The information in this video will be completely unrelated to maize farming.

The video interventions will be targeted at the individual (household) level as opposed to group screenings. Videos will be shown in the house or in the field if necessary. This option guarantees consistency in the treatment and requires the least amount of observations. The videos will be developed in-house by the research team and produced by a professional media production company (NOTV).

For IVR, we will partner with Human Network International. Human Network International’s 3-2-1 develops content for farmers on a variety of crops and agricultural practices, and make the service available to farmers for free or at low cost. Since it is not possible to exclude farmers from this service and at the same time, we can not force farmers to use the service, we will use an encouragement design for this factor. In particular, to two thirds of the sample that gets to see a video, a small leaflet with information on how to use the IVR system. In addition, we will also send SMS messages to half of this last sub-group to remind them of the existence of the IVR service.

Crop and Technology

The interventions for which we will investigate various attributes will be on maize intensification. Maize is widely consumed, yet its value to weight

	subcounties	parishes	villages
Bugiri	11	72	399
Mayuge	13	73	462
Iganga	16	82	360
Namayingo	7	43	308
Namutumba	7	37	348
Total	54	307	1877

Table 1: Administrative structure of study area

ratio is sufficiently high to also make it an important traded commodity. Therefore, increasing maize productivity at the farm household level has the potential to lead to improvements in both nutritional outcomes and income. Maize yields in Uganda are relatively low. While on-station trials report potential yields average about 1.6 metric tons per acre (improved varieties, no fertilizer used), according to Uganda National Household Survey 2005/06 data, average maize yields are much lower, at about 618 kg per acre for the main growing season of 2014. There is a lot of variation in yields, with the top 10 percent of best farmers getting yields in excess of 1.1 metric tons per acre. At the same time, the use of modern inputs such as inorganic fertilizer and modern technologies such as row planting is very low in Uganda. For example, use of inorganic fertilizer averages only 1 kg of nutrient per hectare per year, compared to Kenya (32 kg/ha); Rwanda (29 kg/ha); and Tanzania (6 kg/ha).

Sampling

Sampling Frame

Maize is especially important in the East. We will sample from five districts from the East known for their maize production: Bugiri, Mayuge, Iganga, Namayingo and Namutumba. Table 1 gives an idea of how villages are distributed over parishes, which are in turn allocated to sub-counties within each of the 5 districts in our study. From this, we removed town councils and also two sub-counties that consisted of islands in lake Victoria.

Our study population consists of maize farmers within this region¹. Because of cost considerations, however, we will use two-stage cluster sampling to obtain a representative sample of this population. In particular, we will first randomly select parishes (proportional to the number of villages within each parish). In the selected parishes, all villages will be included in the study. Within each village, we will then list all the households, from which we will then sample households to be included in the study. At the same time, we suspect that outcomes within villages will be correlated, for instance due to local weather conditions, or development programs that are implemented in certain areas. We therefore use the village as a blocking factor. In other words, in each village, we will make sure all possible treatment combinations related to the first 2 factors (plus control) are administered. From the research strategy above, we learn that the messenger and the recipient factor combine into 9 different treatment combinations. Adding the control leads to 10 different households being selected in each village. Of the 9 farmers that are shown a video (not the placebo), about six will also receive the IVR treatment, and of these six, three will also get SMS reminders.

Statistical Power

We determine total sample size and the distribution over the different treatment combinations by running various power analyses for different research questions with different underlying assumptions. Instead of determining power analytically, we use simulation techniques. Simulation allows us to sample from actual data on outcome variables instead of from a theoretical distribution with an assumed mean and standard deviation, which is a much more intuitive way to think about statistical power. It is straightforward to build in flexibility, such as allowing for more than one treatment arm, or account for sampling design effects such as cluster sampling. The algorithm that was used to perform the power calculations can be found in the git repository.

The power calculations will be based on the first set of research questions (related to the gender composition of the messenger and the receiver), as

¹In particular, given the subject of our study, we target monogamous households that are growing maize. However, we will also include some female headed households as they are an important disadvantaged group in Uganda. These households can obviously not get all the treatments. For these households, we will only vary the messenger between male and female.

the hypotheses we formulate here are most complex and the factorial design allows us to recycle observations for the third factor related to adding IVR to the video content. Not only do the two gender related factors in the first set of research question both have three levels (male, female, and both), we are also interested in particular interactions between the two factors, which needs to be accounted for when determining sample size. For the third factor, which is the one related to adding IVR to the video treatment, we are only interested in the main effects (video, video + IVR, and video + IVR + SMS reminders). Therefore, in this factor, we can allocate two thirds of the households where whoever gets to see any kind of video to the video+IVR factor and half of this subgroup can then be allocated to the video + IVR + SMS treatment. For reasonable effect sizes, these sample sizes are likely to result in sufficient power to estimate the additional effect of this form factor.

A first set of power calculations takes **maize yield** data, taken from the Uganda National Household Survey of 2005/06, as the outcome variable of interest². In particular, we calculate average yields of maize (defined as quantity harvested per acre planted) during the second season of 2005 at the household level³. We will sample from this distribution for the power calculations. Mean yields in the sample was about 618 kg per acre with a standard deviation of about 314 kg per acre. This is within the 500-800 kg per acre range that experts estimate for maize yields in the area. We will also use location data from this study to get an idea of the impact of stratifying at this level on statistical power⁴.

The corresponding expected effect size for each treatment combination on average yield at the household level is summarized in Figure 1. They range from a 2.5 percent increase in the treatment combination where the receiver is the woman and the messengers is a male to a 22.5 percent increase for the treatment combination where the video messages is shown to the couple

²While this is a relatively old survey, there are several reasons why we preferred it to more recent similar surveys such as the Uganda National Panel Survey of 2013/14. The most important reason relates to sample size. The UNHS interviewed about 6,000 households, while the UNPS interviewed only about 3,000 households.

³Using the UNHS may provide an overly pessimistic view. Experience with UNHS data suggests the yield data is very noisy. In addition, the first set of power calculations we used in the funding proposal for this study, we relied on rice yield data we collected ourselves in a previous study. When this data was used, we obtained substantially higher power for the same hypotheses.

⁴We use UBOS's Enumeration Area (EA) as location indicators, which is the lowest level of location identification that is provided without jeopardizing anonymity.

and the message in the video is also delivered by a couple. The expected effect sizes are based on the interplay of the three effects mentioned above: gender-matching effect, a knowledge-is-power-effect, and the projection of a household approach. For instance, the effects on the diagonal in Figure 1 are higher because here the gender-matching effect is playing: We expect that men learn more from other man and women learn more from other women, which in turn would lead to higher yields among gender matched sub-samples. In Mozambique, Kondylis et al. (2016) find evidence of such a gender-matching or homophily effect in extension, whereby women seem to learn more from fellow women and men seem to learn more from fellow men. In particular, they find that among men who received information about pit planting from a male messenger, the proportion that are aware of pit planting is 10 percentage points higher. Female awareness, knowledge and adoption is the same irrespective of the presence of a male extension worker. However, if a female extension worker is added, awareness, knowledge and adoption among female farmers also increases by roughly the same proportions. We use these results to get an idea about the gender matching effect. In particular, we model a 10 percent increase in yields when messenger and recipient is male. We model a slightly lower effect when messenger and recipient is female (7.5 percent), as it is assumed women have less bargaining power within the household.

The expected effects emanating from the gender composition of the receiver is related to information deficiencies and a knowledge-is-power-effect. This can be directly when women are also important actors in agriculture. For instance, Kabunga, Dubois, and Qaim (2012) find that female farmers are less likely to adopt the tissue banana culture technology in Kenya, but that they would have an equal chance to adopt innovations, provided that they acquire sufficient knowledge about the innovation. In addition, knowledge may also affect technology adoption and subsequent yields through changes in relative bargaining power of the actors. Doss and Morris (2001) find that adoption of agricultural technologies among female farmers is lower than among male farmers. They find that this is due to gender-linked differences in access to complementary inputs. We assume only a very small effect (+1 percent) for the subset of women farmers who get to see a video where the messenger is a male farmer. As this effect seems to be partially driven by differences in access to complementary inputs, we assume a higher effect when the recipient is male and the messenger is female (+1.5 percent).

Lambrecht, Vanlauwe, and Maertens (2016) investigate the effect of par-

Control 618 kg/ac		Messenger		
		Male	Female	Both
Receiver	Male	680 kg/ac 10.0 %	649 kg/ac 5.0 %	711 kg/ac 15.0 %
	Female	627 kg/ac 1.5 %	664 kg/ac 7.5 %	695 kg/ac 12.5 %
	Both	711 kg/ac 15.0 %	695 kg/ac 12.5 %	742 kg/ac 20.0 %

Figure 1: Expected Effect Sizes for Yields

ticipating in extension training as a couple. They investigate whether participation of female farmers in an agricultural extension programme in South-Kivu increases adoption of three technologies: improved legume varieties, row planting and mineral fertilizer. In their study, joint male and female programme participation leads to the highest adoption rates. In particular, they find that joint participation increases fertilizer adoption by 12.4 percentage points and row planting by about 13.4 percentage points. While these are large effects (as both fertilizer use and row planting is very rare in the area, with baseline percentages of 6 percent for fertilizer and 10 percent for row planting), the non-random nature of the intervention may lead to overestimation of the couple effect, and we model much more modest effects. Also, due to the homophily effect and the fact that man are likely to have more power within the household, we model a slightly higher increase in case the messenger is a man (+15 percent) versus a woman (+12.5 percent).

We also expect larger effects when the message is brought by a couple, as a cooperative approach to farming is likely to reduce inefficiencies within the farm household. We are not aware of previous studies that may guide us in the choice of the effect size, but we think it may be similar to the effect of providing training to a couple. We again expect this effect to be highest when the receiver is a male, due to his bargaining position (+15 percent). We expect only a slightly lower effect when the women receives this treatments (+12.5 percent). Finally, when information inefficiencies are addressed by showing the video to both husband and wife and a cooperative approach to farming is projected, we expect the largest effect (+20 percent)

At one extreme, we could calculate sample size required to identify all possible treatment combinations in Figure 1. For instance, we could com-

pare control to MM (receiver=male and messenger=male), control to FM (receiver=female, messenger=male), control to MF (receiver=male, messenger=female), etc. Testing all interactions in this way would require a sample size well above what is possible. We therefore will base our power calculations on comparing groups of treatments based on the research questions outlined above.

A first question will be to find out if videos work, irrespective of who they are shown to and who features in the video. We thus calculate sample size by taking all treatment arms and interactions in our factorial design together and we simply test the difference between treatment and control⁵. This scenario is shown in the first graph in Figure 2. We find that we would be able to obtain sufficient power (.8) with a total sample size of about 500 observations. As these observations need to be balanced across the different cells in the experiment, this means we need at least 250 observations in the control and a minimum of 28 observations in each treatment combination.

A second question relates to the gender-matching or homophily effect. This is likely to be the most stringent, as it is based on only 4 cells of the design in Figure 1. In particular, we will compare cells with matched gender (MM and FF) with those that have opposite sexes matched (MF and FM). The results for these power calculations are shown as the second figure in the first row in figure 2. We find that we would need about 1,800 observations in total to get sufficient power. This means that in each of the relevant cells (MM, FF, MF and FM), we would need about 450 observations.

A third question relates to the gender composition of the recipient. Here, we want to compare a situation where there is information asymmetry (where only one of the spouses was shown the video) to a situation where both husband and wife received the same information (both were shown the video), irrespective of who features in the video. In this case, we use all cells in figure 1 and compare BB, BF and BB to MM, MF, MB, FM, FF and FB. The results of these power simulations is shown as the third graph in figure 2. We find that we obtain sufficient power for a total sample size of 2,000 observations. Taking into account the fact that we need to balance across all other cells, we need 333 observations in each of BB, BF and BB and 167 observations in each of MM, MF, MB, FM, FF and FB.

⁵In the control, the placebo video also needs to be shown to somebody. We also make sure there is balance here between man, woman and couple. As such, in one third of our control, the placebo video is shown to the man, in one third it is shown to the female and in one third it is shown to the couple.




Figure 2: Power Curves for Maize Yield

A fourth question is related to the gender composition of the animator, where we expect that when the message is brought by a couple, and a household cooperative approach is thus promoted, leads to better outcomes than when the message is given by an individual (either a male or a female farmer). We calculate sample size that is needed to determine a significant difference in yields between these two groups and plot this in last figure in Figure 2. The fact that we expect relatively large effects from a cooperative approach means we obtain sufficient power with a total sample size of about 2150 observations. Again taking into account balance across other cells mean we need at least 359 observations in each of MB, FB and BB and at least 180 observations in each of the other cells.

A second set of power calculations was done for an outcome variable that captures equity within the household. The **gender productivity gap**, where women managed plots are farmed less extensively than male managed plots resulting in significantly lower yields, has been observed throughout Africa South of the Sahara and for a variety of crops (Udry, 1996). In Uganda, Ali et al. (2015) find that female managed plots are almost 20 percent less productive than male managed plots. The gap is caused by a general lack of access to production factors compared to men. For instance, women may be forced to farm on inherently lower productivity plots and they may have less access to productivity enhancing tools, technologies and inputs such as fertilizer. Knowledge related to crop intensification is one such factor where access between men and women differs, and so finding out how changing particular attributes to extension videos affect this gap is an important outcome.

Using data from the latest Uganda National Panel Survey (2012-2013), we also find a significant gender gap among maize farmers in Uganda. Male farmers obtain yields of about 1427 kg per hectare. Women managed maize plots only get about 1385 tons. We thus model a difference of about 42 kg per hectare, with a standard error of about 40 kg per hectare⁶. The difference we find seems lower than what others find. For instance, ? find the gender gap to be about double of what we find, although they base their analysis on male versus female headed households, while we look at the gender of the person that manages the plot. According to FAO, female managed plots are in general 20-30 percent less productive. Ali et al. (2015) estimate the gap to be 23 percent, but they aggregate all crops by weighing by prices and

⁶In this case, the simulations are not based on draws from actual data, but from a normal distribution with mean 42 and standard deviation of 40.

Control 42 kg/ha		Messenger		
		Male	Female	Both
Receiver	Male	48.30 kg/ha +15.0 %	46.20 kg/ha +10.0 %	43.05 kg/ha +2.5 %
	Female	37.80 kg/ha -10.0 %	33.60 kg/ha -20.0 %	35.70 kg/ha -15.0 %
	Both	43.05 kg/ha +2.5 %	40.95 kg/ha -2.5 %	28.00 kg/ha -33.3 %

Figure 3: Expected Effect Sizes for Gender Productivity Gap

expressing yields in monetary terms. Peterman et al. (2011) estimate it as high as 50 percent.

There is no literature we are aware of that looks at the effect of the gender composition of the messenger and the gender composition of the receiver of agricultural extension messages on the agricultural productivity gap. Therefore, we will use our best judgment to determine expected effect sizes. Modeled effect sizes are summarized in Figure 3. Starting from a baseline situation where maize yields are about 42 kg/ha higher on male managed plot, showing a video to a man only is likely to further increase this gap. We therefore simulate a 15 percent higher yield gap in this group. Due to the homophily effect, we expect that the increase in male productivity is smaller when the messenger is a woman (as the message will be less effective), and so the gap will increase less (+10 percent). Finally, if the message is brought by a couple and a household approach to farming is projected, the man may be encouraged to share some of the knowledge with his wife. However, we expect the increase in yield on male managed plots to still be higher than any potential increase of yield on the female managed plot. As such, we expect the effect on the yield gap to be marginally positive (+2.5%).

When the information is shown to the woman but not to the man, it is expected that yields on female managed plots will increase, while yields on male managed plots are likely to remain the same. This would mean that the gap reduces. Again due to the homophily effect, we expect women to learn most from women, and so the reduction in the gap will be highest when the messenger is also a woman (-20 %). The reduction will be smallest when the information is given by a man (-10 %). Finally, if the woman is sensitized on the importance of a household approach to farming, some of

the information may be shared with the husband, also reducing the effect on the gap somewhat (-15 %).

When the information is given to the couple, we expect little or no effect on the gap if the message is given by either male or female. We may expect a slight increase in the gap if the information is given to the male (+2.5%) and a slight decrease if the information is given by a female (-2.5%). Finally, if both spouses have all the information and the message is given that the household farm should be managed as a unit, we expect that the gap reduces by a third (-33.3%).

With respect to intra-household outcomes, we are also interested in four hypotheses. First, instead of comparing the treatment to control, we will compare the status quo (where a man is targeted by a male extension worker) to any other treatment combination. Second, we want to test the importance asymmetric information, and we will test if the gap significantly differs between households where the man was shown the video and households where the woman was shown the video. Third, we want to test if there is a difference depending on whether a household cooperative approach was promoted or the information was brought by an individual. Finally, we also want to separately compare changes in the gender yield gap for households that are both shown the video that also projects a household approach to all the other households.

We again ran a series of power simulations corresponding to these four hypotheses. The results are presented in Figure 4. To be able to answer the first question with sufficient power, we need a total sample size of about 600 observations. Of these, at least 300 observations should be in the MM cell, and about 38 observations in each of the other cells. To answer the second question, we need about 500 observations. These need to be equally divided over the first two rows of the matrix in figure 3, meaning at least 83 observations in each cell. To answer the third question, we need a total of about 1,100 observations. These observations should be distributed as follows: at least 183 in each of the cells in the last column and at least 92 in each of the other cells. The final question can already be answered with less than 300 observations. Of this, 150 should be in the BB cell, and about 20 in each of the other cells.

On the basis of the above, we come to the following ideal division of observations across the different treatment cells : {ctrl=250, MM=450, MF=450, MB=359, FM=450, FF=450, FB=359, BM=333, BF=333, BB=359}, leading to a total number of observations of 3,793. However, due to various con-




Figure 4: Power Curves for Gender Productivity Gap

straints, we will relax power for the most stringent research hypothesis, which is the second question relating to the gender-matching or homophily effect. If we only require 60 percent power, we can reduce sample size for MM, MF, FM and FF to 375, leading to a sample size of about 3,500 observations in total. To account for attrition, we add another 100 observations to each group according to their weight in the total, {ctrl=257, MM=385, MF=385, MB=369, FM=385, FF=385, FB=369, BM=342, BF=342, BB=369} or 3,588 observations.

Assignment to Treatment

We use cluster sampling to draw our sample. We randomly select parishes (within subcounties within the five districts) proportional to the number of villages within each parish. We include all villages within each parish. We use village as a blocking variable. That is, within each village, we randomly allocate up to 10 households to any of the 10 treatment combinations defined by the messenger gender composition and recipient gender composition factors plus control. Two third of them also receives the IVR encouragement and half of those that also get the IVR treatment will also get weekly SMS reminders that promote the IVR service. The code that does the sampling and treatment allocation can be found [here](#). The resulting sample with treatment allocation can be accessed [here](#).

Fieldwork

Calendar

In the East, there are two maize cropping seasons. For our study, we will concentrate on the second maize growing season, which runs from about August to January. In August, fields are prepared and planting is done in September. First weeding happens around the end of September or the beginning of October. By late October, a second round of weeding takes place as maize starts to flower. Mid to late December, harvesting starts. In higher areas, harvesting can go on until mid January. During the second season, farmers prefer early maturing varieties, as the rains are short. However, in general, early maturing varieties have lower yields. The full cycle from planting to harvesting in lowland areas takes about 3 to 3.5 months. In high altitude areas this can take up to 5 months. In our study, we will concentrate on

lowland areas.

Barriers to maize intensification

The treatments consist of the provision of information that is assumed to increase maize productivity through encouraging adoption of modern technologies and recommended practices. The goal was to select those practices and technologies that are likely to have the largest impact on yield, household income and intra-household equity. The topics to be included in the video script were obtained from interviews with key stakeholder and experts, such as maize farmers, value chain actors, maize breeders, extension workers and other government staff such as the district agricultural officer. These interviews took place in May 2017.

The main factors affecting maize productivity are related to pests, poor soils, and poor seed quality. The three most important **pests** that affect maize farming are Striga, maize stalk borer, and now also Armyworm. Striga (in particular the Striga hermonihica variety), also called witchweed, is a parasitic plant that feeds off the roots of maize. Doing so, it starts to draw nutrients from the host, leading to severe stunting of the host crop. Since most of the damage is already done by the time the parasite emerges from the ground, weeding is not very effective. In addition, each Striga plant produces up to 50,000 seeds, which remain dormant in the soil until new maize is planted. Striga can reduce production from about 30 percent up to total loss of the crop. No single method is effective to control striga. A new hybrid seed called Longe 7 HR, boosted with a herbicide, is said to be less affected by striga, but it is expensive at 10,000 UGX/kg and does poorly under dry conditions and in areas where the rains are short such as in the East⁷. Crop rotation with sweet potato or beans can reduce striga infestation. Furthermore, (additional) inorganic fertilizer use has also been recommended to counter striga. It has been observed that striga proliferates when soils are poor. In addition, one can make up for the nutrients drained by the parasite by boosting the maize plant with DAP or NPK. Weeding, especially the first time the striga comes up before it has had the chance to dig into the roots of the maize, is recommended, also to reduce the spread of seeds.

The stemborer is actually a moth that is active during night, laying eggs on maize plants. It is the larva of the moths that cause the damage. They

⁷In the East, early maturing varieties are preferred, such as Longe 10. In the future, it a similar HR strain of Longe 10 is expected to be released.

enter inside maize stem, where they feed and grow for about three weeks. If larvae enter the maize plant at an early stage, this can kill the plant. At a later stage, since larvae make extensive tunnels inside the stem, the maize plants weaken. The stem may break, or the maize plants may fall over. Maize stemborer can reduce production by between 20 and 40 percent.

The fall armyworm is relatively new in Uganda⁸. As the stemborer, the damage is done by a larva from a moth. Currently, it is present in about 20 districts. In some districts, up to 40 percent of maize is affected. Yield losses are estimated at 10 to 15 percent nationwide. However, experts fear this is only the beginning, and armyworm may overtake stemborer as the most destructive pest. Armyworm is very difficult to control. One expert had some success with a chemical called Dudu Accelamectin, which required spraying only once and hence is cheap. However, other experts warned that the effectiveness of one single type of chemical depends on many factors such as location, type of seed used, etc. Other chemicals such as Rocket were also mentioned to be somewhat effective.

Poor soils, exhausted by continuous cultivation with little rotation, is a second main limiting factor. Soils lack macro-nutrients such as nitrogen (N), phosphorous (P) and potassium (K). Farmers rarely use organic fertilizer. Experts say it is virtually impossible to get decent yields without using inorganic fertilizer. In particular, at planting stage, DAP or NPK should be used in the soil. After about 4 weeks, a nitrogen based fertilizer such as Urea should be applied on the topsoil near the plants to boost vegetation. Sometimes it is also advised to split the application of urea, administering half a doze at 4 weeks and half a dose at tasseling stage. Fertilizer can easily double yields. Especially used in combination with improved seeds, the effects of fertilizer are significant.

Within Africa, farmers mainly rely on saved **seeds**, and maize in Eastern Uganda is no exception (McGuire and Sperling, 2016). However, as maize is cross pollinated, recycling leads to fast degeneration. However, hybrid seeds such as Longe 7, Longe 9 are available and have also been distributed through Operation Wealth Creation. Hybrid seeds potentially triple yields. In addition to hybrid seeds, which need to be procured every seasons as they can not be recycled, the government also released Open Pollinated Varieties (OPV). These are improved varieties that can be reused for about 3 or 4

⁸Armyworm was discovered in Uganda in the first season of 2017. However, experts told us that it was also present in the second season of 2016 but was mistaken for stemborer.

seasons. In general, they are lower yielding than hybrid seeds. Still, yield are likely to be about 30 to 40 percent higher than when saved seeds are used, up to 100 percent if OPVs are also used with fertilizer. OPVs are generally recommended to poorer farmers as they can be reused as seeds to some extent.

Most hybrid seeds take long to mature. Farmers also complain that hybrids do not withstand drought very well. However, new varieties have entered the market that are both high yielding and early maturing. For example, UH5354, marketed as Bazooka yields about 20 percent more than other hybrids and is drought resistant. Farmers that use these seeds report up to 40 bags of maize per acre. Today, yields are about 7-8 bags per acre. Similarly, there are several OPVs on the market that are also early maturing.

Finally, there are also **agricultural practices** that affect yields, but the effects are generally smaller than those related to improved seeds and inorganic fertilizer use. The most important are:

- **Timely planting:** maize should be planted at the onset of the rains. However, many farmers delay planting for several reasons. For instance, they may not be certain that the rains have really started. There may also be bottlenecks in land preparation, where the poor are often the last in line for the use of oxen or tractors. It is estimated that each day a farmer delays planting, yields reduce by 1.5 percent.
- **Plant spacing:** Plant that are too close to each other compete for light and nutrients, reducing yields. Plants that are too far from each other means space is wasted, also leading to lower yields. Related is the number of seeds that farmer plant in each hill. The recommended practice is to put in two, but farmers want to be sure of germination and put in more, sometimes up to 5. If they all germinate, there is competition for nutrients and sun. Usually there is also not enough thinning, or this is done too late when competition has already occurred. One expert told us that standard spacing is 75cm x 60cm with two plants per hill. However, they now recommend 75cm x 30cm with 1 plant per hill. The expert estimated the latter results in a 35 percent higher yield than the former.
- **Weeding:** The first weeding should be at 18-20 days after planting at the three leaf stage, and the second weeding at 2-3 weeks after first weeding. From emergence to 8 leaves (knee height), maize plant is a

very poor competitor and unwanted stress such as weed competition must be prevented. Weeding must therefore be done within that period. A third weeding is optional. It is only dependent on weed pressure after the second weed. If needed, it should be done at tasseling stage. Weeding removing Striga at early stage before it can attach to the roots of the maize can also have a significant impact on yields. In addition, removing striga before it flowers reduces the spread of the pest. Proper weeding can increase yield by about 50 to 70 percent.

Instruments

We will use video to convey information to farmers with the aim of changing adoption behaviour. We will focus on providing information on seed selection, soil nutrient management (including promoting organic fertilizer application), weeding, timely planting and plant spacing⁹. From our conversations with experts, we learn that many farmers may already be aware of the existence and use of these technologies or practices. Therefore, simply providing information about the existence of modern technologies and recommended practices and on how to use them may not be sufficient to change behavior. Often, access and affordability was mentioned as a problem¹⁰. In our videos,

⁹As mentioned above, pest control also ranks high among technologies that increase yields. However, we felt uncomfortable promoting chemicals in our videos that are potentially hazardous to humans. In Uganda, Endosulfan, DDT, Glyphosate, Linden and others—are still being used. Many of these are on the World Health Organization’s list of banned chemicals and blacklisted by the Stockholm Convention on Persistent Organic Pollutants.

¹⁰Farmers indeed often mention they do not use modern inputs because they have no money, suggesting liquidity is the main constraining factor, and thus instead of providing information, a more effective strategy would be to provide credit or subsidize inputs. Recent research, however, finds that despite what farmers claim, liquidity is often not the most important constraining factor. Duflo, Kremer, and Robinson (2011) note that in Kenya, fertilizer is relatively cheap and can be bought in small quantities, putting it in reach of even the poorest households. They suggest the main reason for low fertilizer adoption rates should be attributed to the farmer’s failure to commit. Ashraf, Gine, and Karlan (2009) find that lack of credit was not the main reason why farmers did not produce high-value export crops, and that farmers that did produce export crops found access to credit on their own. Emerick et al. (2016) find that increasing access to drought tolerant seed that reduced downside risk increased the use of credit from existing sources. Finally, Karlan et al. (2014) compare outcomes related to agricultural production between households that were given cash to households that were provided with insurance and find that when provided with insurance, farmers are able to find resources to increase

we will thus also try to alter the belief that seeds and fertilizers are “too expensive” by pointing out the costs and benefits of the different technologies and practices we promote¹¹. In addition, we will encourage farmers to start small, using fertilizer and seeds on a small area of their field to experiment and see for themselves, and reinvest in subsequent years. Inter-temporal decision making, where costs today have to be compared to uncertain outcomes in a distant future, is often challenging for poorly educated farmers¹². Furthermore, Duflo, Kremer, and Robinson (2011) point out farmers may have difficulties committing to fertilizer use in Kenya. In our video, we attempt to make farmers aware of this, and suggest some techniques to overcome this bias. We also pay considerable attention to the way the message is delivered. For instance, the message is brought by “peer farmers”, as it is found that farmers find communicators who face agricultural conditions and constraints most comparable to themselves to be the most persuasive (BenYishay and Mobarak, 2014). The information is also presented as a success story, which is assumed to affect a range of non-cognitive farmer characteristics such as aspirations, locus of control and self-esteem (Bernard et al., 2015).

The video starts with a farmer (a male farmer, a female farmer, or a couple, see below) introducing themselves. He talks about how he used to struggle with his maize gardens and how at one point in time, he decided things needed to change. It is shown how the farmer sells a hen, and obtains a small loan from a friend. This money is then used to buy small quantities of improved seed and fertilizer in a local shop. It is then shown that, before planting the improved seed, the farmer prepares the garden. He is shown collecting manure wherever he can find it, and applies it to a small corner of the field of 20m by 20m. Next, it is shown in detail how the maize seeds are spaced 75cm x 30cm with 1 plant per hill and how the DAP should be applied. The viewer is reminded to plant in time. The next scene depicts the field after about 10 to 12 days when the maize has emerged from the ground.

expenditure on their farms.

¹¹The fact that farmers do not engage in a cost benefit analysis, where inputs are used to generate output, was illustrated by a complaint made by one of the farmers. He complained that farmers get low prices. To him, it did not make sense to buy seed at 6000 per kg and sell maize at only 300-400 per kg.

¹²Under-investment caused by low perceived returns in the future has also been documented in the demand for schooling (Jensen, 2010). A recent experiment investigates if rice farmers in Uganda are aware of the returns to different recommended rice growing practices such as early transplanting and fertilizer use (?).

At this stage, it is recommended that the farmer engages into gap filling to replace seeds that did not germinate with new seeds to preserve optimal plant density. The next shot shows the field at 18 to 20 days after planing, when first weeding is done. Particular attention is paid to identification of striga in an early stage. It is also advised to weed again two to three weeks later. The next scene zooms in on urea fertilizer application. Here, the field is shown at about 4 weeks after planting when the maize is knee high. It is shown how Urea topsoil dressing should be applied. Finally, it is recommended to do one more round of weeding around the tasseling stage of the maize.

We then spend some time comparing the costs to the benefits the different improved inputs and recommended practices. For fertilizer and improved seed, the costs for one acre is calculated and compared to the value of what is harvested. The profit is then compared to the value of what would have been harvested on that acre without improved seed and fertilizer. This would be less than half of what the profit would be when improved inputs are used. For recommended practices, we report what the expected yield increase would be if, for instance, recommended spacing was used or weeding is done in time.

Next, we try to promote a long run perspective, where the farmer is encouraged to start small (one tenth of an acre) and grow bigger over time. We also pay some attention to the commitment problem. We advise the farmer to, at time of harvest when the farmer sells most of his maize, immediately go to the farm supply store and purchase seeds and fertilizer, and store this in a safe place. If inputs are not available, farmers are encouraged to keep the money needed to buy the inputs in a separate, labeled container. It is sometimes argued that such mental accounting can be an effective commitment mechanism (Dupas and Robinson, 2013). In a last scene, the farmer recapitulates and once more directly addresses the viewer and encourages him or her to try this as well.

A total of three such videos will be produced. They will be identical in terms of the information that will be in the video, but they will differ according to the gender of the messenger, corresponding to one of the factors in the factorial experimental setup. In one version of the video, the actor will be a man. In a second version of the video, the actor will be female. In a third version, the video will feature a couple as the messenger. Here, both male and female will feature in the different scenes, and the person talking to the viewer will be altered between the man and the women. Finally, we will use a placebo video for the control. This will be a neutral video on a non-related topic, such as the tourist potential of the region. Farmers will

be shown any of these four videos according to random treatment allocation. Videos will be shown twice to each farmer in the sample, once before planing (July) and once immediately after planting (August).

The videos will be embedded in a small Computer Assisted Personal Interviewing (CAPI) survey application implemented in Open Data Kit (ODK). The application will start by asking some identifying information about the farmer and recording consent. It will also be used to collect some basic farmer characteristics, which can be used to investigate balance at baseline (see the section on balance below). The application will also embed the video, which will be cued to make sure the correct video is shown to the right farmer. Finally, information will be gathered on viewing conditions, and some information will be collected to see if farmers understood the information that was given in the video.

The IVR system consists of a toll free number that can be called by the farmer. Farmers that are assigned to this treatment will be given the toll free number and leaflet that provides some more information about the system. When calling the IVR system, the caller is greeted in Lusoga and a tutorial is provided within the system. The caller is presented with a menu and can choose to learn more about different aspects of maize farming, such as soil management, seed selection, pests and diseases, post harvest practices,... After selecting the desired option, the farmer hears a conversation between two farmers, where one farmer explains recommended practices to another farmer. While the IVR system has more information than the videos, we made sure that the information that we provide in the videos is the same as what is provided in the IVR system. The SMS messages that will be sent out to half of the farmers that are assigned the IVR treatment will simply be a reminder of the existence of the service and the phone number they can dial to access it.

After harvesting of maize in January or February 2018, an farmer level endline survey will collect information on a range of outcomes related to learning, adoption behaviour, maize production and farmer well-being more generally. We will also inquire about intra-household issues. This endline survey will also be implemented as CAPI, either using ODK, SurveyBe or Survey Solutions.

Empirical Analysis

Variables

We will evaluate effectiveness using the following variables. We will also drop outcomes from our analysis for which 95 percent of observations are the same value. This is done to reduce the influence of outcomes with limited variation. For continuous variables, 5 percent trimmed distributions will be used¹³.

- knowledge: this will be assessed using short multiple choice “quiz” type questions. The result will be binary variable (correct or not) and proportions will be compared between groups using both Chi-squared tests for difference in proportions and linear probability models.
- adoption of practices, dis-aggregated by gender of main manager and assessed at plot level. Recommendations are those that were shown in the videos:
 - organic fertilizer used (yes/no)
 - quantity of organic fertilizer used (kg) and application rates (kg/acre)
 - fertilizer use Urea (yes/no)
 - quantities of fertilizer used Urea (kg) and application rates (kg/acre)
 - fertilizer use DAP (yes/no)
 - quantities of fertilizer used DAP (kg) and application rates (kg/acre)
 - use of improved seed, defined as either hybrid or OPV (yes/no)
 - quantity of improved seed used (kg) and application rates (kg/acre)
 - recommended spacing of maize used (yes/no)
 - planted in time (assessed as the square of the time difference between onset of rains and time of planting)
 - practiced gap filling (yes/no)
 - practiced weeding according to recommended timing and frequency (yes/no)

¹³When new variables are derived, trimming is done on the end result, that is $\text{trim}(\text{kg}/\text{acre})$ instead of $\text{trim}(\text{kg})/\text{trim}(\text{acre})$

- secondary effects:
 - hired labour for maize farming (yes/no)
 - quantity of hired labour (man-hours)
 - use of herbicides/pesticides/fungicides (yes/no)
 - quantity of herbicides/pesticides/fungicides (kg) and application rates (kg/acre)
 - access to credit during the last 6 months (yes/no) and amount accessed (UGX)
 - Agricultural advice sought on maize farming in the last 6 months? (yes/no)
 - use of labour saving technology such as oxen, power tiller,... (type, use)
- production: at household level and also dis-aggregated by gender of main manager and assessed at plot level.
 - log of maize yields in kg per acre
 - maize yield gap within household (difference in kg/are on women managed plot and male managed plot)
 - log of maize production in kg
 - area of maize cultivated (acre)
 - area of maize cultivated as a share of total area used for cultivation (percentage)
 - labour allocation decision (women/men/children hours worked in maize production)
- Welfare outcomes
 - welfare: consumption expenditure per capita, based on a selection of most consumed items.
 - poverty: consumption expenditure relative to the official poverty threshold.
 - market participation: was maize sold (yes/no), quantity sold, price at which maize was sold (to capture quality differences), and to who the maize was sold.

- women empowerment in agriculture (assessed by a standard index (Alkire et al., 2013))
- Non-cognitive characteristics: We also want to investigate if viewing successful farmers alters attitudes and aspirations, such as viewing farming as a business and changes in locus of control:
 - Do you consider maize farming your main business? (yes/no)
 - Do/did you want your children to become maize farmers? (yes/no)
 - Do you agree that chance determines what happens in your life? (likert scale 1-5)
 - Do you agree that other people such as your wife/husband, pastor, other family members, local leader etc determine what happens in your life? (likert scale 1-5)
 - When you make plans, do you stick to them? (likert scale 1-5)
 - Do you agree that important events (good and bad) that happen in your life were meant to happen? (likert scale 1-5)
 - Do you feel you have mostly good qualities? (likert scale 1-5)
 - Do you agree that, poor people are poor because they do not work hard enough to improve their lives? (likert scale 1-5)
 - In farming do you usually adhere to the goals you have set out? (likert scale 1-5)
 - Do you sometimes spend money and later admit or realize it wasn't worth it? (likert scale 1-5)
 - When things look bad, do you usually expect the best? (likert scale 1-5)
 - Are you a person that likes to try out new things even if you know there is a high chance things go wrong? (likert scale 1-5)

Balancing Checks

While we are not planning to do a dedicated baseline survey, we will ask some questions before the first intervention to investigate balance. The choice of variables was based on what variables other researchers in similar studies use in their orthogonality tests. In particular, we looked at balance tables in

studies that investigate the adoption of yield improving methods and technologies using RCTs. These studies include Duflo, Kremer, and Robinson (2011), Karlan et al. (2014), Ashraf, Gine, and Karlan (2009) and Bulte et al. (2014). In addition, we also add some variables that may be useful to investigate heterogeneity in treatment effects. In particular, we collect some household characteristics such as household size, age and education level of both wife and husband. We then ask more specific questions related to maize farming, such as acreage and by whom this is primarily managed, as well as quantities produced in the last season. Furthermore, we ask if the household received agricultural extension, whether improved maize seeds were used, and whether fertilizer was applied. We also collect data on housing conditions (material of wall and number of bedrooms) and access to off-farm income. The complete baseline questionnaire can be found on github.

Treatment Effects

Intent to Treat

For the research questions related to gender composition of messenger and recipient, we assume full compliance, and as such assume that the average treatment effect (or Intention tot Treat effect) is equal to the Treatment of the Treated effect. Therefor, we will simply compare means between the different groups in a regression framework. In particular, we identify the treatment effects (β) by estimating the average treatment effect of an outcome variable (y):

$$y_h = \alpha + \beta T_h + \varepsilon_h \quad (1)$$

where T is an indicator function that is one if household h was in the treatment group and zero otherwise. We will control for the gender of the enumerator and also add parish fixed effects. In addition, in a factorial design, orthogonal treatments are likely to increase the variance of the outcome variable. As such, we also need to control for the treatment status of the other factors (and add dummies for, for instance IVR and IVR+sms). For the treatments where compliance can not be completely controlled, we also control for ex post outcomes derived from the endline survey (eg. an indicator if the man used IVR, an indicator if the woman used IVR,...). For the few outcome variables that are also collected at baseline such as fertilizer use

or use of improved seed (see balancing checks above), difference-in-difference models will be estimated.

It is important to realize that treatment and control groups need to be balanced over all other factors. This means that often, we can not use all observations. For example, if we want to simply test if being shown any video was effective in increasing maize yields, we can not simply compare the 257 control households that were shown a video to all households that were shown a video (3,331 households). We need to make sure that within the treatment group, there are equal numbers in each cell. As such, we need to take the treatment combination with the least observations, which are according to the power calculations the combinations where the video is shown to the couple and the messenger is either a man or a women (BM and BF). In these cells, we need at least 342 observations. Therefore, to test the effectiveness of videos, we compare 257 control households to 3,078 households, which corresponds to 342 households in each of 9 treatment combination cells. Therefore, in each cell that has excess observations, we will take a random sample. For example, according to the power calculations, we will have 385 households in the treatment combination where the messenger is male and the recipient is male (MM). We will take a random sample of 342 from these 385 households.

Treatment on the Treated

For the research question related to different ways to deliver extension information, we compare videos to IVR. As mentioned above, we have much less control over compliance in the IVR treatment. We therefore estimate the treatment of the treated effect using two stage least squares. In particular, we estimate:

$$y_h = \alpha + \beta C_h + \varepsilon_h \quad (2)$$

Where C_h is an indicator that takes the value of one if the household called the IVR system and zero otherwise. This indicator C_h will be instrumented by an indicator that represents the allocation of the household to the IVR treatment T_h .

Heterogeneous Effects

While we did not consider heterogeneous treatment effects in the power calculations, there are some groups of people we expect our treatments will be more or less effective for. One key group of farmers we want to focus on is youth. On the one hand, youth appear to be less interested in the rural livelihoods of their parents. In Uganda, the youth often abandon the countryside as soon as they can to work in Kampala. On the other hand, the youth may be more interested in video and conversant with the IVR tools we promote. We also expect that the “farming as a business” message we try to convey appeals more to young people. Age categories (i.e. a youth indicator dummy variable) will therefore be constructed on the basis of age data collected before the first intervention and treatment indicators will be interacted with the youth dummy.

We will also run separate analysis depending on whether the household has access to income from sources other than farming. Often, such households differ substantially from households that only rely on farming in terms of access to capital and ability to manage risk. We therefor also collect data on access to income from other sources at baseline and estimate different treatment effects for both types of farmers.

Third, our videos heavily promote the use of modern inputs. Inputs are costly, something we try to address in our videos by advising to start small and grow over time. However, the ability to use the inputs also depends on availability. Therefore, we will also look at heterogeneity in the treatment effect that is related to the distance to the nearest farm supply shop. This is measured at baseline using a continuous variable, which will be interacted in a regression with the treatment variable.

Finally, we will also investigate how education levels of both the men and woman in the farm household interacts with the effectiveness of our various treatment combinations. This is again done by constructing different categories (eg. less versus more educated households) and comparing treatment effects.

Standard Error Adjustments

While our main outcome variables are ultimately household welfare and maize yields, we will also estimate the impact of our interventions on a range of intermediate variables as indicated above. The fact that we have many

such variables may lead to the so-called “look elsewhere” effect, where one is bound to find significant effects simply due to the sheer number of hypotheses that are tested. Therefore, some form of multiple-inference correction is in order. In general, there are two ways in which to reduce the likelihood of false positives that result of multiple hypothesis testing. One can either reduce the number of hypotheses, or one can make the statistical test stricter by for instance reducing the significance threshold (such as the popular Bonferroni adjustment). We will address false positives arising from multiple hypothesis testing using two approaches.

First of all, we will use the groupings presented in the section that lists the variables to create indices (knowledge, adoption, secondary effects, production, welfare,...). At the most basic level, each of the indices is a weighted mean of the several standardized outcomes within each group. In particular, for each variable within each group, we make sure positive direction always means better, otherwise we switch sign. We then demean the outcome and standardize by scaling by the (pooled) standard deviation. We then create weighted averages for the outcomes in each group at the household level, using as weights the inverse of the co-variance matrix of the transformed outcomes within the group. This is done for each of the groups. The resulting variables can then be used to assess the impact of the particular intervention using the specifications outlined above.

However, we may be interested in identifying differential effects within each of the groups. For example, we may want to differentiate between the effect of our intervention fertilizer application and the effect of our intervention on improved seed adoption. We will therefore also use Family Wise Error Rate Control. In particular, we will use the free step-down re-sampling method proposed in Westfall and Young (1993).

References

- Ali, D., D. Bowen, K. Deininger, and M. Duponchel. 2015. *Investigating the Gender Gap in Agricultural Productivity: Evidence from Uganda*. Washington, DC: World Bank.
- Alkire, S., R. Meinzen-Dick, A. Peterman, A. Quisumbing, G. Seymour, and A. Vaz. 2013. “The Women’s Empowerment in Agriculture Index.” *World Development* 52: 71 – 91.

- Ashraf, N., X. Gine, and D. Karlan. 2009. "Finding missing markets (and a disturbing epilogue): Evidence from an export crop adoption and marketing intervention in Kenya." *American Journal of Agricultural Economics* 91 (4): 973.
- BenYishay, A. and A. M. Mobarak. 2014. *Social Learning and Communication*. NBER Working Paper 20139, National Bureau of Economic Research, Cambridge, MA.
- Bernard, T., S. Dercon, K. Orkin, and A. Seyoum Taffesse. 2015. "Will video kill the radio star? Assessing the potential of targeted exposure to role models through video." *World Bank Economic Review* 29 (sup 1): S226.
- Bernard, T., S. Makhija, K. Orkin, A. S. Taffesse, and D. J. Spielman. 2016. *Video-based agricultural extension: Analysis of a pilot project in Ethiopia*. Project note, International Food Policy Research Institute, Washington, DC.
- Bulte, E., G. Beekman, S. Di Falco, J. Hella, and P. Lei. 2014. "Behavioral responses and the impact of new agricultural technologies: Evidence from a double-blind field experiment in Tanzania." *American Journal of Agricultural Economics* 96 (3): 813.
- Casaburi, L., M. Kremer, S. Mullainathan, and R. Ramrattan. 2014. "Harnessing ICT to increase agricultural production: Evidence from Kenya." *Unpublished working paper*.
- Doss, C. R. and M. L. Morris. 2001. "How does gender affect the adoption of agricultural innovations? The case of improved maize technology in Ghana." *Agricultural Economics* 25 (1): 27 – 39.
- Duflo, E., M. Kremer, and J. Robinson. 2011. "Nudging farmers to use fertilizer: Theory and experimental evidence from Kenya." *American Economic Review* 101: 2350–2390.
- Dupas, P. and J. Robinson. 2013. "Why don't the poor save more? Evidence from health savings experiments." *American Economic Review* 103 (4): 1138–71.

- Emerick, K., A. de Janvry, E. Sadoulet, and M. H. Dar. 2016. "Technological innovations, downside risk, and the modernization of agriculture." *American Economic Review* 106 (6): 1537–61.
- Fletschner, D. and D. Mesbah. 2011. "Gender disparity in access to information: Do spouses share what they know?" *World Development* 39 (8): 1422–1433.
- Humphreys, M., R. Sanchez de la Sierra, and P. van der Windt. 2013. "Fishing, commitment, and communication: A proposal for comprehensive non-binding research registration." *Political Analysis* 21 (1): 1.
- Jack, B. K. 2013. *Market inefficiencies and the adoption of agricultural technologies in developing countries*. White paper, Agricultural Technology Adoption Initiative, Cambridge, MA.
- Jensen, R. 2010. "The (perceived) returns to education and the demand for schooling." *Quarterly Journal of Economics* 125 (2): 515.
- Kabunga, N. S., T. Dubois, and M. Qaim. 2012. "Heterogeneous information exposure and technology adoption: The case of tissue culture bananas in Kenya." *Agricultural Economics* 43 (5): 473–486.
- Karlan, D., R. Osei, I. Osei-Akoto, and C. Udry. 2014. "Agricultural decisions after relaxing credit and risk constraints." *Quarterly Journal of Economics* 129 (2): 597.
- Kondylis, F., V. Mueller, G. Sheriff, and S. Zhu. 2016. "Do female instructors reduce gender bias in diffusion of sustainable land management techniques? Experimental evidence from Mozambique." *World Development* 78 (C): 436–449.
- Lambrecht, I., B. Vanlauwe, and M. Maertens. 2016. "Agricultural extension in eastern Democratic Republic of Congo: Does gender matter?" *European Review of Agricultural Economics* 43 (5): 841.
- McGuire, S. and L. Sperling. 2016. "Seed systems smallholder farmers use." *Food Security* 8 (1): 179–195.
- Peterman, A., A. Quisumbing, J. Behrman, and E. Nkonya. 2011. "Understanding the complexities surrounding gender differences in agricultural

- productivity in Nigeria and Uganda.” *Journal of Development Studies* 47 (10): 1482–1509.
- Udry, C. 1996. “Gender, agricultural production, and the theory of the household.” *Journal of Political Economy* 104 (5): 1010–1046.
- Westfall, P. H. and S. S. Young. 1993. *Resampling-based multiple testing: Examples and methods for p-value adjustment*, vol. 279. John Wiley & Sons.

Script for Video

Scene 1: Opening scene: A male farmer stands in front of a well managed, high yielding maize field with big cobs and introduces himself. Farmer in view, talks to the audience.

Hello everyone. My name is ____ and I am a maize farmer, just like you. I used to struggle with the same problems as many of you, and as a result, my gardens produced barely enough maize to feed my family. However, over the years, I have learned how to increase the number of bags of maize I can get from my gardens.

I started using modern inputs such as hybrid seeds and inorganic fertilizers and started managing my garden following recommended agricultural practices such as improving my soil with organic manure, optimal spacing of the maize plants and timely planting and weeding.

Since I had little money, I started small. However, season after season, I re-invested the extra money I made and after a few seasons, I managed to use modern inputs on my entire field. I now have sufficient maize to feed my family and even sell on the market!

Do you also want to grow more maize? Let me show you how I did it!

Scene 2: In front of a Striga infested, poorly looking maize field. Farmer in view, talks to the audience.

This is what my garden used to look like. My maize performed poorly because I kept using recycled seed. The soil was exhausted because I kept growing maize season after season without using fertilizer. I thought farming using recommended plant spacing was tedious and cumbersome, and weeding was a waste of time. I often delayed planting. After one particular bad harvest, my family suffered badly. It was at that point that I sat down and decided things needed to change: I needed to grow more maize on the land that I had.

I realized that improving my soil with organic manure would not cost me money, only labour. I also recalled being told that I could increase my yields through timely planting, optimal spacing of the maize plants and frequent weeding.

Scene 3: In the village, show farmer selling a chicken to a neighbor and shaking hands with a friend upon securing a loan. In a small shop that sells farm supplies, show how the farmer buys 1 kg of “bazooka”, 6 kg of DAP and 6 kg of Urea. Make sure the prices are clearly indicated on the products (bazooka 8,500 per kg, DAP 3,000 per kg, urea 2,500 per kg). Here the farmer is again acting and information is given as a voice-over.

It was more challenging to get improved seeds and inorganic fertilizer. However, in the end, I managed to obtain enough money by selling a cock to my neighbour and borrowing a little from a friend. I used the money to buy 1 kg of hybrid seed, 6 kg of DAP and 5 kg of Urea. These supplies were sufficient to produce maize on an area of 5 by 5 sticks, or one tenth of an acre of my field. I thus choose a corner of my field where I would farm according to recommended practices.

Scene 4: Show how the farmer collects cow dung from his chicken and from public grazing land or the road in a bag. Show how the farmer throws mango peels/banana peels/sweet potato peels... on the field. Show how this is plowed into the land.

Before planting the improved seeds, when I was preparing the land, I applied organic fertilizer such as animal droppings or green manure from plant residue that I collected.

Scene 5: In the small corner of the field, focus on good agricultural practices. Show each practice as detailed as possible! Eg, show how the farmer measures the spacing using something that is typically about 30 cm and about 75 cm. Show the farmer making a hole of 10 cm deep and adding 1 water bottle cap DAP in the hole after which he puts some soil on top. Show the farmer adding one improved maize seed in the hole and covering it with soil. Please consult with Wiberforce on how to do these things. Again, farmer acts and voice-over explains what farmer is doing.

At planting time, I decided to use recommended spacing, carefully measuring one foot between plants and 2 and ½ foot between rows. I first made a hole of about 4 inches deep and added 1 water bottle cap of DAP. I then added some soil on top of it. I then put one improved maize seed in it and cover with soil. It is also very important to not delay planting: immediately after the onset of the rain, start planting.

Scene 6: Show a field where the crop is just emerging to illustrate gap filling. Farmer acts and voice-over explains what farmer is doing.

After 10 to 12 days, carefully check if all seeds have germinated. To keep optimal plant density, it is important to resow where seeds did not germinate.

Scene 7: Field at three leaf stage, focus on weeding. Show how a striga looks like when really young. And show a flowering striga so everyone recognizes the plant. Show the farmer removing weeds and striga. Farmer acts and voice-over explains what farmer is going on.

Weeding is particularly important in the first 4 weeks. Start weeding 18 to 20 days after planting. It is particularly important to remove Striga at this stage. In this early stage, the Striga did not reach the roots of the maize plant yet, so damage can still be avoided. Try to remove the Striga before it flowers to avoid it from spreading to the rest of the garden. A second round of weeding should be done at 2 to 3 weeks after the first weeding.

Scene 8: Show a field that is knee high. Show the farmer applying 1 water bottle cap of urea around each plant according to recommended practices. Farmer acts and voice-over explains what farmer is doing.

About 4 weeks after planting, or when the plants are knee high, apply Urea. Apply 1 water bottle cap of urea around each plant.

Scene 9: Show a growing/ flowering field and show how the farmer is removing weeds. Farmer acts and voice-over explains what farmer is doing.

Weeding should be done one last time around the tasseling stage of the maize. And that's it really.

Scene 10: Doing the math – any ideas to make this engaging are welcome! Here the farmer can address the audience again. Can we use visual aids to make this appealing to low educated farmers???

I used to think that farming maize according to recommended practices would be a lot of work and cost me a lot of money. But inorganic fertilizer is essentially free, I just needed to collect it and throw it on the garden. Applying inorganic manure is done only once while preparing the field, proper spacing is done while planting and weeding is done three times. But soon I realized the increase in maize production due to these recommended practices was well worth the extra effort!

Similarly, I used to think that modern inputs such as improved seed and inorganic fertilizer were too expensive. But once I compared the expenditures to the additional maize I would get in the future, I found out these inputs are not expensive at all!

Let us do the math together. To cultivate one acre using improved seeds such as bazooka, I need about 10 kg of seed. Bazooka sells at about 8500 per kilogram, so I would need 85,000 UGX to use Bazooka on an entire acre. In addition, I would need 60 kg of DAP and 60 kg of Urea. DAP is sold at 3,000 shillings per kg and Urea is sold at 2,500 shillings per kg. This means 180,000 shillings for DAP and 150,000 shillings for Urea to use fertilizer according to recommendations on an entire acre. Thus, to use modern inputs on one acre of land, I would need a total of about 395,000 UGX. But look at the result of this investment! Using these inputs, I will be able to harvest about 25 bags from this same acre. For each bag, I can get about 50,000 UGX, so the value of my harvest is 1,250,000. Subtracting the cost of the fertilizer and seed, this means I remain with a profit of 855,000. Now let us compare this to a situation in which I did not use modern inputs: With recycled seed and without fertilizer, I used to get only 8 bags from this acre. Selling these at 50,000 each would mean my harvest is worth 400,000. Remember with modern inputs my profit is 855,000: This is more than double!

Similarly, I used to think that adhering to recommended spacing was tedious and weeding a waste of time. But also here, there are important returns to these practices. Take for example timely planting: each day you are late in planting, your yield will reduce. For instance, if I would normally get 8 bags from one acre, planting one week too late will result in only getting 7 bags. Plant spacing also has a big effect on production. I noticed that by using one foot between plants and 2 and ½ foot between rows as optimal spacing, I could increase my yield per acre from 8 bags to 10 bags. The same holds for keeping the garden free of weeds, as maize is poor at competing. I found out that I can get an extra 2 bags of maize per acre if I weed three times and pay particular attention to removing stirgas at an early stage.

Scene 11: Include flashbacks: Maybe show the farmer selling his chicken and buying small amounts seeds and fertiliser in shop again and then show him in front of the small corner of the plot

I know farming according to these recommendations mean more work, and that 400,000 is a lot of money! But you can always start small, like I did. Remember I started in a corner of my garden of only 5 by 5 sticks? The improved seeds, DAP and Ureas required to cultivate on this area was only 41,500. But on that small plot, I was able to get 2.5 bags, for which I got 125,000. My profit was 83,500.

Remember that when I would not have used recommended practices, fertilizer and improved seed on this small plot, I wouldn't even have been able to harvest one full bag of maize! I would have made only 40,000 shillings.

Scene 12: can we show the farmer with some money in hands, in front of a somewhat bigger plot (quarter acre) and in a shop buying somewhat larger quantity improved seeds, DAP and urea.

Over time, I would use the extra money I made by using improved inputs to buy more seed and fertilizer. On my small plot of 5 sticks by 5 sticks plot, I made a profit of 83,500 shillings. If I would not have used modern inputs, I would have had 40,000. So I reinvested the difference, 43,500 and bought more improved seeds and chemical fertilizer. This allowed me to cultivate almost a quarter

of an acre using modern inputs in the following season, generating a profit of more than 170,000. I used the extra money again to buy even more modern inputs. Pretty soon, I was able to cultivate an entire acre using improved seeds and fertilizers.

Scene 13: Start by letting farmer talk to audience. Then show the farmer selling a bag of maize and going immediately to the store to buy inputs. Show the farmer putting money in a container that has a clear label on it reading “seed and fertilizer”.

From experience, I also learned that it is often difficult to commit to re-investing my money. Even though at harvest, I tell myself I will use part of the money to buy fertilizer and improved seed, there are many other competing issues that require money and I would often find myself without money when it is time to start planting maize again.

Therefore I now plan well ahead. During harvest, I calculate how much seed and fertilizer I need, and I immediately go to the shop to buy the inputs and store them in a safe place in my house. If the inputs are not available in the store, I put the money aside in a special container and regularly visit the store to see if the inputs are there. In this way, I am sure that when time for planting comes, I have the seeds and fertilizer and can plant timely.

Scene 14: Show the farmer in front of the nice field again, where he recapitulates (maybe throw in a few flashbacks?). Maybe also show the varieties in the store, with a clearly visible price tag. Eg Longe 10 H @ 6000/kg, Bazooka @8,500/kg,...

So, to recapitulate, to become a successful maize farmer, it is first and foremost important to get good improved seed. Preferably, get high yielding hybrid seeds that are early maturing, such as longe 10H or UH5354 (Bazooka). Note that such hybrid seeds can not be recycled: you need to purchase new ones every year. If you can not afford this, try at least to buy Open Pollinated Varieties such as Longe 5 (Nalongo) or Longe 4. These are cheaper and can be reused three or four seasons. Second, apply fertilizer to the soil. Use organic fertilizer from animals or crop residue when preparing the field. During planting, DAP should be used in the soil. When the maize is about knee high, urea should be applied around the plants as top dressing. Fertilizers not only boost maize production, it also reduces Striga infestation. Combining this with timely planting, proper spacing of 1 foot by 2,5 foot and keeping the garden free of weed, you are bound to get much more maize! I strongly encourage you to try this as well. Start small and see for yourself! Good luck!

Mzee Musa: Now the land is ready – uhuh – do I now start planting?

Mr. Onesmus: Yes, it is time for planting, but you must always remember – you need to choose the right seeds if you want to get high returns on your harvest.

Mzee Musa: But maize is maize, I can plant anything, right?

Mr. Onesmus: There are different varieties of maize. If you want a good harvest, make sure you use high yielding varieties for better yields.

Mzee Musa: What kind of varieties are there?

Mr. Onesmus: Quite a few.

For hybrids, buy early maturing seeds from an approved agro-input store every growing season, such as *longe 10H* or *UH5354* (*Bazooka*).

Then for open pollinated varieties such as the early maturing *Longe 5* (*Nalongo*) or *Longe 4*, you can also get these from approved agro-input stores or use saved seeds from the previous harvest. But remember to change seed after 3 or 4 seasons.

Mzee Musa: Ok, I have the seeds – do I now plant?

Mr. Onesmus: Yes, you can plant now – however, For good establishment, plant your maize after two heavy rains at the beginning of the rain season. Do not delay planting as it will reduce yields.

Mzee Musa: But how much seed do I need?

Mr. Onesmus: You will need at least 10Kg of seed to plant 1 acre of land.

Mzee Musa: Do I just throw the seeds anywhere in the garden?

Mr. Onesmus: No. You will need to plant in lines at a spacing of 2½ ft between lines and 1 ft between plants. Place 1 seed per hole for optimum plant population

Mr. Onesmus: For your maize to grow well, you will need to maintain good soil health by using fertilizer from animal refuse and crop residue or fertilizer manufactured from industries

Mzee Musa: But when and how do I apply it?

Mr. Onesmus: Apply farmyard fertilizer from animal droppings or green manure from plant residue or both during land preparation. Alternatively, you can use manufactured fertilizer from industries.

Mzee Musa: And the fertilizer?

Mr. Onesmus: Apply 1 bottle-top full of DAP per hole and cover with little soil before placing the seed.
You will need at least 60kg of DAP to cover 1 acre.

Mzee Musa: Do I apply fertilizer just once and that's it?

Mr. Onesmus: No, you will need to apply again.
Apply 1 bottle-tops around the plant when the plant is knee high. You will need 60kg of urea to plant 1 acre. But is also advisable that you always consult with your agricultural extension worker about your soil nutrient needs before use of fertilizers. For best results however, use both organic and inorganic fertilizers in combination

Mr. Onesmus: Weeds compete for water, nutrients and space with your maize, so weeding is very important, especially during the first 4 weeks. Start weeding 18 to 20 days after panting. Then again 2 to 3 weeks later.

Mzee Musa: Is there any particular weed I should look out for?

Mr. Onesmus: Striga. It grows in close association with maize and can be identified by its dark green leaves and purple or white flower. The most basic and easiest way to get rid of this weed is by hand pulling striga plants early before they flower. *In this early stage, the Striga did not reach the roots of the maize plant yet, so damage can still be avoided. Removing Striga before it flowers also prevents it from spreading to the rest of the garden.*

SMS messages

1. Okungula duma mungi bwosimba ensigo enongosemu okusinga ngosimbye ensigo yaduma wabulidho-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (159)
2. Simba duma mumabanga ga futi 1 mubikolo ku futi 2.5 mulaini ngokozesa mpeke 1 mubuli kinnha-Kuba kwisimu eritasasuza 0200522420 owabulwe ebisingawo ku duma (157)
3. Enaku 10 ku 12 ngomaze okusimba duma, irizaamu empeke buli awatamera munimiro yonayona-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (156)
4. Okudhuba enimiro yaduma kya mugaso inho musasira 4 edhisoka ngomaze okusimba-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (146)
5. Okwanguwa okutoola kayongo munimiro ngakali muto kikendeza kubulabe bwatusa kuduma-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (154)
6. Oludhuba lwokubiri lwetagisa musasira 2 ku 3 oluva-inhuma lwoludhuba olusooka-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (147)
7. Duma bwatuka mumaviivi pima YUREYA akasandikira 1 akokucupa omute kubuli kikolo ngokyetoloza-Kuba kwisimu eritasasuza 0200522420 owabulwe ebisingawo ku duma (155)
8. Duma bwatandika okutoluyo oluyange enimiro eba yetagisayo oludhuba lulala olusembayo-Kuba kwisimu eritasasuza 0200522420 owabulwe okusingawo ku bya duma (154)

Translations

You will get much more maize if you use hybrid seed instead of recycled seed – Call the maize hotline on 0200522420 free for more advise!

Maize should be spaced 1 foot between plants and 2.5 feet between rows, 1 seed per hill – Call the maize hotline on 0200522420 free for more advise!

10 to 12 days after planting your maize, resow where seeds have not germinated – Call the maize hotline on 0200522420 free for more advise!

Removing weeds from your maize garden is very important in the first 4 weeks -
Call the maize hotline on 0200522420 free for more advise!

Removing striga early, before it flowers, can reduce much of the damage - Call the
maize hotline on 0200522420 free for more advise!

A second round of weeding should be done about 2-3 weeks after the first weeding
- Call the maize hotline on 0200522420 free for more advise!

When your maize is knee high, apply 1 water bottle cap of urea around each plant -
Call the maize hotline on 0200522420 free for more advise!

A final round of weeding should be done when the maize starts tasseling - Call the
maize hotline on 0200522420 free for more advise!