

Diagnostic performance of quantitative measures from FDG-PET/CT, FEC-PET/CT and DW-MRI in the detection of lymph node metastases in endometrial and cervical cancer.

Ben King¹

¹The Institute of Genetics & Cancer, The University of Edinburgh.

Contributing authors: b.g.king@sms.ed.ac.uk;

Abstract

This will be an absrract test!

1 Introduction

The prognosis for both cervical and endometrial cancers is associated with presence of lymph node metastasis. In endometrial cancer the 5-year disease free survival is estimated at 90% for patients without lymph node metastasis, 60-70% for those with malignant pelvic lymph nodes and 30-40% with malignant para-aortic lymph nodes [1]. In cervical cancer the 5-year overall survival is estimated at 92% with localised disease and 58% with regional disease such as lymph nodes metastasis [2]. Treatment options differ for patients with metastatic lymph nodes. Patients with localised disease will receive surgery as their primary treatment whereas patients with malignant lymph nodes will typically be referred for chemoradiotherapy.

Accurate staging of lymph nodes is therefore critical in treatment planning and improving prognosis for patients. Although surgical lymphadenectomy is considered the most accurate technique for the detection of lymph node metastases, systematic lymphadenectomy is controversial as the procedure

carries significant risks of lymphoedema, lymphocyst formation, and injury to surrounding organs and tissues [3, 4]. Furthermore, this procedure has not been shown to reliably convey any significant survival benefit and is therefore not widely recommended as a therapeutic option [5]. Accurate pre-operative staging of lymph nodes would therefore have a significant impact on reducing morbidity of unnecessary lymphadenectomy procedures, surgical time, and cost of staging.

Multi parametric magnetic resonance imaging (MRI) is currently considered the optimal imaging technique for local tumour staging of endometrial cancer through assessment of myometrial and cervical invasion but the major limitation of all imaging techniques is poor detection of metastatic lymph nodes [1]. Morphological signs of nodal involvement, including size criteria, on CT and MRI vary and have been found to be unreliable markers of malignancy [6]. Tumour may have spread to non-enlarged nodes (micrometastases) and nodes can enlarge due to benign conditions. Functional imaging techniques that can detect changes in the microenvironment and metabolic properties of tissues pose an alternative strategy for diagnosis of lymph node status.

Integrated positron emission tomography/computed tomography (PET/CT) with ¹⁸F-fluoro-2-deoxy-D-glucose (¹⁸F-FDG), a glucose analogue, allows combined anatomical and functional imaging. Visual diagnosis of nodal status in endometrial and cervical using FDG-PET has been evaluated in a number of small, mainly retrospective, clinical trials. Reported sensitivities of FDG-PET/CT for nodal involvement are heterogeneous (63%-80%) whilst specificities are high (93%-97%) [5, 7]. Low sensitivity of FDG-PET/CT is considered to be the main limitation of the imaging modality due to the limited spatial resolution of PET imaging [8].

Maximum standardised uptake value (SUVmax) is a semi-quantitative measure derived from FDG-PET/CT scans that represents relative glucose uptake in tissues. SUVmax of the primary tumour has been shown to be higher in patients with higher grade endometrial cancer, myometrial invasion, cervical invasion, and lymph node metastasis [9, 10]. Reports on the utility of SUVmax in the detection of nodal involvement are limited and most studies to date have used visual assessment of the level of uptake in nodes compared to background. Kim et al. retrospectively assessed SUVmax in histologically confirmed benign and malignant lymph nodes and reported moderate sensitivity (69%) and high specificity (92%) when using an SUVmax cut-off of 2.8 [11].

Diffusion-weighted MRI (DW-MRI) detects changes in the diffusion properties of water in tissues which is primarily influenced by cellular density and the nuclear-to-cytoplasm ratio and can therefore enhance subtle underlying abnormalities in tissues. DW-MRI is established for routine use in endometrial cancer staging and is shown to be promising in diagnosing extent of

myometrial invasion adjunct to conventional T2-weighted imaging [12–14]. There has been some research investigating the utility of the quantitative DW-MRI measure, the apparent diffusion coefficient (ADC), in the diagnosis of lymph node metastasis. The measurement of ADC from DW-MRI allows for quantitative assessment of cellular density. Whilst some studies report metastatic nodes have a lower ADC_{mean} than benign nodes, other studies have not suggesting there is a significant overlap in ADC_{mean} values between benign and malignant nodes with no reliable cut-off [15–17]. Furthermore, in endometrial cancer, retrospective studies have also produced heterogeneous results with some showing ADC is significantly different in metastatic and benign lymph nodes and others finding no difference [15, 18].

Choline is a critical component of cell membranes. In rapidly dividing endometrial cancer cell lines increased uptake of choline has been observed [19]. Radiolabelled choline (¹¹C or ¹⁸F choline) can be incorporated with PET for functional imaging of cell membrane metabolism in vivo and has established use in detection of malignant lymph nodes in prostate cancer exhibiting a high specificity and moderate sensitivity [20]. In small feasibility studies, ¹¹C-choline PET has shown similar diagnostic performance to FDG-PET/CT in the identification of gynaecologic tumours [21]. Additionally, ¹¹C-choline PET/CT had high accuracy for nodal staging in uterine carcinoma [22]. Despite positive early findings no large prospective studies have reported the performance of ¹⁸F-fluoroethyl-choline-PET/CT (FEC-PET/CT) in the diagnosis of nodal involvement in endometrial or cervical cancer. Furthermore, no investigations into the diagnostic utility of nodal SUV_{max} on FEC-PET/CT in gynaecological cancers have been conducted to date.

The present study aims to investigate the ability of quantitative measurements from FDG-PET/CT, FEC-PET/CT and DW-MRI in differentiation of metastatic and benign lymph nodes and determine the diagnostic utility of these imaging techniques.

This study forms part of the secondary analysis of the prospective UK MAPPING study [23].

2 Methods

2.1 Patients

This study analyses data from the MAPPING study: a multicentre, prospective study evaluating the diagnostic accuracy of FDG-PET/CT, FEC-PET/CT and DW-MRI in the detection of nodal metastases. A full summary of the Materials and Methods can be found in Rockall et al. (2021); key methods are detailed here [23].

Participants were recruited from 5 gynae-oncology tertiary referral centres in National Health Service (NHS) hospitals in England. Eligible patients were over the age of 18 years and had newly diagnosed, histologically confirmed cervical or endometrial cancer. Age, tumour grade, pre-and post-treatment FIGO stage and the final treatment received were collected.

One-hundred and sixty-two patients were enrolled in the study. Of these, thirty-three cervical cancer and 66 endometrial cancer patients met the primary reference standard of nodal histology, had quantitative imaging data, and were therefore eligible for analysis (Fig. 1).

2.2 FDG-PET/CT and FEC-PET/CT protocol

FDG and FEC PET/CT scans were acquired using a standardised protocol based on UK NCRI PET Research Network guidance at accredited centres. Participants were injected with FDG and FEC and scans were acquired at a median of 60 mins post injection on separate days. Low dose CT was performed followed by emission study (base of skull to upper thighs) as 3D acquisitions. Images were reconstructed using ordered subset expectation maximisation and the PET images were attenuation corrected using the CT data.

2.3 DW-MRI protocol

Standard of care conventional MRI scan for staging was performed as per local protocol which included a minimal dataset of MR sequences. The MRI field of view included the pelvis and para-aortic regions up to the level of the left renal vein. For nodal evaluation, readers had a minimal dataset that included axial T1, axial T2 and axial diffusion weighted MRI (b values 0, 300, 600, 900 and 1200) with associated calculated ADC maps.

2.4 Surgical staging and pathologic evaluation

The primary reference standard was confirmed nodal histology by surgical lymphadenectomy. The surgical lymphadenectomy was undertaken as per the MAPPING study protocol and anatomical location, presence of metastatic disease, and lymph node dimensions in each region (right pelvis, left pelvis, and para-aortic) were recorded [23]. A region was determined to be positive for nodal metastases with the presence of at least one malignant node at histology.

2.5 Image evaluation

All images were read by two central MRI and PET readers; the central reads were co-ordinated through the UK NCRI PET core lab and the trials unit. A third central PET reader reviewed all PET measurements, and recorded additional readings. All readers were accredited radiologists that were core members of the gynae-oncology multi-disciplinary team and/or PET/CT experts. Readers were made aware of the clinical diagnosis (endometrial or cervical cancer) from the pre-operative biopsy results, as per standard clinical

practice. Readers were masked to other imaging studies and the final nodal histology.

2.6 FDG-PET/CT and FEC-PET/CT image evaluation

The anatomical location of any focally increased tracer uptake that was higher than background adjacent tissue corresponding to a node of any size on the CT was recorded. An elliptical ROI was manually placed on the slice with the highest activity and a measurement of SUVmax was recorded. Short-axis and long-axis diameter was recorded for each identified node. SUVmax of the primary tumour was also measured by a single central expert PET reader by placing a regular ROI over the primary tumour on the slice with the highest activity on a dedicated PET/CT workstation (Hermes Medical Solutions).

2.7 DW-MRI image evaluation

Readers were aware of the nodal size from the standard of care MRI, and DW-MRI was additive to the standard conventional imaging. Nodes were identified on the high b-value ($b = 1200 \text{ mm}^2/\text{s}$) DW-MRI as non-continuous high signal intensity (SI) round or ovoid structures that corresponded with a node on the anatomical images. SA diameter on MRI was measured for each identified node. The ADCmean of identified nodes were measured by manually placing a regular ROI over nodes greater than 5 mm short axis diameter, avoiding the margin and noted with anatomical location. ADCmean of the primary tumour, if visible, was also measured in each patient by both readers in an identical fashion.

2.8 Dataset

The highest SUVmax, for FDG-PET/CT and FEC-PET/CT, and lowest ADCmean, for DW-MRI, from the left pelvis, right pelvis, and para-aortic regions from each reader was selected for analysis. In cases where both readers provided measurements in a region, the readings were averaged and included in the final analysis. Where only one reader provided a measurement in a region, this measurement was included in the final analysis. Where no central readers provided a measurement in a region, the third external reader's measurement was included in the final analysis. The mean SUVmax from FDG-PET/CT, SUVmax from FEC-PET/CT, and ADCmean of the primary tumour from both readers was calculated and included in the final analysis. Reads in regions that had corresponding nodal histology were included in the analysis.

Where data for both were recorded, nodal-to-tumour ratio (NTR) was calculated by dividing nodal by tumour SUVmax or ADCmean for all measurement methods. SUVmax:ADCmean ratio (STAR) was calculated by dividing nodal SUVmax by nodal ADCmean for FDG-PET/CT and FEC-PET/CT. Finally, for FDG-PET/CT and FEC-PET/CT, SUVmax normalised to SA diameter (SNSA) was calculated with SA diameter from the corresponding measurement method.

A total of 270 nodes had histology and were included in the final dataset, of which 178 and 82 were from endometrial cancer and cervical cancer patients, respectively. 115 left pelvic, 119 right pelvic, and 36 para-aortic nodes were included in the final dataset. A total of 134, 66, and 154 nodes had SUVmax from FDG-PET/CT, SUVmax from FEC-PET/CT, and ADCmean from DW-MRI respectively.

2.9 Statistical analysis

Statistical analysis was performed in R. Patients were stratified into two cohorts by their diagnosis of endometrial or cervical cancer. Nodes from all regions were grouped together for analysis. Nodal-to-tumour ratios (NTR) were calculated by dividing the nodal quantitative measure (SUVmax or ADCmean) by the quantitative measure from the primary tumour. SUVmax-to-ADCmean ratio (STAR) was calculated by dividing the SUVmax by the ADCmean of the corresponding node. SUVmax normalised to SA diameter (SNSA) was calculated by dividing the SUVmax by the SA diameter of each node for FDG-PET/CT and FEC-PET/CT. The Mann-Whitney U-test was selected to compare quantitative measures in histologically confirmed benign versus malignant lymph nodes. One-way ANOVA with Tukey's post-hoc test was used to determine significance when comparing left pelvic, right pelvic, and para-aortic regions. P-values were adjusted using the false-discovery rate (FDR) method.

Pairwise Pearson correlation coefficients were calculated between features and p-values were adjusted using the false-discovery rate method.

To determine if a regression model could assist in diagnosing benign from malignant lymph nodes, univariate and multivariate logistic regression models were built. Due to the low number of measurements for cervical cancer and FEC-PET/CT, only FDG-PET/CT and DW-MRI data from endometrial cancer was considered when building models. Additionally, due to strong correlation between features, calculated variables (NTR, STAR, SNSA) were excluded as features.

A multivariate logistic regression model with L1 regularisation penalty (LASSO) was employed. Rows with any NAs were omitted prior to analysis, leaving a total of 57 nodes for analysis. Logistic regression predicts a probability of malignancy between zero (benign) and one (malignant). The dataset was randomly split into train and test sets at a 70%/30% split. Features in the training set were scaled between 0 and 1, and the same scaling function was applied to the test set. The 'glmnet' package was used to find the optimal value of lambda by cross-validation. A logistic regression model with alpha set to 1.0 for the L1 penalty was fit to the training data and used to predict probabilities for the test data. Optimal cutoff probability was found through maximisation of the F1-score. This process was repeated 5,000 times to retrieve the average optimal probability cutoffs and lambda. The model was refit using these hyperparameters and inference was run another 5,000 times to estimate model performance.

Logistic LASSO regression shrunk all coefficients of features, excluding SUVmax and SA diameter, from FDG-PET/CT to zero. To determine if simpler models would perform better, univariate logistic regression was undertaken on these features using the 'glm' package in R. Rows with NAs were omitted prior to analysis, leaving a total of 104 nodes for analysis and the dataset was randomly split into train and test sets at a 70%/30% split. Logistic regression models were fit to SUVmax and SA diameter separately. Optimal cutoffs and model performance were determined as with logistic LASSO regression.

Optimal cut-offs for all quantitative measures were determined using the 'cutpointr' package in R. Direction was set to greater than or equal to for all measures except ADCmean and ADCmean NTR where direction was set to less than or equal to. Cutpoints were selected through maximisation of the F1 score. Stratified bootstrapping was repeated 2000 times. Out of bag estimates of area under the curve (AUC), sensitivity, and specificity were calculated.

Receiver operating characteristic (ROC) analysis was applied, using the 'pROC' package in R, for each quantitative measure to generate ROC curves.

Interobserver agreement for each FDG-PET/CT, FEC-PET/CT, and DW-MRI was evaluated by calculating intraclass correlation coefficients (ICC) for endometrial and cervical cancer cohorts separately using the R package 'irr'. A two-way model was used and agreement between readers was evaluated. Confidence level of the interval was set at 95%.

3 Results

3.1 Demographics

The demographics of patients eligible for analysis are summarised in Table 1. A total of 99 patients were eligible for analysis: 33 cervical cancer patients and 66 endometrial cancer patients. 67 and 73 patients received left pelvic and right pelvic lymphadenectomy, respectively, while just 26 patients underwent para-aortic lymphadenectomy. 88% of cervical cancer patients were diagnosed as FIGO stage IB1 and no patients in this cohort had a FIGO stage greater than II. In the endometrial cancer cohort, FIGO stage ranged from IA to IVB (Table 1).

3.2 Quantitative measures in benign and malignant lymph nodes

SUVmax and ADCmean of lymph nodes in endometrial and cervical cancer was analysed. No significant difference was found between regions in SUVmax or ADCmean on FDG-PET/CT and DW-MRI (Figure 1). Lymph nodes of the para-aortic region on FEC-PET/CT had significantly higher SUVmax than left and right pelvis. Interestingly, average SUVmax of all nodes on FDG-PET/CT was significantly higher in endometrial than in cervical cancer (Figure 2).

Quantitative measures, both raw and calculated, were compared in benign and malignant lymph nodes in endometrial cancer. All raw and calculated

Characteristic	n = 99
Median age (range), years	63.5 (24-80)
Median BMI (range), kg/m ²	28 (18-50)
Histology	
Cervical (n = 33)	
Adenocarcinoma	11 (33%)
Adenosquamous	6 (18%)
Squamous cell	15 (45%)
Unknown	1 (3%)
Endometrial (n = 66)	
Endometrioid	35 (53%)
Serous/clear cell	26 (39%)
Other	2 (3%)
Mixed (2 or more)	3 (5%)
Surgical Procedure	
Laparoscopic	42 (42%)
Robotic	31 (31%)
Open	26 (26%)
Lymphadenectomy	
Left pelvic	67 (68%)
Right pelvic	73 (74%)
Para-aortic	26 (26%)
FIGO stage (on MRI)	
Cervical (n = 33)	
IB1	29 (88%)
IB2	1 (3%)
IIA1	2 (6%)
IIB	1 (3%)
Endometrial (n = 66)	
IA	18 (27%)
IB	16 (24%)
II	9 (14%)
IIIA	3 (5%)
IIIC	14 (21%)
IVB	5 (8%)
Unknown	1 (2%)

Table 1 Demographics of patients enrolled onto the MAPPING study and eligible for analysis by meeting the primary reference standard of surgically confirmed nodal histology.

measures were significantly higher in malignant than benign lymph nodes on FDG-PET/CT (Figure 3A). Short-axis diameter, SUVmax, and STAR were all significantly elevated in malignant lymph nodes on FEC-PET/CT (Figure 3B). ADCmean and ADCmean NTR was not significantly different between benign and malignant lymph nodes in endometrial cancer (Figure 3C).

Low numbers of measured nodes in cervical cancer may limit the power of statistical analysis. No calculated or raw measures were significantly different between benign and malignant nodes on FDG-PET/CT and FEC-PET/CT,




Fig. 1 Quantitative measures of lymph nodes of each region on FDG-PET/CT (left), FEC-PET/CT (middle), and DW-MRI (right). Nodes from endometrial and cervical cancer were pooled together for this comparison.




Fig. 2 Quantitative measures of lymph nodes in cervical and endometrial cancer on DG-PET/CT (left), FEC-PET/CT (middle), and DW-MRI (right). All regions are pooled together for this comparison.

although mean measures were higher in malignant nodes (4A & 4). Interestingly, ADCmean and ADCmean NTR were significantly lower in malignant nodes in cervical cancer (4C).

Previous literature has shown elevated primary tumour SUVmax and depressed ADCmean in patients presenting with malignant lymph nodes (5). SUVmax of the primary tumour was significantly elevated on FDG-PET/CT but not on FEC-PET/CT, with no significant difference found on DW-MRI (5). No significant difference in any quantitative measure was found in primary tumours of the cervix.




Fig. 3 (A) Quantitative and calculated measures on FDG-PET/CT were significantly higher in malignant (M) than benign (B) lymph nodes in endometrial cancer. (B) SUVmax, short-axis diameter (SA), and SUVmax-to-ADCmean ratio (STAR) on FEC-PET/CT were significantly higher in malignant than benign lymph nodes in endometrial cancer. (C) No significant difference was observed between quantitative measures from DW-MRI in endometrial cancer. Nodes of all regions are pooled together for this analysis. Significance was tested between groups using Mann-Whitney U-test. ***, $p < 0.001$, **, $p < 0.01$, *, $p < 0.05$. B, benign; M, malignant; SA, short-axis diameter, LA, long-axis diameter; NTR, nodal-to-tumour ratio (SUVmax for FDG-PET/CT and FEC-PET/CT, and ADCmean for DW-MRI); STAR, SUVmax-to-ADCmean ratio; SNSA, SUVmax normalised to SA diameter.

Pearson correlations coefficients were calculated between quantitative measures in endometrial cancer (Figure 6). Short-axis diameter, long-axis diameter, and SUVmax of the lymph node were highly correlated between FDG-PET/CT and FEC-PET/CT, but did not correlate well with SUVmax of the primary tumour. SUVmax of the primary tumour correlated weakly with SUVmax of the lymph node. Furthermore, these measures were not correlated with ADCmean of the primary tumour or lymph node, except for tumour SUVmax which was inversely correlated with tumour ADCmean.




Fig. 4 (A) No significant difference in quantitative measures was found between malignant (M) and benign (B) lymph nodes on FDG-PET/CT in cervical cancer. (B) No significant difference in quantitative measures was found between malignant and benign lymph nodes on FEC-PET/CT in cervical cancer. (C) ADC_{mean} and nodal-to-tumour ratio (NTR) from DW-MRI were significantly decreased in malignant compared to benign nodes. Nodes of all regions are pooled together for this analysis. Significance was tested between groups using Mann-Whitney U-test. *, $p < 0.05$. B, benign; M, malignant; SA, short-axis diameter, LA, long-axis diameter; NTR, nodal-to-tumour ratio (SUV_{max} for FDG-PET/CT and FEC-PET/CT, and ADC_{mean} for DW-MRI); STAR, SUV_{max}-to-ADC_{mean} ratio; SNSA, SUV_{max} normalised to SA diameter.




Fig. 5 (A) SUVmax was significantly elevated in primary tumours on FDG-PET/CT in patients with malignant (M) lymph nodes compared to those with benign (B) lymph nodes in endometrial cancer. (B) No significant difference in quantitative measures of the primary tumour were found for any imaging modality in cervical cancer. **, $p < 0.01$.




Fig. 6 Pairwise Pearson correlation coefficients between raw quantitative measures from FDG-PET/CT, FEC-PET/CT, and DW-MRI in endometrial cancer. ***, $p < 0.001$, **, $p < 0.01$, *, $p < 0.05$.

3.3 Diagnostic performance of quantitative measures

Diagnostic performance of each measure, along with optimal cutpoints, in endometrial cancer is summarised in Figure 7. ROC curves are shown in Figure 8.

No raw quantitative measure clearly outperformed the others on FDG-PET/CT. SUVmax from FDG-PET/CT showed relatively high sensitivity and specificity, with the highest AUC of the quantitative measures. Short-axis diameter had the highest sensitivity of all measures from FDG-PET/CT.

The AUC of SUVmax from FEC-PET/CT was lower than that of FDG-PET/CT. Large error bars for FEC-PET/CT quantitative measures suggest low sample numbers may be restricting analysis and optimal cutpoints should be interpreted with caution. ADCmean from DW-MRI showed moderate performance with moderate sensitivity, low specificity, and moderate AUC.

Of the calculated measures, STAR had the highest AUC for FDG-PET/CT with similar performance to SUVmax.

A univariate logistic regression model trained on SUVmax performed well compared to both raw and calculated quantitative measures. The probability of malignancy based on SUVmax from FDG-PET/CT is shown in Figure 9. An optimal cut-off of () was selected to maximise the F1 score, yielding an AUC of () and relatively high specificity of ().




Fig. 7 Performance of all raw and calculated quantitative measures from FDG-PET/CT, FEC-PET/CT, and DW-MRI as well as performance of the univariate logistic regression (LR) model trained on SUVmax from FDG-PET/CT in endometrial cancer. Red crosses on box and whisker plots represent mean performance from bootstrapped data.




Fig. 8 ROC curves for quantitative measures from FDG-PET/CT (top left), FEC-PET/CT (top right), DW-MRI (bottom left), and the LR model (bottom right) in endometrial cancer. Line colour represents a quantitative measure for that imaging modality.




Fig. 9 Predicted probability (blue line) of malignancy of lymph nodes from a logistic regression model trained on nodal SUVmax from FDG-PET/CT in endometrial cancer. The red dashed line represents the optimal SUV cut-off. Points represent the SUVmax of benign (bottom) and malignant (top) lymph nodes the logistic regression model was trained on.

3.4 Interobserver agreement

Intraclass correlation coefficients (ICC) were calculated for each imaging modality. FDG-PET/CT and FEC-PET/CT had excellent interobserver agreement at 0.999 (95% CI 0.998-0.999) and 0.983 (95% CI 0.963-0.992) respectively. DW-MRI had poor interobserver agreement at 0.549 (95% CI 0.345-0.702). Interestingly, the interobserver agreement between ADCmean measurements of the primary tumour on DW-MRI was higher with an ICC of 0.797 (95% CI 0.546-0.916).

Readers were more likely to take a measurement in metastatic regions. For all imaging modalities, the frequency of both readers providing a measurement in metastatic regions was higher than the frequency of both readers providing a measurement in benign regions (FDG-PET/CT: 90% vs 67%; FEC-PET/CT: 83% vs 57%; DW-MRI: 70% vs 31%).

4 Discussion

4.1 Start here!

References

- [1] Morice, P., Leary, A., Creutzberg, C., Abu-Rustum, N. & Darai, E. Endometrial cancer. *Lancet (London, England)* **387** (10023), 1094–1108 (2016). [https://doi.org/10.1016/S0140-6736\(15\)00130-0](https://doi.org/10.1016/S0140-6736(15)00130-0) .
- [2] Siegel, R. L., Miller, K. D., Fuchs, H. E. & Jemal, A. Cancer statistics, 2022. *CA: A Cancer Journal for Clinicians* **72** (1), 7–33 (2022). <https://doi.org/10.3322/caac.21708> .
- [3] Querleu, D. *et al.* Audit of preoperative and early complications of laparoscopic lymph node dissection in 1000 gynecologic cancer patients. *American Journal of Obstetrics and Gynecology* **195** (5), 1287–1292 (2006). <https://doi.org/10.1016/j.ajog.2006.03.043> .
- [4] Achouri, A. *et al.* Complications of lymphadenectomy for gynecologic cancer. *European Journal of Surgical Oncology (EJSO)* **39** (1), 81–86 (2013). <https://doi.org/10.1016/j.ejso.2012.10.011> .
- [5] Bollineni, V. R., Ytre-Hauge, S., Bollineni-Balabay, O., Salvesen, H. B. & Haldorsen, I. S. High Diagnostic Value of 18F-FDG PET/CT in Endometrial Cancer: Systematic Review and Meta-Analysis of the Literature. *Journal of Nuclear Medicine* **57** (6), 879–885 (2016). <https://doi.org/10.2967/jnumed.115.170597> .
- [6] Sala, E., Rockall, A., Rangarajan, D. & Kubik-Huch, R. A. The role of dynamic contrast-enhanced and diffusion weighted magnetic resonance imaging in the female pelvis. *European Journal of Radiology* **76** (3),

- 367–385 (2010). <https://doi.org/10.1016/j.ejrad.2010.01.026> .
- [7] Hu, J. *et al.* Diagnostic accuracy of preoperative 18F-FDG PET or PET/CT in detecting pelvic and para-aortic lymph node metastasis in patients with endometrial cancer: A systematic review and meta-analysis. *Archives of Gynecology and Obstetrics* **300** (3), 519–529 (2019). <https://doi.org/10.1007/s00404-019-05207-8> .
 - [8] Garau, L. M. *et al.* The role of 18F-FDG PET/CT in endometrial adenocarcinoma: A review of the literature and recent advances. *Clinical and Translational Imaging* **8** (5), 357–364 (2020). <https://doi.org/10.1007/s40336-020-00385-x> .
 - [9] Antonsen, S. L. *et al.* SUVmax of 18FDG PET/CT as a predictor of high-risk endometrial cancer patients. *Gynecologic Oncology* **129** (2), 298–303 (2013). <https://doi.org/10.1016/j.ygyno.2013.01.019> .
 - [10] Husby, J. A. *et al.* Metabolic Tumor Volume on 18F-FDG PET/CT Improves Preoperative Identification of High-Risk Endometrial Carcinoma Patients. *Journal of Nuclear Medicine* **56** (8), 1191–1198 (2015). <https://doi.org/10.2967/jnumed.115.159913> .
 - [11] Kim, H. J., Cho, A., Yun, M., Kim, Y. T. & Kang, W. J. Comparison of FDG PET/CT and MRI in lymph node staging of endometrial cancer. *Annals of Nuclear Medicine* **30** (2), 104–113 (2016). <https://doi.org/10.1007/s12149-015-1037-8> .
 - [12] Rechichi, G. *et al.* Myometrial invasion in endometrial cancer: Diagnostic performance of diffusion-weighted MR imaging at 1.5-T. *European Radiology* **20** (3), 754–762 (2010). <https://doi.org/10.1007/s00330-009-1597-x> .
 - [13] Beddy, P. *et al.* Evaluation of Depth of Myometrial Invasion and Overall Staging in Endometrial Cancer: Comparison of Diffusion-weighted and Dynamic Contrast-enhanced MR Imaging. *Radiology* **262** (2), 530–537 (2012). <https://doi.org/10.1148/radiol.11110984> .
 - [14] Koplay, M. *et al.* Diagnostic efficacy of diffusion-weighted MRI for pre-operative assessment of myometrial and cervical invasion and pelvic lymph node metastasis in endometrial carcinoma. *Journal of Medical Imaging and Radiation Oncology* **58** (5), 538–546 (2014). <https://doi.org/10.1111/1754-9485.12209> .
 - [15] Nakai, G. *et al.* Detection and Evaluation of Pelvic Lymph Nodes in Patients With Gynecologic Malignancies Using Body Diffusion-Weighted Magnetic Resonance Imaging. *Journal of Computer Assisted Tomography* **32** (5), 764–768 (2008). <https://doi.org/10.1097/RCT.0b013e318153fd43>

- [16] Roy, C., Bierry, G., Matau, A., Bazille, G. & Pasquali, R. Value of diffusion-weighted imaging to detect small malignant pelvic lymph nodes at 3 T. *European Radiology* **20** (8), 1803–1811 (2010). <https://doi.org/10.1007/s00330-010-1736-4> .
- [17] Chen, Y. B., Liao, J., Xie, R., Chen, G. L. & Chen, G. Discrimination of metastatic from hyperplastic pelvic lymph nodes in patients with cervical cancer by diffusion-weighted magnetic resonance imaging. *Abdominal Imaging* **36** (1), 102–109 (2011). <https://doi.org/10.1007/s00261-009-9590-z> .
- [18] Rechichi, G. *et al.* ADC maps in the prediction of pelvic lymph nodal metastatic regions in endometrial cancer. *European Radiology* **23** (1), 65–74 (2013). <https://doi.org/10.1007/s00330-012-2575-2> .
- [19] Trousil, S. *et al.* Alterations of Choline Phospholipid Metabolism in Endometrial Cancer Are Caused by Choline Kinase Alpha Overexpression and a Hyperactivated Deacylation Pathway. *Cancer Research* **74** (23), 6867–6877 (2014). <https://doi.org/10.1158/0008-5472.CAN-13-2409> .
- [20] Contractor, K. *et al.* Use of [11C]Choline PET-CT as a Noninvasive Method for Detecting Pelvic Lymph Node Status from Prostate Cancer and Relationship with Choline Kinase Expression. *Clinical Cancer Research* **17** (24), 7673–7683 (2011). <https://doi.org/10.1158/1078-0432.CCR-11-2048> .
- [21] Torizuka, T. *et al.* Imaging of Gynecologic Tumors: Comparison of 11C-Choline PET with 18F-FDG PET. *Journal of Nuclear Medicine* **44** (7), 1051–1056 (2003) .
- [22] Sofue, K. *et al.* Role of carbon-11 choline PET/CT in the management of uterine carcinoma: Initial experience. *Annals of Nuclear Medicine* **23** (3), 235–243 (2009). <https://doi.org/10.1007/s12149-009-0230-z> .
- [23] Rockall, A. G. *et al.* Diagnostic Accuracy of FEC-PET/CT, FDG-PET/CT, and Diffusion-Weighted MRI in Detection of Nodal Metastases in Surgically Treated Endometrial and Cervical Carcinoma. *Clinical Cancer Research* **27** (23), 6457–6466 (2021). <https://doi.org/10.1158/1078-0432.CCR-21-1834> .