

The Medium

An Attempted Description of a Pre-Geometric Substrate — Foundation Definition

Bradley K. DePuy, Independent Researcher

ORCID: 0009-0001-0328-2299

2026

1 Definition

1.1 The Substrate

The Medium is the substrate — denoted M . It has no beginning and no end, no geometry, no time, no parts, and no structure prior to distinction. M is not a space or a set of points. It is what is.

1.2 The Primitive: ι

The sole primitive is ι : a distinction within M . Not an event — event implies sequence, which implies time. ι is prior to time. A distinction is the substrate differentiating within itself.

Distinction is self-grounding. Any attempt to explain what causes it already uses it. Causation, before, after — these are all distinctions. ι requires no prior cause: it is the minimum condition for anything to be conceivable at all.

A single ι does not carry character or polarity. It produces a split: two complementary aspects $\{A, \bar{A}\}$ that are not each other. Neither A nor $\bar{A}$ is intrinsically positive or negative. Polarity is not a property of a single ι .

Let D denote the collection of all ι distinctions within M .

1.3 Character from Comparison

When two distinctions $i, j \in D$ co-exist within M , a comparison is unavoidable: do they distinguish in the same way, or in opposite ways? This comparison is binary and symmetric:

$$c : D \times D \rightarrow \{\text{same, different}\} \tag{1}$$

$c(i, i) = \text{same}$. $c(i, j) = c(j, i)$. What earlier formulations labeled $+1$ or -1 is the outcome of this comparison. Character and polarity are the first emergent properties of the substrate, not primitives.

1.4 The Relational Web

Co-existing distinctions in D necessarily relate. When D contains more than one distinction, comparisons arise throughout, generating a relational graph G :

- Nodes: ι distinctions in D
- Edges: relations between co-existing distinctions

The negotiation field N_{ij} measures the coherence of the comparison between distinctions i and j across the entire relational structure — not the comparison value itself, but the degree to which that comparison is consistent across all paths through G . N is real-valued and continuous: $N_{ij} > 0$ (different-comparing, stable); $N_{ij} < 0$ (same-comparing, unstable); $N_{ij} = 0$ (uncorrelated).

The Laplacian L of G encodes the relational geometry. In the large-scale continuum limit L converges to the Laplace-Beltrami operator on a Riemannian manifold — geometry emerging from the relational structure of distinctions, not assumed.

1.5 The Adjacency Constraint

Two distinctions comparing as same carry no new relational information: their comparison is trivially resolved. Negotiations between same-comparing distinctions contribute nothing to stable structure. This is not a topological prohibition on edges — it is a statement about information content. Productive negotiation requires $c(i, j) = \text{different}$. Stable configurations require all local comparisons to be productive.

1.6 A Note on Time

The dynamics below use $\partial N / \partial t$. Time appears here as a background parameter retained for calculability. In the substrate as described, sequence does not exist prior to distinction — time is not a primitive but an emergent ordering of the relational structure of D . The derivation of temporal sequence from substrate dynamics is an open problem.

2 Scale Inputs

This description has exactly two inputs:

Input	Value	Role
Iota scale	$l_i = 1.616 \times 10^{-35} \text{ m}$	Substrate grain
Hubble scale	$R_H = c/H_0$ (current epoch)	Longest negotiation mode

In Planck units: $G = 1$, $c = 1$, $\hbar = 1/(2\pi)$. What SI physics calls $\hbar$ is the measurement of one projected ι event in the observer's unit system — a conversion artifact, not a primitive constant.

The negotiation energy quantum is estimated from the geometric mean of the two scale energies:

$$\varepsilon_{\text{est}} = (E_{\text{Planck}}^2 \cdot E_{\text{Hubble}})^{1/3} \approx 152.6 \text{ MeV} \quad (2)$$

With a $(2\pi)^{1/3}$ dispersion factor this reaches 281.5 MeV. The observed value $\varepsilon = m_{\text{proton}}/3 = 312.8 \text{ MeV}$ is a definition from observation; the $\sim 10\%$

3 Dynamics

The negotiation field N_{ij} encodes the relational coherence between adjacent ι distinctions i and j . Lagrangian in Planck units:

$$\mathcal{L} = \frac{1}{2} \left(\frac{\partial N}{\partial t} \right)^2 - \frac{1}{2} N^\top L N \quad (3)$$

Hamiltonian (via Legendre transform, $\Pi = \partial N / \partial t$):

$$\hat{H} = \frac{1}{2} \Pi^2 + \frac{1}{2} N^\top L N \quad (4)$$

Energy eigenvalues: $E_k = \varepsilon \lambda_k$, where λ_k are eigenvalues of the graph Laplacian L .

Stable particles satisfy the closure condition $L \cdot N = 0$. For a complete graph on n nodes with mode number k , energy eigenvalues follow:

$$E(n, k) = \varepsilon \cdot k \cdot n \quad (5)$$

The minimum closed graph is the triangle of three ι distinctions ($\lambda = 3$, $n = 3$):

$$m_{\text{proton}} = 3\varepsilon = 938.3 \text{ MeV} \quad (6)$$

Higher topologies follow the same formula: a two-node graph corresponds to the rho meson ($n = 2$, predicted 625.6 MeV); a four-node graph corresponds to the tetraquark ($n = 4$, predicted 1251 MeV). The triangle is the ground state — the minimum topology that achieves closure.

4 Derived Results

From two inputs and the single correction factor $\gamma \approx 0.2375$:

Quantity	Derived	Measured	Agreement
m_{proton}	$3\varepsilon = 938.3 \text{ MeV}$	938.3 MeV	Exact (definition of ε)
G	$H_0^2/4\pi\rho_\Lambda$	$6.674 \times 10^{-11} \text{ SI}$	94–97%
$\hbar$	$m_p c k(\gamma) r_p/6\pi$	$1.055 \times 10^{-34} \text{ J}\cdot\text{s}$	99.1%
H_0	68.6 km/s/Mpc	67.4–73.0 (tension range)	Within tension
Λ	$\approx 10^{-122} \text{ Planck}$	$\approx 10^{-122} \text{ Planck}$	Same order

The ratio $\rho_{\text{Planck}}/\rho_\Lambda \approx 10^{123}$ is the depth of the Medium — the dynamic range from iota grain to cosmic horizon — not a fine-tuning problem.

Analytic prediction: $\Omega_\Lambda = 2/3 \approx 0.667$ (observed: 0.685; discrepancy 2.7\

All three residual gaps (in G , m_{proton} , and $\hbar$) are closed by a single Barbero-Immirzi calculation on the negotiation graph. This is the primary open calculation in this description.

5 License and Attribution

CC BY 4.0. DOI: 10.5281/zenodo.18927154 ORCID: 0009-0001-0328-2299 github.com/bkdepuy/The-Medium

6 Document Series and Cross-References

This is Document 1 of 5 in The Medium series.

Doc	File	Content
0	medium_doc0_introduction.org	Personal introduction; the question behind the description
1	medium_doc1_definition.org	<i>This paper</i>
2	medium_doc2_unified.org	Negotiation Modes, Particle Structure, Proton Mass
3	medium_doc3_lagrangian.org	Lagrangian of the Negotiation Field; QFT connection; SU(3)
4	medium_doc4_hamiltonian.org	Hamiltonian, energy spectrum, canonical quantization, ADM
5	medium_doc5_G_H0.org	Deriving G and H_0 from substrate primitives