

# 1 Sum from 1 to n

We want to prove that  $1 + 2 + \dots + n$  can be calculated with  $\frac{n(n+1)}{2}$

## 1.1 Definitions

We will define a function to let us talk about the sum of numbers from 1 to  $n$ .  
Let:

$$F(n) = 1 + 2 + \dots + n \quad (1)$$

We will define a predicate to let us talk about the relationship between  $F(n)$  and the shortcut calculation. Let:

$$P(n) : F(n) = \frac{n(n+1)}{2} \quad (2)$$

Note that  $P(n)$  evaluates to a boolean. It can be true or false for any particular  $n$ . It is true for a particular value of  $n$  if  $F(n)$  does in fact equal  $\frac{n(n+1)}{2}$  and it is false if these two things are not equal.

## 1.2 Goal

Our goal is to prove that  $P(n)$  holds (is true) for all values of  $n$  greater than 0.  
Prove:

$$\forall n \in N : P(n) \quad (3)$$

## 1.3 Proof by induction

### 1.3.1 Base case

To show our base case  $P(1)$  is true, we will state the base case, then show that the left side does in fact equal the right side. Prove:

$$P(1) : F(1) = \frac{1(1+1)}{2} \quad (4)$$

$$F(1) = 1$$

$$\frac{1(1+1)}{2} = \frac{2}{2} = 1$$

### 1.3.2 Inductive step

We will prove that **if**  $P(k)$  holds (is true) for some  $k \in N$ , **then**  $P(k+1)$  is also true. Prove:

$$P(k) \implies P(k+1), \forall k \in N \quad (5)$$

We start with the *inductive hypothesis*, we assume for the time that  $P(k)$  holds.  
Assume:

$$P(k) : F(k) = \frac{k(k+1)}{2} \quad (6)$$

Now, assuming that  $P(k)$  is true, prove:

$$P(k+1) : F(k+1) = \frac{(k+1)((k+1)+1)}{2} \quad (7)$$

By definition:

$$F(k+1) = 1 + 2 + \dots + k + (k+1)$$

which is by definition:

$$F(k+1) = F(k) + (k+1)$$

which by our inductive hypothesis is:

$$F(k+1) = \frac{k(k+1)}{2} + (k+1)$$

simplifying is:

$$F(k+1) = (k+1)\left(\frac{k}{2} + 1\right)$$

which is equivalent to:

$$F(k+1) = (k+1)\left(\frac{k}{2} + \frac{2}{2}\right)$$

which simplifies to:

$$F(k+1) = \frac{(k+1)(k+2)}{2}$$

which is clearly:

$$F(k+1) = \frac{(k+1)((k+1)+1)}{2}$$

And so we have proved  $P(k+1)$  (7) by showing that the left side is equal to the right side (assuming that  $P(k)$  is true).

## 1.4 Conclusion

We have proved that  $P(n)$  holds for a base case of  $P(1)$  and that for all  $k \in N$ ,  $P(k)$  being true implies that  $P(k+1)$  is also true. Therefore  $P(n)$  holds for all  $n > 0$  (all natural numbers).

$$P(1) : F(1) = \frac{1(1+1)}{2}$$

$$P(k) \implies P(k+1), \forall k \in N$$

$$\therefore P(n), \forall n \in N$$

## 2 Making postage with 3 and 5 cent stamps

We want to prove that all postage amounts greater than 7 cents can be made with combinations of 3 and 5 cent stamps

## 2.1 Definitions

We will define a predicate to talk about whether a particular number can be represented as a summation of a non-negative multiple of 3 and a non-negative multiple of 5.

$$P(n) : n = 3a + 5b \mid a, b \in \mathbb{Z}_{\geq 0} \quad (8)$$

Note that  $P(n)$  may be true or false for any given number  $n$ . For example,  $P(2)$  is false, as 2 cents of postage cannot be made with 3 and 5 cent stamps. However,  $P(11)$  is true, because 11 cents of postage can be made with a 5 cent stamp and two 3 cent stamps.

## 2.2 Goal

Our goal is to prove that  $P(n)$  holds for all values of  $n$  greater than 7.

$$\forall n \in \mathbb{Z}_{>7} : P(n) \quad (9)$$

## 2.3 Proof by induction

### 2.3.1 Base case

To show that the base case  $P(8)$  is true, we will state the base case, then show that we can find suitable non-negative integers  $a$  and  $b$ . Prove:

$$P(8) : 8 = 3a + 5b \quad (10)$$

$$a, b = 1$$

### 2.3.2 Inductive step

We will prove that **if**  $P(k)$  holds (is true) for some  $k \in \mathbb{Z}_{>7}$ , **then**  $P(k + 1)$  is also true. Prove:

$$P(k) \implies P(k + 1), \forall k \in \mathbb{Z}_{>7} \quad (11)$$

We start with the *inductive hypothesis*, we assume for the time that  $P(k)$  holds. Assume:

$$P(k) : k = 3a + 5b \mid a, b \in \mathbb{Z}_{\geq 0} \quad (12)$$

Now, assuming that  $P(k)$  is true, prove:

$$P(k + 1) : k + 1 = 3c + 5d \mid c, d \in \mathbb{Z}_{\geq 0} \quad (13)$$

If  $b > 0$ , then  $d = b - 1$  and  $c = a + 2$  results in:

$$3c + 5d = 3(a + 2) + 5(b - 1) = 3a + 5b + 1 = k + 1$$

By our inductive hypothesis, we assumed that  $a$  and  $b$  were non-negative integers, and so  $c$  and  $d$  will be too.

If  $b = 0$ , then  $k$  must be a multiple of 3. The smallest multiple of 3 in  $\mathbb{Z}_{>7}$  is 9, which is  $3 * 3$ . All other multiples of 3 in  $\mathbb{Z}_{>7}$  will have a greater or equal

number of 3 in their all-3 representation ( $b = 0$ ). Therefore if  $b = 0$ , then  $a \geq 3$ . So  $d = b + 2$  and  $c = a - 3$  results in:

$$3c + 5d = 3(a - 3) + 5(b + 2) = 3a + 5b + 1 = k + 1$$

By our inductive hypothesis and our reasoning that  $a$  must be greater than or equal to 3 in the case where  $b = 0$ , then  $c$  and  $d$  will be non-negative integers too.

And so we have proved  $P(k + 1)$  (13) by assuming  $P(k)$  (12) was true. If  $P(k)$ , then  $P(k + 1)$  (11).

## 2.4 Conclusion

We have proved that  $P(n)$  holds for a base case of  $P(8)$  and that for all  $k \in \mathbb{Z}_{>7}$ ,  $P(k)$  being true implies that  $P(k + 1)$  is also true. Therefore  $P(n)$  holds for all  $n > 7$ .

$$P(8) : 8 = 3(1) + 5(1)$$

$$P(k) \implies P(k + 1), \forall k \in \mathbb{Z}_{>7}$$

$$\therefore P(n), \forall n \in \mathbb{Z}_{>7}$$

## 3 Proof with inequality

We want to prove that for any integer  $n$  greater than 6,  $n!$  is greater than  $3^n$ .

$$n! > 3^n \mid n > 6$$

### 3.1 Definitions

We will define a predicate to let us talk about an inequality relationship between  $n!$  and  $3^n$ . Let:

$$P(n) : n! > 3^n \tag{14}$$

Note that  $P(n)$  may be true or false for any particular choice of  $n$ . For example,  $P(2)$  is false because  $2!$  is not greater than  $3^2$ . But  $P(7)$  is true because  $7!$  (5040) is greater than  $3^7$  (2187).

### 3.2 Goal

Our goal is to prove that  $P(n)$  holds for all values of  $n$  greater than 6.

$$\forall n \in \mathbb{Z}_{>6} : P(n) \tag{15}$$

### 3.3 Proof by induction

#### 3.3.1 Base case

To show that the base case  $P(7)$  is true, we will state the base case, then show that the inequality is true. Prove:

$$\begin{aligned} P(7) : 7! &> 3^7 \\ 7! &= 5040 > 2187 = 3^7 \end{aligned} \tag{16}$$

#### 3.3.2 Inductive step

We will prove that **if**  $P(k)$  holds (is true) for some  $k \in \mathbb{Z}_{>6}$ , **then**  $P(k+1)$  is also true. Prove:

$$P(k) \implies P(k+1), \forall k \in \mathbb{Z}_{>6} \tag{17}$$

We start with the *inductive hypothesis*, we assume for the time that  $P(k)$  holds. Assume:

$$P(k) : k! > 3^k \mid k \in \mathbb{Z}_{>6} \tag{18}$$

Now, assuming that  $P(k)$  is true, prove:

$$P(k+1) : (k+1)! > 3^{k+1} \tag{19}$$

Starting with the left side:

$$(k+1)! = (k+1)k!$$

Using our inductive hypothesis:

$$\begin{aligned} (k+1)! &> (k+1)3^k \\ (k+1)! &> \frac{1}{1}(k+1)3^k \\ (k+1)! &> \frac{3}{3}(k+1)3^k \\ (k+1)! &> \frac{k+1}{3}3 * 3^k \\ (k+1)! &> \frac{k+1}{3}3^1 * 3^k \\ (k+1)! &> \frac{k+1}{3}3^{k+1} \end{aligned}$$

And if  $\frac{k+1}{3}$  is greater than 1:

$$(k+1)! > \frac{k+1}{3}3^{k+1} > 3^{k+1}$$

therefore:

$$(k+1)! > 3^{k+1}$$

Since  $k \in \mathbb{Z}_{>6}$ , we know that  $\frac{k+1}{3}$  will always be greater than  $\frac{6}{3}$  which is greater than 1. And so we have proved  $P(k+1)$  (19) by assuming  $P(k)$  (18) was true. If  $P(k)$ , then  $P(k+1)$  (17).

### 3.4 Conclusion

We have proved that  $P(n)$  holds for a base case of  $P(7)$  and that for all  $k \in \mathbb{Z}_{>6}$ ,  $P(k)$  being true implies that  $P(k+1)$  is also true. Therefore  $P(n)$  holds for all  $n > 6$ .

$$\begin{aligned} P(7) : 7! &> 3^7 \\ P(k) &\implies P(k+1), \forall k \in \mathbb{Z}_{>6} \\ \therefore P(n), \forall n &\in \mathbb{Z}_{>6} \end{aligned}$$

## 4 Another summation

We want to prove that  $1 + 4 + 7 + \dots + (3n - 2)$  can be calculated with  $\frac{n(3n-1)}{2}$