

---

# Chapter 1

## The Standard Model of Particle Physics

Particle physics, or high-energy physics, is the study of the most fundamental constituents of matter and their interactions. The best current description of these interactions is known as The Standard Model of Particle Physics (SM); a group of theories that cover all currently known particles and their interactions. The SM was developed through-out the latter half of the 20th century and has seen tremendous success in predicting the behaviour of our universe at the most fundamental level. The SM has stood the test of time and rigorous examination by numerous experiments. Additionally many of its parameters have been measured with tremendous precision e.g. the electron magnetic moment  $g$  is known to  $10^{-13}$  [1]. The last piece to be confirmed was the existence of the Higgs boson, which in turn points to the existence of the so-called Higgs field. Evidence of the elusive Higgs were observed by the ATLAS and CMS experiments at CERN [2,3]. Despite its tremendous success, the SM cannot account and explain for all observed phenomenon in the universe. Firstly, the theory requires many of its parameters to be measured empirically. The theory does not a priori provide a value for these parameters such as the number of particle generations. Additionally the theory does not describe the most familiar of the forces, gravity. Furthermore, the SM does not provide a candidate for dark matter or dark energy, which according to recent measurements accounts for more than 90% of the total energy density in the universe [4]. The asymmetry between

22 matter and antimatter is also not fully explained by the SM. rite something about  
 23 radiative corrections to higgs mass and the hierarchy problem As such there is a strong  
 24 focus on developing theories which go beyond the standard model (BSM) to provide an  
 25 answer to these open questions. The discussion in this chapter is largely based on [5]  
 26 and [6].

27 The SM describes the nature of the interactions of the fundamental constituents  
 28 of our universe in terms of the three different fundamental forces: strong, weak and  
 29 electromagnetic, each described by a specific theory. As mentioned before, the most  
 30 familiar of the forces, gravity, is not described by the SM. The SM classifies particles  
 31 into several categories depending on their properties and allowed interactions. Particles  
 32 which have a half-integer spins (e.g.  $S = \frac{1}{2}, \frac{3}{2}, \dots$ ) are known as *fermions*, these are the  
 33 basic constituents of matter. Particles with integer spins (e.g.  $S = 0, 1, \dots$ ) are known as  
 34 *bosons*, these mediate interactions between fermions and other bosons.

35 Fermions can be divided into two subgroups: quarks, which can interact via the  
 36 strong, weak and electromagnetic forces and leptons which can only interact by the  
 37 weak and electromagnetic forces. There are six known leptons: electron  $e$ , muon  $\mu$  and  
 38 tau  $\tau$ , which all have electric charge<sup>1</sup>  $Q = 1$ , and the corresponding electrically neutral  
 39 neutrino  $\nu_e$ ,  $\nu_\mu$  and  $\nu_\tau$ . Analogously, six quark *flavours* are known:  $u$ ,  $c$  and  $t$ , with  
 40 electric charge  $Q = +2/3$  and  $d$ ,  $s$  and  $b$ , with electric charge  $Q = -1/3$ .

41 Quarks and leptons are divided into three generations, which differ only by the  
 42 mass and flavour of their constituent fermions, each generation being heavier than the  
 43 previous. A summary of all elementary particles described by the SM can be found in  
 44 Table 1.1.

45 For every matter fermion ( $f$ ) there is an equivalent antimatter partner ( $\bar{f}$ ) which  
 46 possesses the same characteristics as its matter companion but is opposite in electric  
 47 charge. Thus 12 matter particles are combined with 12 antimatter partners for a total  
 48 of 24 elementary particles which form all visible matter in the universe.

49 The interaction between fermions occur via the exchange of spin one particles known  
 50 as bosons. Each force is mediated by one or more bosons (Table 1.2). The strong force  
 51 is mediated by a set of massless bosons known as the gluons. The weak force is mediated

---

<sup>1</sup>The electric charge is always stated in units of elementary charge  $e$

by a neutral massive boson known as the  $Z$  boson and a pair of charged massive bosons known as the  $W$  bosons. Finally, the electromagnetic force is mediated by a massless boson known as the photon. Note that each boson has an antimatter partner however some are indistinguishable from their matter partner. A summary of their properties is shown in Table 1.1.

Each fermion has a set of so-called quantum numbers which dictate the type of interactions that can occur. For example each lepton has a lepton number associated with it, electrons have an electron lepton number ( $L_e$ ) of  $+1$ , while the positron has  $L_e = -1$ . Muons and taus have their own respective lepton number ( $L_\mu$  and  $L_\tau$ ). Each neutrino has lepton number  $L_f = 1$  and their anti-matter counterpart have  $L_f = -1$ . Each of these lepton numbers is approximately conserved separately across interaction vertices. The conservation is only approximate due to the non-zero mass of neutrinos. Another example of a quantum number is baryon number ( $B$ ), each quark has  $B = \frac{1}{3}$  and anti-quarks have  $B = -\frac{1}{3}$ .

## 1.1 Quantum Electrodynamics

The interaction of particles via the electromagnetic force is described by Quantum Electrodynamics or QED. These interactions are mediated by the massless neutral boson known as the photon and the strength of the interaction is characterized by the fine-structure constant  $\alpha$ . All electrically charged fermions are allowed to interact, since the photon itself is not charged, no self-interaction is allowed within QED. Figure 1.1 shows the single vertex described by QED, where two fermions interact via a photon. Note that the electric charge is conserved across the vertex, so for example  $\gamma \rightarrow e^+e^+$  is not allowed within QED.

By combining different forms of this vertex one can build every possible QED interaction. The interaction  $e^+e^- \rightarrow e^+e^-$  is known as Bhabha scattering. Two leading order (LO)<sup>2</sup> diagrams contribute to this interaction, annihilation (Figure 1.2a) and scattering (Figure 1.2b).

---

<sup>2</sup>The simplest diagram with the least vertices

		Fermions ( $s = \frac{1}{2}$ )			Bosons ( $s = 0$ )	Higgs ( $s = 1$ )
		I	II	III		
Quarks		$+\frac{2}{3}$ 2.3 MeV $u$ Up	$+\frac{2}{3}$ 1.275 GeV $c$ Charm	$+\frac{2}{3}$ 173.07 GeV $t$ Top	0 0 MeV $\gamma$ Photon (EM)	0 126.07 GeV $H^0$ Higgs boson
		$-\frac{1}{3}$ 4.8 MeV $d$ Down	$-\frac{1}{3}$ 95 MeV $s$ Strange	$-\frac{1}{3}$ 4.18 GeV $b$ Bottom	$\pm 1$ 80.4 GeV $W^\pm$ W boson (Weak)	
Leptons		$-1$ 0.511 MeV $e$ Electron	$-1$ 105.7 MeV $\mu$ Muon	$-1$ 1.777 GeV $\tau$ Tau	0 91.2 GeV $Z$ Z boson (Weak)	
		0 < 2.2 eV $\nu_e$ Electron Neutrino	0 < 0.17 MeV $\nu_\mu$ Muon Neutrino	0 15.5 MeV $\nu_\tau$ Tau Neutrino	0 0 MeV $g$ Gluon (Strong)	q mass <b>symbol</b> name (force)

Table 1.1: A summary of all elementary particles described by the SM [7]. Note the various groupings and divisions including by spin, generation and particle type. For each particle the charge ( $q$ ), mass and name are shown as per the legend on the bottom-right.

Name	Relative Strength	Boson
Strong	$10^{38}$	Gluons
Electromagnetic	$10^{36}$	Photon
Weak	$10^{25}$	$W^\pm$ and $Z^0$
Gravity	1	Graviton*

Table 1.2: A summary of the four fundamental forces ordered by relative strength. These are approximate relative strengths for the purpose of demonstrating the hierarchy of forces as a function of their strength. A more accurate determination of the interaction strength depends on the details of the interaction itself. Note however the order-of-magnitude differences in the relative strengths of these forces. Note that the graviton is the theoretical boson responsible for mediating gravitational interactions, it is not part of the SM.






Figure 1.1: The interaction vertex described by QED. One can obtain all possible vertex shapes by rotating this basic vertex and assigning the appropriate electric charge and making sure to conserve lepton number across the vertex.



(a) Electron-Positron pair annihilation mediated by a photon.



(b) Electron-Positron pair scattering via the emission of a photon.

Figure 1.2: LO feynman diagrams of the process  $e^+e^- \rightarrow e^+e^-$  allowed in QED. Additional vertices can be added to produce higher-order diagrams of the same process.

## 79 1.2 Quantum Chromodynamics

80 Interactions via the strong force are described in the theory of Quantum Chromodynam-  
 81 ics or QCD. These interactions are mediated by a set of massless neutral bosons known  
 82 as gluons. QCD introduces the concept of colour, which similarly to electrical charge,  
 83 determines the possible interactions that can occur via the strong force. Colour can take  
 84 three states, red (antired), blue (antiblu), green (antigreen):

$$r = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad g = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad b = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (1.1)$$

$$\bar{r} = \begin{pmatrix} 1 & 0 & 0 \end{pmatrix}, \quad \bar{g} = \begin{pmatrix} 0 & 1 & 0 \end{pmatrix}, \quad \bar{b} = \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \quad (1.2)$$

Both quarks and gluons possess colour and as a result gluons, unlike photons, can self-interact in a three gluon vertex (Figure 1.3b) or a four gluon vertex (Figure 1.3c). As with electrical charge, colour-charge must also be conserved. Thus in the scattering process  $q \rightarrow q + g$  shown in Figure 1.3a the flavour of the quark may not change but the colour-charge does and the gluon carries away the difference in colour. Thus each gluon has a colour-anticolour charge associated with it. Naively one would expect nine different types of gluon that participate in interaction, owing to the nine possible combinations of colour and anticolour, however the SU(3) symmetry on which QCD is based results in a colour octet:

$$\begin{array}{ll} (r\bar{b} + b\bar{r})/\sqrt{2} & -i(r\bar{g} - g\bar{r})/\sqrt{2} \\ -i(r\bar{b} - b\bar{r})/\sqrt{2} & (b\bar{g} + g\bar{b})/\sqrt{2} \\ (r\bar{r} + b\bar{b})/\sqrt{2} & -i(b\bar{g} - g\bar{b})/\sqrt{2} \\ (r\bar{g} + g\bar{r})/\sqrt{2} & (r\bar{r} + b\bar{b} - 2g\bar{g})/\sqrt{6} \end{array}$$

85 and a “colour singlet”:

$$(r\bar{r} + g\bar{g} + b\bar{b})/\sqrt{3} \quad (1.3)$$

86 which is overall colourless.




Figure 1.3: Diagrams of the fundamental interaction vertices described by quantum chromodynamics. Shown are (a) Gluon emission from a quark, (b) Gluon emission from a gluon and finally (c) the four-gluon vertex.

87 There are then eight different gluons that can participate in QCD interactions each  
 88 with a different colour-charge combination. Additionally there is a ninth combination  
 89 which is overall colorless so it cannot take part in interactions.

90 In an analogous fashion to screening which occurs with electric charges, quark-  
 91 antiquark pairs act like dipoles which screen the true colour charge of the central quark.  
 92 However since gluons also carry colour, they cause the opposite effect (anti-screening) to  
 93 amplify and change the observed colour of the quark. Which effect wins out depends on  
 94 the number of colours in the theory and the number of quark flavours. As it is with three  
 95 colour states and six different quark flavours, anti-screening is the overall dominant ef-  
 96 fect. As a result the colour potential decreases with distance and quarks experience very  
 97 little potential when very near to each other. This effect is known as asymptotic freedom  
 98 and results in quarks only existing within colorless bound states known as *hadrons*.

99 Hadrons can be divided into two categories: *mesons*, which contain a quark and an  
 100 antiquark ( $q\bar{q}$ ); and *baryons* which are made of three quarks (or antiquarks) each with  
 101 a different (anti)colour-charge to result in a colourless composite particle. Common  
 102 examples of baryons are protons ( $uud$ ) and neutrons ( $udd$ ) which are the building blocks  
 103 of atomic nuclei. While  $\pi^0$  ( $u\bar{u}/d\bar{d}$ ) is a commonly produced meson in hadron colliders.  
 104 Note that due to the quark configuration, baryons have baryon number  $B = +1$  while  
 105 mesons have  $B = 0$ .




Figure 1.4: The neutral current and charged current vertices allowed via the weak force. Where  $f$  can be an  $e$ ,  $\mu$  or  $\tau$  and  $\nu_\ell$  is the corresponding lepton neutrino of the same flavour. One can obtain all possible interaction vertices by rotating these basic vertices and assigning the appropriate electric charge and making sure to conserve lepton flavour across the vertex.




Figure 1.5: Neutral current weak scattering vertex

### 1.3 Weak Interactions

The final type of interaction involves the so-called weak force. The weak force is responsible for  $\beta^-$  decay ( $n \rightarrow p + e^- + \bar{\nu}_e$ ) and  $\beta^+$  decay. Interactions via the weak force are mediated by a single neutral massive boson and two charged massive bosons. Since the bosons responsible for weak interactions are massive, the range of interaction is very short, unlike electromagnetic interactions via a massless photon.

All fermions can take part in interactions via the weak force. Let us consider weak interactions involving only leptons. The weak neutral vertex is very similar to the basic vertex seen in QED (1.1). A valid interaction via the weak force is then formed by combining these simple vertices (Figure 1.4) while taking care to conserve electric charge and lepton flavour. An example of a leptonic weak interaction is muon decay ( $\mu \rightarrow \nu_\mu e^- \bar{\nu}_e$ ) shown in Figure 1.5.

Let us consider weak interactions involving quarks. The neutral vertex is similar to that of the leptonic version, a quark can emit a  $Z$  boson or a  $Z$  can decay to a quark-antiquark pair. The charged current then changes the flavour of an up-type quark into a



121 down-type quark (or vice-versa) with a  $W$  boson of the appropriate charge (Figure 1.4c).  
 122 Weak interactions can also change the flavour of a quark across generations. A well-  
 123 known example of such an interaction is Kaon decay ( $K^+ \rightarrow \mu^+ \nu_\mu$ ). In order to account  
 124 for this interaction and preserve the universality of weak interactions, Nicola Cabibbo  
 125 postulated [8] that the states that couple to the charged current are really a mixture of  
 126 'rotated' quark states:

$$\begin{pmatrix} u \\ d' \end{pmatrix} = \begin{pmatrix} c \\ s' \end{pmatrix} \quad (1.4)$$

127 where

$$d' = d \cos \theta_c + s \sin \theta_c \quad (1.5a)$$

128

$$s' = -d \sin \theta_c + s \cos \theta_c \quad (1.5b)$$

129 This introduces an arbitrary parameter into the theory known as the quark mixing  
 130 angle or the Cabibbo angle  $\theta_c$ . The introduction of quark mixing has the effect of  
 131 attenuating the interaction strength at vertices involving multiple quark generations.  
 132 Interactions which cross one generation are said to be Cabibbo Suppressed while those  
 133 that cross two generations are Doubly Cabibbo suppressed.

134 Taking into account the three quark generations, quark mixing can be expressed  
 135 in matrix notation as shown in Equation 1.6. This unitary matrix is known as the  
 136 Cabibbo-Kobayashi-Maskawa Matrix (CKM Matrix) after Cabibbo which initially pos-  
 137 tulated quark mixing and Makoto Kobayashi and Toshihide Maskawa who later added  
 138 an additional generation, containing the top and bottom quarks, to the matrix [9].

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = V_{CKM} \begin{pmatrix} d \\ s \\ b \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} \quad (1.6)$$

139 Several parameterizations of the CKM matrix exist, the Chau-Keung parametriza-  
 140 tion [10] uses angles  $\theta_{12}$ ,  $\theta_{23}$ ,  $\theta_{13}$  and a phase  $\delta_{13}$ :

$$V_{CKM} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}\exp(-i\delta) \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}\exp(i\delta) & c_{12}c_{23} - s_{12}s_{23}s_{13}\exp(i\delta) & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}\exp(i\delta) & -c_{12}s_{23} - s_{12}c_{23}s_{13}\exp(i\delta) & c_{23}c_{13} \end{pmatrix} \quad (1.7)$$

141 where  $c_{ij} = \cos \theta_{ij}$  and  $s_{ij} = \sin \theta_{ij}$  for  $i=1,2,3$ . This parametrization has the advantage  
 142 that each angle  $\theta_{ij}$  relates to a specific transition from one generation to the other. If  
 143  $\theta_{13} = \theta_{23} = 0$  the third generation is not coupled to the other two and the matrix reduces  
 144 to the original matrix postulated by Cabibbo. Note that  $\theta_{12}$  is the Cabibbo angle,  $\theta_c$ ,  
 145 described earlier.

146 Another parameterization due to Wolfenstein [11] expresses all elements in terms of  
 147 the Cabibbo angle by defining  $\lambda \equiv s_{12} = \sin \theta_{12}$  and then expressing the other elements  
 148 in terms of powers of  $\lambda$ :

$$V_{CKM} \approx \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} \quad (1.8)$$

149 where  $A$ ,  $\rho$  and  $\eta$  are all real numbers intended to express the order of magnitude  
 150 differences between  $s_{12}$  and the other elements in the matrix. Of course, all the elements  
 151 should be the same irrespective of which parametrization is used.

152 The elements of the CKM matrix have been measured and the latest accepted results  
 153 are summarized in 1.10 [7]. The interaction strength is then proportional to  $|V_{ij}|^2$ .  
 154 Unitarity of the CKM implies that the probability of transition from any up-type quark  
 155 to any down-type is the same,

$$\sum_k |V_{ik}|^2 = \sum_i |V_{ik}|^2 = 1 \quad (1.9)$$

156 for all  $i$  quark generations [12]. Note that the term  $V_{tb}$  is approximately unity and by far  
 157 dominates over the other  $V_{tj}$  terms. This means that the top-quark transitions almost  
 158 exclusively into a  $b$ -quark ( $t \rightarrow Wb$ ) with transitions  $t \rightarrow Ws$  and  $t \rightarrow Wd$  having a  
 159 probability of less than 1%. The soft muon tagger which is the focus of this thesis relies

on weak semileptonic decays of  $b$ -quarks. From 1.10 one can see that the transition  $b \rightarrow c$  dominates over  $b \rightarrow u$ . Additionally the focus of this thesis is on semileptonic  $t\bar{t}$  events, where one of the  $W$  bosons in the event decays to quarks as per the magnitude of  $V_{ij}$ .

$$V_{CKM} = \begin{pmatrix} 0.97427 \pm 0.00015 & 0.22534 \pm 0.00065 & 0.00351^{+0.00015}_{-0.00014} \\ 0.22520 \pm 0.00065 & 0.97344 \pm 0.00016 & 0.0412^{+0.0011}_{-0.0005} \\ 0.00867^{+0.00029}_{-0.00031} & 0.0404^{+0.0011}_{-0.0005} & 0.999146^{+0.000021}_{-0.000046} \end{pmatrix} \quad (1.10)$$

An additional unique feature of weak interactions is that the charge conjugation-parity ( $CP$ ) symmetry is violated. The operator  $C$  denotes the change of a particle by its antiparticle partner and  $P$  denotes a reversal of helicity (the projection of spin onto the momentum of a particle). A clear violation of  $C$  and  $P$  was observed in the radioactive decay of Cobalt-60, where the resulting electrons were preferentially emitted in the opposite direction of the nuclear spin of the Cobalt [13]. Thus weak currents only couple to left-handed neutrinos (or right-handed antineutrinos) which is a violation of parity. Additionally charge symmetry is also violated since a left-handed neutrino is preferentially picked over a left-handed antineutrino. Finally in 1964  $CP$  violation was observed in the decay of neutral kaon [14].

Thus the probability of  $\bar{a} \rightarrow \bar{b}$  is not equal to that of  $a \rightarrow b$ . The existence of  $CP$  violation has interesting consequences for the formation of the early universe. The preferential production of matter over antimatter in  $CP$  violating interactions would shift the balance in favour of matter resulting in a universe similar to our own. Finally as with QCD, weak interactions couple weak bosons to each other. Thus the vertex  $Z \rightarrow W^-W^+$  is allowed via the weak force.

### 1.3.1 Electroweak Unification and the Higgs mechanism

The unification of the electromagnetic and weak theories was first proposed by Glashow and later developed by Weinberg and Salam into the electroweak theory [15–17]. The theory postulates that while at low energies the two forces are to be treated separately, at higher the two can be seen as a single force. Thus the two forces are different manifestations of the same “electroweak” interaction. There were several stumbling blocks to

the unification of the forces. Firstly, the boson which drives the electromagnetic interaction, the photon, is massless while the weak bosons are both massive. Evidence for the massive nature of these bosons has been established by experimental results from [18] at CERN.

Thus the symmetry of the theory must be spontaneously broken in some way. A mechanism for ElectroWeak Symmetry Breaking (EWSB) was postulated by Higgs, Brout, Englert and others which introduces mass to the weak bosons and posits the existence of an additional scalar (spin  $S = 0$ ) boson known as the Higgs boson.

### Gauge Theories

Gauge invariance is one of the underlying invariances which underpins the Standard Model. Given the so-called Dirac lagrangian<sup>3</sup>

$$\mathcal{L} = i\hbar c \bar{\psi} \gamma^\mu \partial_\mu \psi - mc^2 \bar{\psi} \psi \quad (1.11)$$

which describes a free particle of spin- $\frac{1}{2}$  with mass  $m$ . Note that it is invariant under the transformation

$$\psi \rightarrow e^{i\theta} \psi, \text{ where } \theta \text{ is a real number} \quad (1.12)$$

since the adjoint  $\bar{\psi} \rightarrow e^{-i\theta} \bar{\psi}$  and the two terms cancel out. This is known as a (*global*) *gauge transformation* since  $\theta$  is the same at all points of space-time. A (*local*) *gauge transformation* occurs when the phase is different for different points in space-time:

$$\psi \rightarrow e^{i\theta(x)} \psi \quad (1.13)$$

The Dirac lagrangian (Equation 1.11) is not invariant under a local gauge transformation since extra terms are created by the derivative. This then implies that the underlying physics of such a theory depends on position in space-time. Thus local gauge invariance must be imposed. In the case of the Dirac lagrangian, this is done by introducing additional terms to the Dirac lagrangian which will cancel the extra terms introduced by the local gauge transformation. As it turns out this results in the introduction of a

---

<sup>3</sup>A Lagrangian is a mathematical function that describes the underlying dynamics of a system as a function of time and space coordinates ( $x^\mu$ ) and their time derivatives.

new massless vector field that couples to  $\psi$ .

The new lagrangian then describes a spin- $\frac{1}{2}$  particle with mass  $m$  that interacts with a free massless field. This new field can be identified as the electromagnetic field and the spin- $\frac{1}{2}$  particles are electrons and positrons. Thus the resulting lagrangian describes all interactions that form part of quantum electrodynamics.

A similar procedure can be applied to the color quark model and obtain a description of all QCD interactions. However requiring that the weak theory be a gauge theory (invariant under local gauge transformation) encounters a problem since the weak bosons are known to be massive. There must be some mechanism via which the  $W^\pm$  and  $Z^0$  obtain mass.

The Higgs mechanism posits the existence of a complex scalar field doublet that when introduced into the electroweak Lagrangian results in the weak fields acquiring a mass term. In other words the  $W^\pm$  and  $Z^0$  interact with the Higgs field and obtain a mass. An additional consequence of introducing the Higgs field is the inclusion of a scalar boson particle, the so-called ‘‘Higgs boson’’. Finally the Higgs field also couples to fermions via the Yukawa coupling generating gauge invariant mass terms for the fermions as well.<sup>4</sup>

The SM Lagrangian in its current form including the Higgs potential is shown in Equation 1.14. This expression describes all possible particle interactions that form part of the SM, of particular interest are the fermion mass term which couples the fermion field ( $\psi$ ) to the scalar Higgs field ( $\phi$ ) and the Higgs kinetic and potential terms.

$$\begin{aligned}
\mathcal{L} = & - \underbrace{\frac{1}{4} W_{\mu\nu}^a W^{\mu\nu a}}_{\text{Weak Field}} - \underbrace{\frac{1}{4} B_{\mu\nu} B^{\mu\nu}}_{\text{EM Field}} - \underbrace{\frac{1}{4} G_{\mu\nu}^a G^{\mu\nu a}}_{\text{Strong Field}} \\
& + \underbrace{\bar{\psi} \not{D}_\mu \psi}_{\text{Fermion Kinetic}} + \underbrace{\lambda \bar{\psi} \psi \phi}_{\text{Fermion Mass}} \\
& + \underbrace{|D_\mu \phi|^2}_{\text{Higgs Kinetic}} - \underbrace{V(\phi)}_{\text{Higgs Potential}}
\end{aligned} \tag{1.14}$$

228

---

<sup>4</sup>For a more complete description of the mathematical procedure see [6].

---

## 229 Bibliography

- 230 [1] B. Odom, D. Hanneke, B. D’Urso, and G. Gabrielse. New Measurement of the  
231 Electron Magnetic Moment Using a One-Electron Quantum Cyclotron. *Phys. Rev.*  
232 *Lett.*, 97:030801, Jul 2006.
- 233 [2] The ATLAS Collaboration. Observation of a new particle in the search for the  
234 Standard Model Higgs boson with the ATLAS detector at the LHC. *Physics Letters*  
235 *B*, 716(1):1 – 29, 2012.
- 236 [3] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with  
237 the CMS experiment at the LHC. *Phys.Lett.*, B716:30–61, 2012.
- 238 [4] P.A.R. Ade et al. Planck 2013 results. I. Overview of products and scientific results.  
239 2013.
- 240 [5] D.H. Perkins. *Introduction to High Energy Physics*. Cambridge University Pres,  
241 2000.
- 242 [6] D. Griffiths. *Introcution to Elementary Particles*. John Wiley & Sons, Inc, 1987.
- 243 [7] J. Beringer et al. (Particle Data Group). The Review of Particle Physics. *Physical*  
244 *Review D*, 86:010001+, 2012.
- 245 [8] Nicola Cabibbo. Unitary Symmetry and Leptonic Decays. *Phys. Rev. Lett.*, 10:531–  
246 533, Jun 1963.
- 247 [9] Makoto Kobayashi and Toshihide Maskawa. Cp-violation in the renormalizable  
248 theory of weak interaction. *Progress of Theoretical Physics*, 49(2):652–657, 1973.

- [10] Ling-Lie Chau and Wai-Yee Keung. Comments on the parametrization of the kobayashi-maskawa matrix. *Phys.Rev.Lett.*, 53:1802, 1984.
- [11] Lincoln Wolfenstein. Parametrization of the Kobayashi-Maskawa Matrix. *Phys. Rev. Lett.*, 51:1945–1947, Nov 1983.
- [12] K. Marathe. *Topics in Physical Mathematics*. Springer, 2010.
- [13] J. J. Aubert, U. Becker, P. J. Biggs, J. Burger, M. Chen, G. Everhart, P. Goldhagen, J. Leong, T. McCorriston, T. G. Rhoades, M. Rohde, Samuel C. C. Ting, Sau Lan Wu, and Y. Y. Lee. Experimental observation of a heavy particle j. *Physical Review Letters*, 33(23):1404–1406, 1974.
- [14] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay. Evidence for the 2 decay of the k20 meson. *Physical Review Letters*, 13(4):138–140, 1964.
- [15] Steven Weinberg. A model of leptons. *Physical Review Letters*, 19(21):1264–1266, 1967.
- [16] S. L. Glashow, J. Iliopoulos, and L. Maiani. Weak Interactions with Lepton-Hadron Symmetry. *Phys. Rev. D*, 2:1285–1292, Oct 1970.
- [17] Abdus Salam. Weak and Electromagnetic Interactions. *Conf.Proc.*, C680519:367–377, 1968.
- [18] G. Arnison, A. Astbury, B. Aubert, C. Bacci, G. Bauer, A. Bzaguét, R. Bck, T.J.V. Bowcock, M. Calvetti, T. Carroll, P. Catz, P. Cennini, S. Centro, F. Ceradini, S. Cittolin, D. Cline, C. Cochet, J. Colas, M. Corden, D. Dallman, M. DeBeer, M. Della Negra, M. Demoulin, D. Denegri, A. Di Ciaccio, D. DiBitonto, L. Dobrzynski, J.D. Dowell, M. Edwards, K. Eggert, E. Eisenhandler, N. Ellis, P. Erhard, H. Faissner, G. Fontaine, R. Frey, R. Frhwirth, J. Garvey, S. Geer, C. Ghesquire, P. Ghez, K.L. Giboni, W.R. Gibson, Y. Giraud-Hraud, A. Givernaud, A. Gonidec, G. Grayer, P. Gutierrez, T. Hansl-Kozanecka, W.J. Haynes, L.O. Hertzberger, C. Hodges, D. Hoffmann, H. Hoffmann, D.J. Holthuisen, R.J. Homer, A. Honma, W. Jank, G. Jorat, P.I.P. Kalmus, V. Karimki, R. Keeler, I. Kenyon, A. Kernan,

- 276 R. Kinnunen, H. Kowalski, W. Kozanecki, D. Kryn, F. Lacava, J.-P. Laugier, J.-  
277 P. Lees, H. Lehmann, K. Leuchs, A. Lvque, E. Linglin, E. Locci, M. Loret, J.-J.  
278 Malosse, T. Markiewicz, G. Maurin, T. McMahon, J.-P. Mendiburu, M.-N. Mi-  
279 nard, M. Moricca, H. Muirhead, F. Muller, A.K. Nandi, L. Naumann, A. Norton,  
280 A. Orkin-Lecourtois, L. Paoluzi, G. Petrucci, G. Piano Mortari, M. Pimi, A. Placci,  
281 E. Radermacher, J. Ransdell, H. Reithler, J.-P. Revol, J. Rich, M. Rijssenbeek,  
282 C. Roberts, J. Rohlf, P. Rossi, C. Rubbia, B. Sadoulet, G. Sajot, G. Salvi, J. Salvini,  
283 J. Sass, A. Saudraix, A. Savoy-Navarro, D. Schinzel, W. Scott, T.P. Shah, M. Spiro,  
284 J. Strauss, K. Sumorok, F. Szoncs, D. Smith, C. Tao, G. Thompson, J. Timmer,  
285 E. Tscheslog, J. Tuominiemi, S. Van der Meer, J.-P. Vialle, J. Vrana, V. Vuillemin,  
286 H.D. Wahl, P. Watkins, J. Wilson, Y.G. Xie, M. Yvert, and E. Zurfluh. Experimen-  
287 tal observation of isolated large transverse energy electrons with associated missing  
288 energy at  $\sqrt{s}=540$  GeV. *Physics Letters B*, 122(1):103 – 116, 1983.