Effective Superluminal Travel: Conditional Geometry, Source Exclusions, and a Gated Research Program

SIGMA AI-assisted research synthesis

Abstract

Can an effective spacetime shortcut remain coherent once its causal structure, source dynamics, and propagation limits are examined together? This theoretical study separates a prescribed geometric shortcut from a physically realizable transport mechanism. A smooth finite-duration metric permits an exterior-clock transit shorter than a matched Minkowski reference while its carrier remains locally timelike and a global temporal function excludes closed causal curves. Under Einstein gravity, however, the target requires negative normal energy and strict null-energy violation. The supplied analytical arguments exclude a canonical scalar and a regular, positive-energy single-scalar extension as sources.

A related eternal superluminal geometry also has incomplete forward axial null geodesics with divergent parallelly propagated curvature, preventing a regular extension through their affected endpoints. A preferred-time wider-cone free field offers a distinct conceptual comparison, but not an interacting transport theory. These results support a staged program of source, finite-duration, quantum, and control tests; they establish neither a working drive nor a universal prohibition on effective superluminal travel.

In brief

The question. Can a traveler reach a distant point sooner than a matched light signal without locally outrunning light?

The answer. A prescribed spacetime geometry can produce that timing effect while the traveler remains locally timelike. Under the examined Einstein-gravity assumptions, it also needs negative energy and defeats the simple scalar sources tested here.

The route. Define the clock comparison, test the geometry's local causality, calculate its required source, then examine propagation and remaining feasibility gates.

The limit. No physically realizable source, stable transport system, or working faster-than-light drive is established.

1 Introduction and physical framework

The answerable question is whether the specified geometric shortcut survives necessary tests of causal consistency, admissible sourcing, and regular propagation—not whether an arbitrarily chosen metric can be called a drive. Effective superluminality is defined by an exterior-clock transit shorter than the corresponding light transit in a matched, unmodified Minkowski geometry. It is distinct from a massive carrier leaving its local metric light cone.

The bounded contribution combines conditional geometric deductions with source-class exclusions and a restricted geodesic obstruction. A preferred-time wider-cone field supplies a conceptually different comparison, but its free-field status prevents an equal-depth assessment of interacting transport. The investigation therefore ends with explicit decision gates rather than a claim that either framework is physically viable.

2 Models, methods, and evidence

The saved analysis prescribed smooth unit-lapse, flat-spatial-slice geometries and examined their local tangent norms, exterior-clock transit protocols, and temporal functions. It treated the finite-duration pulse separately from the eternal constant-speed geometry. The geometric calculations used signature $(-,+,+,+)$; source calculations assumed Einstein gravity with zero cosmological constant and $G_{ab} = 8\pi G T_{ab}/c^4$.

The source analysis used the Hamiltonian and momentum constraints, spatial evolution equations, and a separate frame-curvature calculation to examine the required density, stresses, fluxes, and null contractions. A spherical-wall estimate applied angular integration and Cauchy–Schwarz. Initial-value uniqueness, canonical-scalar stress, and the quadratic Hamiltonian and principal symbol of a minimally coupled $P(X,\phi)$ scalar then supplied distinct formation and source-stability tests.

The propagation analysis solved axial affine equations using a translating Killing symmetry and distinguished a subluminal complete-ray integral from superluminal rays approaching characteristic separatrices. Parallelly propagated tidal curvature tested whether the latter endpoints were removable coordinate effects, while the backward family supplied a control. These were analytical arguments; the supplied evidence contains no completed numerical simulation, measured observation, or inspected domain-literature assessment.

This synthesis uses a bounded selection of the archived research. Unreviewed records could affect its conclusions; the full omission ledger remains available with the generation record.

3 An operational shortcut without local superluminal motion

The first question is geometric: can a prescribed spacetime yield an earlier arrival on matched distant clocks while the carrier remains locally slower than light? Use signature $(-,+,+,+)$ and a smooth, nondecreasing center path $x_s(t)$ that starts at 0 before $t = 0$ and ends at L after $t = T$, with $L > cT$. Let $v(t) = dx_s/dt$, $r^2 = [x - x_s(t)]^2 + y^2 + z^2$, and let the smooth profile $f(r)$ equal one near the center and vanish for $r \geq R$. The assigned metric is

$$ds^2 = -c^2dt^2 + [dx - v(t)f(r)dt]^2 + dy^2 + dz^2.$$

It is a target geometry, not a sourced solution.

Example: What the distant clocks compare

Place two exterior stations a distance L apart. A light signal in the matched flat-space reference needs L/c to make the trip. The prescribed pulse carries the central traveler between them in a time $T < L/c$, while the traveler remains timelike in the pulse geometry. The earlier arrival compares two specified spacetime models; it does not demonstrate a buildable source or a local speed above light.

First test local motion. The invertible coframe $(c\,dt,\ dx - v f\,dt,\ dy,\ dz)$ gives Lorentzian signature and $\det(g_{\mu\nu}) = -c^2$, even where the coordinate speed v exceeds c. On the central path $x = x_s(t)$, $y = z = 0$, the shifted spatial differential vanishes, so $ds^2 = -c^2dt^2$ and $d\tau = dt$. The carrier is timelike throughout. Relative to observers normal to the time slices, the causal-cone condition is

$$(dx - v f\,dt)^2 + dy^2 + dz^2 \le c^2 dt^2.$$

The carrier's spatial displacement in that local frame is zero.

Next test distant arrival. Put stationary stations A and B at $x = -D$ and $x = L + D$, with $D > R$, outside the moving region; synchronize their clocks in the common Minkowski past. At $t = -D/c$, A sends a light signal through the flat pre-pulse region to the carrier at $(0, 0)$. The carrier carries that information to (T, L) and relays it through the flat post-pulse region to B, which receives it at $t = T + D/c$. The exterior-clock interval is

$$\Delta t_{\text{shortcut}} = T + 2D/c < (L + 2D)/c = \Delta t_{\text{Minkowski}}.$$

The strict inequality is exactly $L > cT$.

This compares the assigned geometry with an unmodified Minkowski light transit between the same stationary endpoints. It does not compare a timelike path against the fastest null path in one geometry. The stations, relays, and carrier are idealized test systems with neglected backreaction; the geometric schedule is specified independently of the newly transmitted information. The clock calculation establishes a conditional arrival advantage, not an on-demand method of creating the pulse.

Finally test this geometry's causal ordering. Its inverse metric has $g^{tt} = -1/c^2$, so the gradient of t is everywhere timelike. With the stated future orientation, t strictly increases on every nontrivial future-directed causal curve. That excludes closed causal curves in this single prescribed spacetime. It says nothing by itself about assemblies of moving corridors, additional transport channels, or repeated trips.

The geometric answer is therefore limited but exact: exterior coordinate speed above c need not mean local superluminal motion or a causal loop in this construction. The unresolved physical question is whether admissible source dynamics can form and maintain the assigned geometry.

4 The Einstein source: negative energy and null stress

The Einstein source calculation turns the prescribed shortcut geometry into a precise material requirement: negative energy measured by the slice-normal observers and, independently, strict




Figure 1: A prescribed finite-duration metric separates local motion from an exterior-clock shortcut. A central carrier remains timelike while the chosen distant-clock comparison can register an earlier arrival; this geometric result does not provide a material source or a working drive.

Source-linked claims: C31, C32; evidence links appear in the claim notes.

negative null stress in the wall. Consider the constant-speed geometry $ds^2 = -(dx^0)^2 + [dx - u\,dx^0]^2 + dy^2 + dz^2$, with $x^0 = ct$, $u = \beta f(r)$, constant $\beta = v/c$, and $r^2 = (x - \beta x^0)^2 + y^2 + z^2$. The dimensionless profile is smooth, equals one for $r \le R$, and vanishes for $r \ge R + \Delta$, where $R, \Delta > 0$ are lengths. Throughout this calculation, assume signature $(-,+,+,+)$, zero cosmological constant, and Einstein's equation $G_{ab} = 8\pi G T_{ab}/c^4$.

The metric has unit lapse, Euclidean spatial metric $h_{ij} = \delta_{ij}$, and shift $B^i = (-u, 0, 0)$. An orthonormal frame consists of the future-directed unit normal $n = \partial_0 + u\partial_x$ and spatial vectors $e_i = \partial_i$. Define the measured projections by $\varepsilon = T(n,n)$, $j_i = -T(n,e_i)$, and $S_{ij} = T(e_i,e_j)$; consequently, the covariant frame components satisfy $T_{0i} = -j_i$. These projections all have units $\mathrm{J\,m^{-3}}$, and the momentum-flux sign is essential when forming null contractions.

Adopt the extrinsic-curvature convention $K_{ij} = -(\mathcal{L}_n h)_{ij}/2$. Since the spatial metric is time independent, $K_{ij} = (\partial_i B_j + \partial_j B_i)/2$, so its nonzero independent components are $K_{xx} = -\beta f_x$, $K_{xy} = -\beta f_y/2$, and $K_{xz} = -\beta f_z/2$, where subscripts on the profile denote spatial partial derivatives. Thus $K^2 = \beta^2 f_x^2$, whereas $K_{ij}K^{ij} = \beta^2 f_x^2 + \beta^2(f_y^2 + f_z^2)/2$. The longitudinal derivative contributes equally to these two quadratic expressions; only the transverse derivatives survive their difference.

The Hamiltonian constraint is ${}^{(3)}R + K^2 - K_{ij}K^{ij} = 16\pi G\varepsilon/c^4$. Because the slices are intrinsically flat, it gives $\varepsilon = -c^4\beta^2(f_y^2 + f_z^2)/(32\pi G)$. For nonzero β, the energy density is therefore strictly negative wherever the transverse wall gradient is nonzero, violating the weak energy condition and hence the dominant energy condition. For a spherical profile, $f_y^2 + f_z^2 = [f'(r)]^2 \sin^2\theta$, with θ measured from the translation axis, so every nontrivial smooth wall contains an open negative-density region.

The angular weighting also fixes the coefficient in the integrated requirement. The proper spatial volume is d^3x, and $\int \sin^2\theta\, d\Omega = 8\pi/3$, giving the normal-frame slice integral

$$E_n \equiv \int \varepsilon\, d^3x = -[c^4\beta^2/(12G)] \int_R^{R+\Delta} r^2[f'(r)]^2\, dr.$$

The radial weight matters: a bound based only on the coordinate thickness would omit the spherical wall's area dependence.

This negative volume integral is not the asymptotic gravitational energy or a demonstrated propulsion expenditure. The standard Arnowitt–Deser–Misner energy surface integral vanishes because the spatial metric is exactly Euclidean, and the compact support of the extrinsic curvature likewise gives zero asymptotic momentum. Thus negative E_n must not be identified with negative total mass, and its magnitude is not an established fuel requirement. There is no contradiction with a positive-energy theorem whose usual dominant-energy premise is violated by the required source.

A profile-independent lower bound follows from the endpoint change, without assuming monotonicity. Since $\int_R^{R+\Delta} f'(r)\, dr = -1$, weighted Cauchy–Schwarz gives

$$1 \le [\int_R^{R+\Delta} r^2[f'(r)]^2\, dr][\int_R^{R+\Delta} r^{-2}\, dr].$$

The second factor is $\Delta/[R(R+\Delta)]$, and therefore $|E_n| \ge [c^4\beta^2/(12G)]R(R+\Delta)/\Delta$. Equality would require the radial derivative to be proportional to r^{-2} throughout the wall, which cannot smoothly join both constant-profile regions.

For a thin wall, $\Delta \ll R$, the leading lower-bound scale is $c^4\beta^2R^2/(12G\Delta)$: reducing the transition thickness increases the required magnitude rather than removing the negative energy. Since c^4/G has units $\mathrm{J\,m^{-1}}$, the bound has energy units. Its scope is the stated spherical, unit-lapse, flat-slice ansatz with $R > 0$, not arbitrary shortcut geometries or a universal cost of effective superluminal travel. The normal density and its slice integral vanish when $\beta = 0$ or the profile is constant.

Negative normal energy alone does not establish violation of the null energy condition; the spatial stresses and momentum flux must also be included. To obtain the relevant components, write $C = c^4/(8\pi G)$, $Q = u_y^2 + u_z^2$, and $L = u_{yy} + u_{zz}$. With the stated signs, the momentum constraint is $j_i/C = \partial_j K^j{}_i - \partial_i K$, yielding $j_x = -CL/2$, while the Hamiltonian result becomes $\varepsilon = -CQ/4$. Both Q and L have units $\mathrm{m^{-2}}$.

The longitudinal stress follows from the spatial Einstein evolution equation rather than an assumed equation of state. With $S = \delta^{ij}S_{ij}$, this equation reads

$$(\partial_0 - \mathcal{L}_B)K_{ij} = KK_{ij} - 2K_{ik}K^k{}_j - [S_{ij} - \delta_{ij}(S-\varepsilon)/2]/C.$$

Define $a = u_x$ and let $D = n^a\partial_a$ act on scalars; for the translating profile, $D = (u-\beta)\partial_x$. If P_{ij} denotes the geometric combination $KK_{ij} - 2K_{ik}K^k{}_j - (\partial_0 - \mathcal{L}_B)K_{ij}$, its longitudinal component and trace are $P_{xx} = Da + a^2 - Q/2$ and $P = Da + a^2$.

Tracing and then inverting the evolution equation gives $S/C = 3\varepsilon/C - 2P$ and $S_{ij}/C = P_{ij} + \delta_{ij}(\varepsilon/C - P)$. Consequently, $S_{xx} = C[(Da + a^2 - Q/2) - Q/4 - (Da + a^2)] = -3CQ/4$. The longitudinal derivative and evolution terms cancel in this stress projection, leaving a negative contribution fixed by the same transverse-gradient quantity that determines the normal density.

For the future-directed null vectors $k_\pm = n \pm e_x$, the measured source components now give $T(k_\pm, k_\pm) = \varepsilon + S_{xx} \mp 2j_x = C(-Q \pm L)$. Equivalently,

$$T(k_\pm, k_\pm) = [c^4/(8\pi G)][-\beta^2(f_y^2 + f_z^2) \pm \beta(f_{yy} + f_{zz})].$$

The second-derivative flux term changes sign between the two directions, but the transverse-gradient term is negative in both.

The same cancellation is visible directly in curvature. Using $R(X,Y)Z = \nabla_X\nabla_Y Z - \nabla_Y\nabla_X Z - \nabla_{[X,Y]}Z$, the orthonormal projections are $R_{nn} = -Da - a^2 - Q/2$, $R_{xx} = Da + a^2 - Q/2$, and $R_{nx} = L/2$. Since contraction with a null vector removes the scalar-curvature term in Einstein's equation, these expressions reproduce $T(k_\pm, k_\pm) = C(-Q \pm L)$. This supplies a second analytic route to the null result, distinct from inferring it from the negative density.

The two contractions sum to $-2CQ$, so at least one is strictly negative wherever $Q > 0$. Their minimum is $C(-Q - |L|)$; if $Q = 0$ but $L \neq 0$, one direction still violates the null energy condition, whereas $Q = L = 0$ leaves other null directions undecided. Every nontrivial smooth spherical wall with $\beta \neq 0$ has an open region with $Q > 0$, independently of monotonicity and without requiring $|\beta| > 1$. The source obstruction is therefore a property of this metric family even below the nominal superluminal threshold, not an effect that first appears at that threshold.

These are pointwise statements, not an averaged null-energy theorem. An averaged condition requires a specified complete, affinely parametrized null geodesic, its tangent normalization, and convergence of the stress integral; the locally defined pair $k_\pm$ cannot simply be integrated as if it supplied those data. Likewise, defining $T_{ab} = c^4 G_{ab}/(8\pi G)$ ensures $\nabla^a T_{ab} = 0$ by the contracted Bianchi identity, but supplies neither a material constitutive law nor preparable data with stable coupled evolution.

The calculation therefore excludes everywhere-null-energy-respecting matter as the source of this exact target and makes reproduction of its negative null stress—not merely its geometry—the necessary next condition for any proposed transport mechanism.

5 Formation from complete data and the canonical-scalar counterexample

Formation is a constraint on complete dynamical data, not merely on the geometry prescribed for a later transit. Under autonomous classical equations with a unique Cauchy development, complete vacuum data cannot evolve into a nontrivial curvature pulse. A canonical scalar with a sign-changing potential provides a substantive test of whether initially hidden, stress-free matter can evade this obstruction: negative energy is algebraically available, but the admissible stress-free data and their subsequent evolution still fail to produce the required source.

The vacuum obstruction requires three explicit hypotheses. The autonomous Einstein–matter theory must admit Minkowski spacetime with matter vacuum Ψ_0; the initial surface must be a complete Cauchy surface $\Sigma \cong \mathbb{R}^3$ carrying $h_{ij} = \delta_{ij}$, $K_{ij} = 0$, $\Psi = \Psi_0$, and every independent matter momentum at its vacuum value; and the smooth maximal globally hyperbolic development must be

unique up to gauge and diffeomorphism. Minkowski vacuum is already a development of these data, so uniqueness excludes an inequivalent later pulse within that development.

Incoming preparation must therefore be represented in the complete initial data, or enter through a boundary if the chosen surface was not actually a Cauchy surface.

Flat gravitational data alone do not establish the matter-vacuum premise. The gravitational constraints fix particular projections of the stress tensor, whereas an exactly Minkowski spacetime region fixes the entire tensor through Einstein's equation. An external actuator likewise cannot be omitted from the source description: its fields, stress, and preparation data belong to the enlarged system. These distinctions separate a genuine vacuum-formation obstruction from the stronger, generally invalid inference that initially flat geometry contains no dynamical matter.

For an explicit classical test, adopt signature $(-,+,+,+)$, natural units $c = \hbar = 1$, ordinary Einstein gravity with zero cosmological constant, and $\kappa = 8\pi G$. The minimally coupled real scalar is specified by

$$S = \int d^4x\,\sqrt{-g}\left[\frac{R}{2\kappa} - \frac{1}{2}g^{\mu\nu}\partial_\mu\phi\,\partial_\nu\phi - V(\phi)\right],$$
$$V(\phi) = \frac{\lambda}{a^2}\phi^2(\phi^2 - a^2)(\phi^2 - 4a^2), \quad \lambda > 0, \quad a > 0.$$

Here ϕ and a have energy units, λ is dimensionless, V has energy-density units, and κ has units of inverse energy squared. This polynomial defines a classical source model, not a demonstrated quantum-complete theory.

The potential is bounded below but negative for $a < |\phi| < 2a$, so rejecting the source cannot rest on an assumed nonnegative potential. Its vacuum at $\phi = 0$ satisfies $V(0) = V'(0) = 0$ and $V''(0) = 8\lambda a^2 > 0$. Variation gives

$$G_{\mu\nu} = \kappa T_{\mu\nu},$$
$$\Box\phi - V'(\phi) = 0,$$
$$T_{\mu\nu} = \partial_\mu\phi\,\partial_\nu\phi - g_{\mu\nu}\left[\frac{1}{2}(\partial\phi)^2 + V(\phi)\right],$$

where

$$(\partial\phi)^2 = g^{\alpha\beta}\partial_\alpha\phi\,\partial_\beta\phi.$$

The analysis concerns smooth solutions in their unique regular globally hyperbolic development; it assumes no extension through a singularity.

Let n denote the unit future normal to an initial slice, h_{ij} its induced metric, $\pi = n^\mu\partial_\mu\phi$ the scalar normal derivative, and $q_i = D_i\phi$ its spatial gradient, with $q^2 = h^{ij}q_iq_j$. The normal energy density, momentum density, and spatial stress are

$$\rho = \frac{1}{2}(\pi^2 + q^2) + V, \qquad j_i = -\pi q_i, \qquad S_{ij} = q_iq_j + h_{ij}\left[\frac{1}{2}(\pi^2 - q^2) - V\right].$$

Complete Einstein–scalar data comprise (h, K, ϕ, π) subject to the gravitational constraints. In particular, $h_{ij} = \delta_{ij}$ and $K_{ij} = 0$ require $\rho = 0$ and $j_i = 0$, but do not by themselves require $S_{ij} = 0$.

Requiring the full tensor to vanish gives a much sharper classification than requiring only its constraint projections to vanish. For every unit spatial vector e, the scalar tensor obeys

$T(n, n) + T(e, e) = \pi^2 + (e \cdot q)^2.$

If $T_{\mu\nu} = 0$, this nonnegative expression vanishes for every e, forcing $\pi = 0$ and $q_i = 0$; the remaining condition $\rho = 0$ then gives $V = 0$. Conversely, these three conditions make every component of the stress tensor vanish.

On a connected slice, all smooth stress-free scalar data therefore have spatially constant field value $\phi = C$, with $C \in \{0, \pm a, \pm 2a\}$. This result uses no assumption that $V \geq 0$ and leaves no smooth localized stress-free profile. The asymptotic matter condition $\phi \to 0$ retains only $C = 0$, but relaxing that condition admits four nonvacuum, stress-free constants. They are genuine alternatives to complete matter-vacuum data, even though their initial stress tensors vanish.

The nonzero roots are not stationary solutions, because $V'(\pm a) = \mp 6\lambda a^3$ and $V'(\pm 2a) = \pm 48\lambda a^3$. Thus the complete data $(\delta, 0, C, 0)$ satisfy both gravitational constraints and have $T_{\mu\nu} = 0$ initially, yet the scalar equation supplies a nonzero acceleration whenever $C \neq 0$. These data are a concrete counterexample to identifying an initially vanishing Einstein tensor with complete vacuum data. They do not contradict vacuum-data uniqueness, because they omit its matter-vacuum premise.

Their evolution can be determined without assuming that the initially hidden field becomes a localized wall. Translations and rotations preserve $(\delta, 0, C, 0)$, and uniqueness preserves those symmetries in the development. Locally, the solution consequently has spatially flat Friedmann–Lemaître–Robertson–Walker form

$ds^2 = -d\tau^2 + b(\tau)^2 d\mathbf{x}^2, \qquad \phi = \phi(\tau),$

where τ is proper time, b is the dimensionless scale factor, and $H = \dot{b}/b$, with a dot denoting differentiation with respect to τ. The initial conditions are $b(0) = 1$, $H(0) = 0$, $\phi(0) = C$, and $\dot{\phi}(0) = 0$.

The homogeneous coupled equations are

$$3H^2 = \kappa\rho, \qquad \dot{H} = -\frac{\kappa}{2}\dot{\phi}^2, \qquad \ddot{\phi} + 3H\dot{\phi} + V' = 0, \qquad \rho = \frac{1}{2}\dot{\phi}^2 + V.$$

The scalar equation gives the conservation law $\dot{\rho} = -3H\dot{\phi}^2$, and hence $d(3H^2 - \kappa\rho)/d\tau = 0$. This verifies propagation of the Hamiltonian constraint without dividing by the initially vanishing H. On every regular portion of the development, it follows that $\rho = 3H^2/\kappa \geq 0$.

This nonnegative energy is not an artifact of choosing the homogeneous normal. With homogeneous pressure $p = \dot{\phi}^2/2 - V$, one has $\rho + p = \dot{\phi}^2$, and any unit timelike observer w with Lorentz factor γ relative to that normal measures

$T(w, w) = (\rho + p)\gamma^2 - p = \rho + (\gamma^2 - 1)\dot{\phi}^2 \geq 0.$

Thus every stress-free flat-slice development classified above satisfies the weak energy condition

throughout its regular evolution, even when the field enters the negative-potential region. Spatial homogeneity also prevents these data from generating a compact moving wall.

A local expansion shows explicitly why initially vanishing stress does not imply a flat time interval. Choose $C = a$ and define $A = -V'(a) = 6\lambda a^3$; for $|V''(a)|\tau^2 \ll 1$ and $\kappa A^2\tau^4 \ll 1$, the smooth equations give

$$\begin{aligned}\phi &= a + \frac{A}{2}\tau^2 + O(\tau^4),\\ H &= -\frac{\kappa A^2}{6}\tau^3 + O(\tau^5),\\ \rho &= \frac{\kappa A^4}{12}\tau^6 + O(\tau^8),\\ p &= A^2\tau^2 + O(\tau^4).\end{aligned}$$

Kinetic and potential energy cancel at quadratic order in proper time, but pressure does not, and the Ricci scalar becomes $R = \kappa(\rho - 3p) = -3\kappa A^2\tau^2 + O(\tau^4)$. This is a short-time analytical expansion, not a stability result; it demonstrates curvature on both sides of the initially stress-free slice rather than an exactly flat prehistory followed by a pulse.

An exactly Minkowski open prehistory eliminates these nonstationary alternatives more directly. Einstein's equation requires $T_{\mu\nu} = 0$ throughout that region, so the stress-free classification forces the scalar to be spacetime-constant there; its field equation additionally requires $V' = 0$. Among the zero-potential roots, only $C = 0$ satisfies both requirements. A complete slice in that prehistory therefore carries precisely the vacuum data $(\delta, 0, 0, 0)$, and uniqueness prohibits subsequent pulse formation, regardless of a prescribed smooth switching schedule.

There is also a source exclusion stronger than the prehistory argument. For every metric-null vector k, the canonical scalar obeys $T(k,k) = (k^\mu\partial_\mu\phi)^2 \geq 0$, independently of the sign of its potential, whereas the exact shortcut target requires a strictly negative null contraction somewhere in its moving wall. Consequently, even inhomogeneous preparation in the negative-potential region cannot make this unchanged scalar source the exact target. The formation analysis therefore changes the task from finding a better preparation of this scalar to changing the source theory or the target geometry; it excludes this source–target combination, not effective superluminal travel in every framework.

6 A local stability gate for nonlinear scalar sources

Nonlinear first-derivative scalar dynamics do not remove the target's source obstruction if the scalar must retain positive high-frequency energy and a regular wave equation on a common matter–gravity Cauchy slicing. The relevant exclusion concerns a single, smooth, minimally coupled real scalar with Lagrangian $P(X, \phi)$ in ordinary positive-sign Einstein gravity; any additional sources are assumed to satisfy the null energy condition. The decisive conflict is local: the sign required for strictly negative null stress also fixes a negative transverse contribution to the scalar perturbation energy.

For this analysis only, adopt natural units $c = \hbar = 1$, signature $(-,+,+,+)$, and $\kappa = 8\pi G$. The action and kinetic invariant are $S = \int d^4x\,\sqrt{-g}\,[R/(2\kappa) + P(X,\phi)]$ and $X = -g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi/2$, with P smooth and its derivatives finite on the background. Taking ϕ to have energy units makes X and P have units of energy to the fourth power, P_X dimensionless, and P_{XX} have units of inverse energy to the fourth power.

The necessary stability gate requires a physical scalar mode with a nondegenerate Lorentzian principal symbol, a common spacelike initial surface with gravity, and positive high-frequency kinetic and spatial-gradient energies relative to the gravitational sector; it is not a sufficient test of nonlinear stability or quantum consistency.

The source requirement entering this test is strict negative null stress, not merely negative normal energy. For the target metric $ds^2 = -dt^2 + (dx - b\,dt)^2 + dy^2 + dz^2$, define $n = \partial_t + b\partial_x$, $e_x = \partial_x$, $k_\pm = n \pm e_x$, and $Q = (\partial_y b)^2 + (\partial_z b)^2$. The curvature contraction gives $T(k_+,k_+) + T(k_-,k_-) = -2Q/\kappa$, including time-dependent profiles, so at least one null contraction is strictly negative wherever $Q > 0$. This is the geometric requirement that the scalar must meet, independently of how its initial data are prepared.

Write $u_\mu = \partial_\mu\phi$, $A = P_X$, and $B = P_{XX}$. Variation of the action gives $T_{\mu\nu} = Au_\mu u_\nu + Pg_{\mu\nu}$ and the scalar equation $\mathcal{E} \equiv \nabla_\mu(A\nabla^\mu\phi) + P_\phi = 0$. For every metric-null vector k^μ, the contraction is therefore $T_{\mu\nu}k^\mu k^\nu = A(k^\mu u_\mu)^2$: strict negativity requires $A < 0$ and a nonzero contraction, while the term proportional to the metric cannot change its sign. The identity $\nabla_\mu T^{\mu\nu} = \mathcal{E}\nabla^\nu\phi$ also ensures covariant stress conservation on scalar solutions.

The same derivatives of P determine the scalar's propagation and perturbation energy. Expanding about a smooth background solution as $\phi = \bar{\phi} + \pi$, and integrating the mixed field–derivative term by parts, gives the fixed-metric quadratic Lagrangian $\mathcal{L}_2 = -Z^{\mu\nu}\nabla_\mu\pi\nabla_\nu\pi/2 + U\pi^2/2$, where $Z^{\mu\nu} = Ag^{\mu\nu} - Bu^\mu u^\nu$ and $U = P_{\phi\phi} + \nabla_\mu(P_{X\phi}u^\mu)$, evaluated on the background. Here $Z^{\mu\nu}$ is also the principal coefficient of the scalar equation, and U has units of energy squared.

Freezing the coefficients supplies a necessary short-wavelength test: lower-derivative terms and background variation cannot reverse its leading derivative signs in a controlled high-frequency regime.

To expose those signs without assuming a particular background gradient, choose a local metric-orthonormal frame adapted to an arbitrary metric-timelike normal. Let $q = \partial_t\bar{\phi}$, $s_i = \partial_i\bar{\phi}$, and $K = A + Bq^2$, where the dot below denotes differentiation along the chosen local time direction. For $K \neq 0$, the conjugate momentum is $p = K\dot{\pi} - Bq(\mathbf{s}\cdot\boldsymbol{\nabla}\pi)$, and the quadratic Hamiltonian density is

$$\mathcal{H}_2 = [p + Bq(\mathbf{s}\cdot\boldsymbol{\nabla}\pi)]^2/(2K) + [A|\boldsymbol{\nabla}\pi|^2 - B(\mathbf{s}\cdot\boldsymbol{\nabla}\pi)^2]/2 - U\pi^2/2.$$

Thus changing the slicing changes the kinetic coefficient and the mixed term, but it does not eliminate the transverse coefficient A.

Choose a perturbation with spatial gradient $\mathbf{d} = \boldsymbol{\nabla}\pi$ perpendicular to $\mathbf{s}$. Its derivative Hamiltonian reduces to $\mathcal{H}_{2,\mathrm{der}} = p^2/(2K) + A|\mathbf{d}|^2/2$, so positive momentum and transverse-gradient energies require both $K > 0$ and $A > 0$. If the required $A < 0$ holds, then $K < 0$ gives negative momentum energy, whereas $K > 0$ leaves negative transverse-gradient energy; $K = 0$ makes the chosen surface characteristic for the scalar. On the formal transverse Fourier modes with wavevector $\mathbf{k} \perp \mathbf{s}$, the

frozen equation gives $\omega^2 = (A|\mathbf{k}|^2 - U)/K$, which is negative at sufficiently large $|\mathbf{k}|$ when $K > 0$, with growth rate approaching $|\mathbf{k}|\sqrt{-A/K}$.

The interpretation of this exponential behavior depends on whether the proposed initial surface is actually a scalar Cauchy surface. For $A < 0$, the two spatial directions transverse to $\mathbf{s}$ already give two negative directions of $Z^{\mu\nu}$; if the symbol is Lorentzian, it must consequently have three negative eigenvalues and one positive eigenvalue. A scalar Cauchy normal then obeys $Z^{\mu\nu}n_\mu n_\nu > 0$, which here means $K < 0$, exposing a ghost—a physical mode with negative kinetic energy relative to the ordinary gravitational sector.

Formal exponential solutions found with $K > 0$ can instead diagnose an inadmissible time slicing, rather than an instability evolved from valid scalar Cauchy data; neither interpretation restores the required positive-energy common evolution.

The gradient classifications make clear why hyperbolicity alone is insufficient. For a timelike scalar gradient, $X > 0$, use its rest frame and define $F = A + 2XB$; then $Z^{\mu\nu} = \mathrm{diag}(-F, A, A, A)$ and $\mathcal{H}_2 = p^2/(2F) + A|\boldsymbol{\nabla}\pi|^2/2 - U\pi^2/2$. Positive high-frequency energy requires $F > 0$ and $A > 0$, with sound speed squared $c_s^2 = A/F$. For the required $A < 0$, choosing $F > 0$ produces a definite rather than Lorentzian symbol and negative squared transverse frequencies, while $F < 0$ restores a Lorentzian symbol only with negative kinetic energy. Thus the hyperbolicity condition $AF > 0$ permits an overall wrong-sign branch and cannot substitute for the energy test.

For a spacelike scalar gradient, $X < 0$, align the local spatial axis with $u_\mu = (0, \sqrt{-2X}, 0, 0)$. The principal symbol becomes $Z^{\mu\nu} = \mathrm{diag}(-A, F, A, A)$, and the derivative Hamiltonian is

$$p^2/(2A) + F(\partial_x\pi)^2/2 + A[(\partial_y\pi)^2 + (\partial_z\pi)^2]/2.$$

Positive energy again requires $A > 0$ and $F > 0$; altering B changes the longitudinal coefficient but cannot repair the negative kinetic coefficient when $A < 0$. With $AF < 0$ the symbol has two eigenvalues of each sign rather than Lorentzian signature, while $A < 0$ and $F < 0$ give a Lorentzian but wrong-sign wave operator.

A nonzero null gradient does not provide an exception. In a local frame with $u_\mu = (q, q, 0, 0)$ and $X = 0$, the determinant is $\det Z = -A^4$, so finite B and $A \neq 0$ retain a Lorentzian symbol, but the general transverse Hamiltonian still fails the positive-energy test when $A < 0$. For $A > 0$, a local boost can make $|q|$ sufficiently small to obtain positive kinetic and spatial-gradient coefficients, but the null stress is then nonnegative. At zero background gradient, $u_\mu = 0$, all null stresses vanish and $Z^{\mu\nu} = Ag^{\mu\nu}$; negative A is simply the wrong-sign scalar, while $A = 0$ removes the quadratic derivative term.

Degenerate boundaries must be separated from regular propagating scalars rather than declared universally inconsistent. For non-null gradients, $\det Z = -A^3F$: at $A = 0$ the null stress vanishes and the principal operator loses rank, while at $F = 0$ it also becomes degenerate. In the timelike-gradient rest frame, $A = 0$ leaves only a quadratic time derivative and no spatial restoring term, whereas $F = 0$ leaves a spatial constraint rather than a scalar wave equation; for spacelike gradients, the corresponding limits remove either the time derivative or the longitudinal principal derivative. For a nonzero null gradient with $A = 0$, $Z^{\mu\nu} = -Bu^\mu u^\nu$ has rank one when $B \neq 0$ and rank zero otherwise.

None of these cases supplies the stipulated nondegenerate wave cone.

A concrete constrained boundary is $P = -\mu^2\sqrt{2X} - V(\phi)$ on $X > 0$, for which $A = -\mu^2/\sqrt{2X} < 0$ but $F = 0$. It can violate the null energy condition, yet its scalar principal part is a spatial constraint in the gradient rest frame, so it lies outside the regular propagating-mode criterion. Its predictivity and stability require a separate coupled constraint and elliptic–hyperbolic analysis, including boundary conditions; failure of the present gate is not a proof that every such formulation is incoherent. Higher-derivative terms likewise change the specified action and require a new test, while smooth passage through $A = 0$ cannot remove the sign obstruction within any open region of strict negative null stress.

For regular minimal coupling, the scalar principal-part test remains a necessary local gate for the coupled Einstein–scalar system in the controlled high-frequency regime. Applied as an effective field theory, it requires a nonempty wavelength window short compared with background variation but still below the theory's cutoff; without that window, the conclusion is restricted to the exact classical model rather than unknown ultraviolet dynamics. The result therefore changes the source search, not the kinematic definition of a shortcut: changing preparation cannot make a regular positive-energy single $P(X, \phi)$ scalar support this exact target.

Further progress requires a changed action, a changed target, or an explicitly analyzed constrained formulation, rather than treating nonlinear dependence on the first-derivative invariant as a stable source mechanism.

7 Axial completeness, tidal curvature, and averaged energy

Affine propagation imposes a distinct test on a prescribed shortcut geometry: local negative null stress, a negative integral along a complete ray, and a finite-affine tidal endpoint are different findings. Consider the eternal constant-speed metric $ds^2 = -(dx^0)^2 + [dx - \beta f(r)dx^0]^2 + dy^2 + dz^2$ on $M = \mathbb{R}^4$, where $x^0 = ct_{\mathrm{phys}}$, $\xi = x - \beta x^0$, and $r^2 = \xi^2 + y^2 + z^2$. The signature is $(-,+,+,+)$, and the smooth spherical profile equals one for $r \le R$ and zero for $r \ge R + \Delta$, with $R, \Delta > 0$. The analysis is classical, with zero cosmological constant and required stress tensor $T_{ab} = CG_{ab}$, where $C = c^4/(8\pi G)$; quantum and test-ray backreaction are not included.

The axial equations make completeness accessible before any averaged-energy interpretation is assigned. Write $a(\xi) = 1 - \beta + \beta f(|\xi|)$ and $w = \beta(1 - f) = 1 - a$, so the comoving metric is $ds^2 = -(dx^0)^2 + (d\xi + w\, dx^0)^2 + dy^2 + dz^2$. Reflection symmetry preserves axial motion, and its future-directed forward null tangent is $\ell = \omega(\partial_0 + a\partial_\xi)$, corresponding to $\omega(n + e_x)$ in the unit-normal frame, with $\omega = dx^0/d\lambda > 0$. Here λ has units of length, ω is dimensionless, and ∂_0 denotes differentiation at fixed comoving coordinates.

The stationary Killing field $K = \partial_0$ gives $E = -g(K, \ell) = \omega a$, while the affine equations give $d\omega/d\lambda = -\omega^2\, da/d\xi$ and $d\xi/d\lambda = E$.

For the restricted subluminal case $0 < \beta < 1$, assume additionally that the nontrivial profile is monotonically decreasing through its wall and satisfies $0 \le f \le 1$. Then $1 - \beta \le a \le 1$, so a never vanishes. Exterior normalization $\omega = 1$ fixes $E = 1 - \beta$, and consequently $1 - \beta \le \omega \le 1$. Since ξ evolves affinely with a nonzero constant derivative and the frequency remains bounded, the forward axial rays extend over a two-sided infinite affine domain; their stress contraction is supported only

in the wall crossings.

Completeness permits the averaged null-energy condition to be tested using $I_\gamma = \int_{-\infty}^{+\infty} T_{ab}\ell^a\ell^b\,d\lambda$. For this subluminal family, the axial contraction and the two symmetric wall crossings give

$$I_+ = 4C\beta(1-\beta)\int_R^{R+\Delta} f'(r)/\{r[1-\beta+\beta f(r)]^2\}\,dr < 0,$$

where f' denotes the radial derivative. The integral is finite because its denominator is bounded away from zero, $R > 0$, and the profile derivative has compact support; its strict negativity follows from the nontrivial monotone decrease.

This is a complete-geodesic averaged-null-energy violation under the stated normalization and profile assumptions, not a result that can be transferred to the superluminal regime by substituting a larger value of β.

For the superluminal analysis, fix $\beta > 1$ and an explicit smooth profile rather than assuming that every compact wall has the same endpoint structure. Define $\eta(s) = \exp(-1/s)$ for $s > 0$ and $\eta(s) = 0$ for $s \le 0$, and set $s = (r-R)/\Delta$ and $f(r) = \eta(1-s)/[\eta(s)+\eta(1-s)]$. This profile is C^∞, is constant in the core and exterior, and decreases strictly within the wall. The constant core also ensures smoothness of the radial construction at the origin.

There is exactly one $h \in (R, R+\Delta)$ satisfying $f(h) = 1 - 1/\beta$, so the forward axial coefficient vanishes precisely at $\xi = -h$ and $\xi = +h$. In the wall, $f'(r) = -f(1-f)[s^{-2} + (1-s)^{-2}]/\Delta < 0$; therefore, with $\kappa = -\beta f'(h) > 0$, the zero slopes are $a'(-h) = \kappa$ and $a'(+h) = -\kappa$. Thus a is positive between the zeros and negative outside them, and both zeros are simple. These statements identify axial characteristic separatrices without requiring the full spherical surface $r = h$ to be a global event horizon.

For $E \neq 0$, the first integral $d\xi/d\lambda = E$ divides the forward rays into three sectors, with the sign of E fixed by $\omega > 0$. In $\xi < -h$, one has $E < 0$: the future runs to $\xi = -\infty$ at infinite affine parameter, whereas the past approaches $\xi = -h$ at finite affine parameter. In $-h < \xi < h$, one has $E > 0$, and the ray travels from $-h$ to $+h$ over the finite affine interval $2h/E$. In $\xi > h$, one again has $E < 0$: the past extends to $\xi = +\infty$ at infinite affine parameter, while the future approaches $\xi = +h$ at finite affine parameter. The exterior ends are genuinely affine-complete because $a = 1-\beta$ is constant there.

The finite-affine boundaries are not finite-coordinate-time points at which these rays can simply cross into another sector. Near a zero ξ_h, let $q = a'(\xi_h) = \pm\kappa$; integrating $dx^0/d\xi = 1/a$ gives $x^0 = q^{-1}\ln(|\xi - \xi_h|/\Delta) + O(1)$. Every approach to $-h$ therefore occurs as $x^0 \to -\infty$, and every approach to $+h$ occurs as $x^0 \to +\infty$. The resulting rays are maximal in the specified manifold because their finite-affine limits have no finite-coordinate-time endpoint in M.

The zero-energy generators must be treated separately because division by a would omit them. A nonzero future null tangent with $E = 0$ lies at one of the zeros, with $\xi = \xi_h$ and $y = z = 0$ for all real coordinate times. Its frequency satisfies $\omega = 1/[q(\lambda - \lambda_c)]$, and integration gives $\lambda - \lambda_c = \exp[q(x^0 - x_0^0)]/(q\omega_0)$ for $\omega_0 > 0$. At $\xi = -h$, the affine domain is (λ_c, ∞), so the generator is past incomplete and future complete; at $\xi = +h$, the domain is $(-\infty, \lambda_c)$, giving the opposite classification.

Together with the three nonzero-energy sectors, these generators establish that every maximal forward axial ray is incomplete in at least one affine direction, even when its orbit covers every finite coordinate time.

A divergent frequency alone would not settle whether the affected endpoint admits a regular extension, so the decisive calculation uses parallelly propagated transverse curvature. With curvature convention $R(X,Y)Z = \nabla_X\nabla_Y Z - \nabla_Y\nabla_X Z - \nabla_{[X,Y]}Z$, the vanishing first transverse metric derivatives on the axis give $R_{AyBy} = -\frac{1}{2}\partial_y^2 g_{AB}$ for longitudinal indices $A, B \in \{0, \xi\}$. Contracting with $\ell^A = \omega(1, a)$ and using $w + a = 1$ yields $R(\ell, e_y, \ell, e_y) = \beta f'(r)\omega^2/r$, with an identical expression in the other transverse direction.

The unit vectors $e_y = \partial_y$ and $e_z = \partial_z$ are parallelly transported along the axial ray, so these are curvature components in a parallel screen frame, not merely components in a rapidly changing observer frame. Their sum gives $T_{ab}\ell^a\ell^b = 2C\beta f'(r)\omega^2/r$, understood as its smooth zero limit in the constant core.

For a nonzero-energy ray approaching a finite-affine endpoint λ_h, one has $\xi - \xi_h = E(\lambda - \lambda_h)$ and $a = qE(\lambda - \lambda_h) + O((\lambda - \lambda_h)^2)$. Hence $\omega \sim 1/[q(\lambda - \lambda_h)]$, and the transverse curvature and null stress behave as $R(\ell, e_y, \ell, e_y) \sim -1/[\kappa h(\lambda - \lambda_h)^2]$ and $T(\ell, \ell) \sim -2C/[\kappa h(\lambda - \lambda_h)^2]$. On either zero-energy generator, the corresponding expressions are exact with λ_h replaced by λ_c. A nondegenerate C^2 Lorentzian extension continuing the affine geodesic through such an endpoint would have continuous curvature with finite contractions against its continued tangent and parallel screen vectors, contradicting this divergence.

The result excludes that regular continuation through the affected endpoints; it neither proves inextendibility everywhere nor classifies lower-regularity extensions.

The endpoint obstruction must nevertheless be distinguished from the regular geometry at a zero at finite coordinate time. The longitudinal metric block has determinant -1 even when $a = 0$, and a different null family can cross those locations. Bounded profile derivatives keep normal-frame curvature and scalar polynomial curvature invariants bounded throughout M, but do not bound the parallelly propagated curvature produced by the unbounded affine frequency. An extension based only on the induced axial two-dimensional metric would therefore miss the decisive four-dimensional transverse tidal terms.

The explicit profile also supplies a geometric timescale without introducing a particle-production model. For $\beta = 2$, its midpoint gives $h = R + \Delta/2$, $f'(h) = -2/\Delta$, and $\kappa = 4/\Delta$. The front generator's affine frequency grows as $\exp(\kappa x^0)$, corresponding to physical e-folding time $\Delta/(4c)$. This is a classical blueshift scale, not a computed quantum emission rate or a proof of quantum instability.

The averaged-energy interpretation now differs sharply from the subluminal result. On each maximal superluminal forward ray, the stress contraction is nonpositive and its integral diverges negatively at every finite-affine endpoint, but none of these rays meets the stipulated two-sided completeness requirement. These are divergent integrals on incomplete rays, not complete-geodesic averaged-null-energy violations. Joining sectors across a zero cannot supply the missing complete ray, because the corresponding asymptotic endpoints fail the regular C^2 continuation test.

The backward axial family provides a complete-ray control with the opposite integral sign. Define

$b(\xi) = 1 + \beta - \beta f(|\xi|) = 2 - a$, so $1 \le b \le 1 + \beta$, and write $\ell_- = \omega(\partial_0 - b\partial_\xi)$. Its Killing constant is $E_- = \omega b > 0$, and $d\xi/d\lambda = -E_-$; the positive upper and lower frequency bounds then give a two-sided infinite affine domain. The axial contraction is $T(\ell_-, \ell_-) = -2C\beta f'(r)\omega^2/r$, yielding

$$I_- = -4C\beta E_- \int_R^{R+\Delta} f'(r)/\{r[1 + \beta - \beta f(r)]^2\}\, dr > 0.$$

This integral is finite, and exterior normalization $\omega = 1$ fixes $E_- = 1 + \beta$.

With length-valued affine parameter, these averaged integrals have units $\mathrm{J\,m^{-2}}$; positive affine rescaling changes their magnitude but not their sign or convergence classification. The complete backward result does not establish positivity on all complete geodesics, and the axial classification leaves complete nonaxial integrals undetermined. What changes in the central argument is therefore precise: the specified eternal superluminal target cannot be repaired by treating its affected forward endpoints as removable regular horizons, while its forward negative divergences cannot be used as a complete-ray averaged-energy exclusion.

Any finite-duration replacement must establish its own affine completeness and tidal behavior rather than inherit either a regularity claim or a prohibition from this eternal model.

8 A wider-cone alternative and the limits of comparison

A preferred-time, wider-cone field provides a distinct comparison because it changes the propagation structure rather than the geometry carrying an ordinary timelike traveler. In the geometric construction, the carrier remains inside the local metric light cone; the travel advantage concerns a comparison between geometries. The alternative instead retains Minkowski geometry for photons and ordinary matter while assigning a separate scalar field a propagation cone wider than the photon cone. Its superluminality is therefore local relative to the photon sector, not the transport of an ordinary craft along a geometrically shortened route.

The assumptions available for this comparison are a physical global time t, positive free-field kinetic energy, a characteristic propagation speed $c_s > c$, and a nonnegative squared-frequency parameter $\omega_0^2 \ge 0$. Here c is the photon speed, c_s has units of metres per second, and ω_0 has units of inverse seconds. Future-directed propagation is required to increase the same preferred time. The supplied analysis supports mathematical coherence at the free-field level under these assumptions; the comparison here does not depend on reconstructing an action or reproducing that free-field proof.

The causal role of the preferred time is substantive rather than a coordinate convention. If every permitted propagation segment advances the same global time, a sequence of such segments cannot return to its initial event. This conditional ordering explains how a wider signal cone can be considered without simultaneously granting arbitrary superluminal channels with independently chosen future orientations. It does not establish causal consistency after interactions are added: the common forward-time restriction must remain valid for the complete coupled system, rather than only for the isolated scalar sector.

The principal missing step is an interacting source–detector theory. A free propagation sector does not establish that ordinary matter can controllably emit the field, encode information in it, or

register its arrival. Those capabilities require specified couplings and a stable coupled evolution; the positive-energy properties of the uncoupled field cannot substitute for that calculation. Conservation must likewise be established for the interacting matter and field system, including their exchanges, and no gravitational completion has been demonstrated.

Even a successful source–detector construction would establish signaling rather than payload transport. Moving material requires a further account of payload coupling, its dynamics during transit, and conservation in the resulting interacting theory. No such material-transport sector is supplied. The wider-cone comparison therefore cannot serve as a mechanism for realizing the prescribed geometric corridor, nor can its free-field consistency be counted as evidence for a working travel system.

Empirical compatibility is also unresolved. Preferred-frame effects, additional radiation channels, and radiative corrections are identified as subjects for calculation and observational comparison, not as quantified predictions or established exclusions. No observational dataset or original domain-literature assessment supports numerical preferred-frame limits, a resolution of contested quantum-instability claims, or a claim of literature novelty here. Consequently, the comparison supplies neither empirical acceptance nor empirical rejection of an interacting wider-cone framework.

The alternative changes the scope of the central argument: a requirement on the Einstein source of a prescribed shortcut is not, by itself, a restriction on every possible propagation framework. It replaces the geometric sourcing question with different obligations concerning preferred-time dynamics, interactions, gravity, and material coupling. The resulting distinction is between a conditionally coherent free signal sector and a transport theory—not between two demonstrated ways of achieving superluminal travel.

9 A conditional corridor and decisive research gates

The proposed corridor is a research target, not a transport device. Its prescribed geometry separates preparation from carriage, so a source could be predeployed or arrive from outside. Neither option has been shown to work. The decisive question is whether the geometric shortcut can survive source dynamics, propagation, quantum effects, and operational control in one consistent model.

Gate 1: source dynamics. A candidate must reproduce the required density, momentum flux, and spatial stresses from a specified action and complete admissible initial data, while satisfying the coupled gravity–matter equations. Defining stress from the Einstein tensor does not construct such a source. The canonical scalar and the regular, minimally coupled single-scalar $P(X, \phi)$ class examined here fail under the stated positive-energy and common-Cauchy-slicing assumptions. A new candidate must change the action, target geometry, or those assumptions and justify the replacement; changing preparation alone does not remove a local source incompatibility.

Complete vacuum data also cannot yield the prescribed pulse in an autonomous theory with unique Cauchy development.

Gate 2: finite-duration propagation. The eternal constant-speed geometry exhibits a restricted axial-ray obstruction, but that result cannot simply be transferred to a switched pulse. The proposed test is to calculate affine-frequency evolution and parallelly propagated curvature

through a finite episode, including whether the relevant spatial velocity gradient has a sufficiently bounded time integral. That criterion remains unproved. Even a frequency bound would leave affine completeness, nonaxial rays, complete-ray energy integrals, and tidal exposure to be checked separately. A negative integral on an incomplete ray is not a global averaged-null-energy theorem.

Gate 3: quantum backreaction. For a specified field theory and state, one must fix boundary and renormalization conditions, calculate stress and fluctuations on the finite-duration configuration, and then show either a self-consistent coupled solution or a controlled backreaction bound. The classical negative-stress and blueshift calculations do not establish a universal quantum prohibition or quantum safety. A fixed-background calculation is therefore an intermediate screen, not the end of this gate.

Gate 4: preparation and control. A proposed system must show which source regions can receive control signals during formation, transit, switching, and shutdown, especially at the leading wall. Predeployment changes the architecture but still needs admissible data and an executable control law. Perturbative and nonlinear stability of the coupled system must also be assessed. The temporal function proved for one prescribed corridor does not settle the causality of multiple corridors or repeated trips.

Gate 5: empirical comparison. Only a sufficiently specified interacting theory can be tested against observations and the domain literature. No such dataset or original literature assessment is supplied here. The wider-cone alternative additionally needs interacting source, detector, and gravitational sectors before its feasibility can be compared at the same depth. The next discriminating tasks are to identify a source that escapes the stated exclusions without sacrificing well-posed evolution, and to test whether finite switching removes the eternal propagation obstruction.

Passing either task alone would not establish a workable drive; failing one would reject that particular source–target combination, not every possible shortcut framework.

10 Limitations

- The source selection is bounded and omits substantial underlying notes. It supports a scoped synthesis, not an exhaustive review or independent reconstruction of every proof.
- The conditional geometry and scalar exclusions concern specified metrics, source classes, and classical assumptions. They do not exclude every possible spacetime shortcut or replacement theory.
- The superluminal endpoint obstruction concerns the specified eternal geometry and forward axial family. It neither proves a finite-duration obstruction nor settles averaged energy on all complete geodesics.
- The spherical energy estimate is a chosen-slice integral, not an asymptotic mass or an engineering energy budget. The supplied preview does not justify a numerical device specification.
- The alternative framework is available only as a reported free-field comparison; its omitted equations prevent an equally detailed mathematical assessment.




Figure 2: The physical-feasibility chain remains open. The chosen geometry requires negative energy and null-energy violation under Einstein gravity; the examined canonical and regular single-scalar sources fail the needed conditions. Unexamined sources, finite-duration propagation, quantum behavior, stability, and control remain gates rather than achieved engineering steps.

Source-linked claims: C14, C15, C16, C33, C35, C38, C39; evidence links appear in the claim notes.

- No measured observation, completed simulation, specified quantum-state backreaction calculation, or inspected domain-literature review establishes physical viability or observational compatibility.

11 Open obligations

- Supply a matter or replacement-gravity action and solve its coupled constraints and evolution for a source matching the required density, pressures, and momentum fluxes. Establish material conservation and a common well-posed evolution rather than relying on geometric conservation alone.
- Specify complete nonvacuum initial data, incoming preparation, or boundary infrastructure. Demonstrate formation without invoking spontaneous evolution from complete Minkowski vacuum data.
- Complete the finite-duration affine calculation proposed in C19, including all relevant ray families and switching regions. Determine whether the eternal blueshift pathology persists or is avoided under explicit bounds.
- Determine averaged null-energy integrals on complete nonaxial geodesics and applicable finite-duration rays; do not use incomplete forward-ray integrals as substitutes.
- Choose quantum fields and states, calculate renormalized stress and fluctuations, identify applicable quantum energy bounds, and solve or bound self-consistent backreaction.

- Establish coupled perturbative and nonlinear stability, tidal exposure, leading-wall control, shutdown, and safe arrival. Distinguish externally prepared corridors from a self-contained craft.
- Analyze interacting and repeated configurations for causal consistency; the single-metric temporal function is insufficient for arbitrary networks.
- For the preferred-time alternative, provide the omitted free-field equations and construct stable source, detector, matter, and gravity interactions with explicit conservation laws and payload coupling.
- Inspect actual observational records and original physical literature before assigning preferred-frame limits, empirical compatibility, contested quantum-instability conclusions, or novelty.
- Define quantitative acceptance and rejection thresholds only after a concrete source theory, quantum state, operational protocol, and observational comparison have been specified.

12 Conclusions

The prescribed finite-duration geometry supplies a conditional counterexample to the assertion that an exterior-clock shortcut must involve locally superluminal carrier motion or a closed causal curve. It does not supply the physical mechanism required to realize that geometry.

The source analysis is restrictive rather than constructive: the target requires strict null-energy violation, while the canonical scalar and the stated regular positive-energy single-scalar extension cannot provide it. The eternal superluminal model also has a genuine axial endpoint pathology that cannot be repaired by treating the affected endpoints as removable coordinate horizons.

Neither a universal prohibition nor a working effective-superluminal transport framework is established. The supported outcome is a conditional corridor concept and a staged research program, with finite-duration propagation, replacement-source dynamics, quantum backreaction, control, and empirical compatibility serving as decisive gates.

A Claim and source notes

These notes identify the saved evidence for claims used in the article. The complete claim register and reasoning are retained in the paper record.

C11 For the specified eternal constant-speed metric with $\beta > 1$ and the explicit smooth monotone spherical profile, every maximal future-directed forward axial null geodesic is affinely incomplete in at least one direction. Its finite-affine endpoint has divergent parallelly propagated tidal curvature and cannot be continued through that endpoint by a regular C^2 extension. *Status:* established. *Evidence:* [Research session 1, artifact 6]

C12 For the same eternal superluminal target, the backward axial null family is complete and has a finite positive averaged null-energy integral. The incomplete forward family does not furnish a complete-geodesic averaged-null-energy test. *Status:* established. *Evidence:* [Research session 1, artifact 6]

C13 A preferred-time wider-cone free-field model provides a formally different comparison with geometric shortcuts, but no interacting source–detector, gravity, or material-transport sector has been established for it. *Status:* limitation. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 2]

C14 The prescribed geometry and its required Einstein tensor do not establish a preparable material source, stable coupled evolution, or operational transport mechanism. *Status:* limitation. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 5], [Research session 1, artifact 6]

C15 Quantum energy constraints, renormalized stress, fluctuations, and self-consistent backreaction remain undetermined for a specified field and state on a finite-duration shortcut. *Status:* unresolved. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 2], [Research session 1, artifact 6]

C16 Admissible formation, switching, leading-wall control, shutdown, tidal safety, nonlinear stability, and causal consistency of combined transport configurations remain unresolved. *Status:* unresolved. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 4], [Research session 1, artifact 5]

C17 Compatibility with existing observations, quantitative preferred-frame limits, contested quantum-instability literature, and literature novelty are not assessed by the supplied evidence. *Status:* limitation. *Evidence:* [Research session 1], [Research session 1, artifact 4], [Research session 1, artifact 5]

C18 A finite-duration, possibly predeployed transport corridor is a conditional conceptual design whose advancement requires successive source, propagation, quantum, control, and observational gates; it is not a demonstrated drive. *Status:* conjecture. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 4], [Research session 1, artifact 5], [Research session 1, artifact 6]

C19 An explicitly finite-duration propagation calculation is proposed to test whether a finite integrated spatial velocity gradient bounds affine frequency and avoids the eternal geometry's endpoint pathology. *Status:* conjecture. *Evidence:* [Research session 1, artifact 6]

C10 For $0 < \beta < 1$ and a smooth monotone spherical profile, the complete forward axial null family has a strictly negative averaged null-energy integral under the stated exterior normalization. *Status:* established. *Evidence:* [Research session 1, artifact 5]