Exact Fibonacci Corrections for Nonconsecutive Subset Sums Modulo Three

SIGMA AI-assisted research synthesis

Needs review: TeX verification is incomplete. Review the archived proofreading diagnostics before relying on the typesetting.

AI-generated research synthesis · SIGMA

Abstract

How many subsets of an initial interval contain no consecutive integers and have sum divisible by three? We derive an exact formula for this count by separating the total number of admissible subsets from a smaller residue correction. A weighted subset recurrence and a cube-root filter reduce the problem to three explicit Fibonacci formulas, one for each index class modulo three. These identities yield a homogeneous order-eight recurrence with integer coefficients, eight sufficient initial values, and the exact value $a_{12} = 128$. They also explain the limiting proportion: the total count grows on the Fibonacci scale, whereas the correction grows on the scale of a Fibonacci sequence indexed at one third of the original index.

An explicit exponential bound proves that a_n/F_{n+2} tends to $1/3$. The argument establishes these conclusions symbolically, rather than inferring a recurrence from finitely many counts.

About this paper

This paper was generated by SIGMA, an AI-assisted research system, from the project's recorded investigations, findings, and supporting evidence. The analysis and conclusions reflect an automated synthesis of that research. Although SIGMA incorporates evidence tracking, critical review, and consistency checks, the resulting paper has not necessarily undergone independent verification or peer review. Readers should evaluate its arguments, supporting evidence, and stated limitations before relying on its conclusions. Research style: Distinguish proved statements from conjectures, failed arguments, and open proof obligations. State assumptions and explain bridges between logical steps, across any mathematical branch.

Evidence coverage limitation

The evidence index omitted 2809 material saved research field fragment(s) across 7 source(s). These fragments were not reviewed for this paper. Interpretations that could depend on omitted material remain provisional; the complete frozen source snapshot and omission ledger are retained with the generation record. *Claims discussed:* C1, C2, C3, C4, C5, C6, C9

1 Introduction

For every integer $n \geq 0$, let a_n count subsets $S \subseteq \{1, \ldots, n\}$ that contain no consecutive integers and satisfy $\sum_{i \in S} i \equiv 0 \pmod 3$. The empty subset is included, so $a_0 = 1$. The question is to determine a constant-coefficient integer recurrence valid from a specified starting index, give sufficient initial values, compute a_{12}, and prove the limiting proportion relative to the total admissible count.

Write $F_0 = 0$, $F_1 = 1$, and $F_{k+2} = F_{k+1} + F_k$ for $k \geq 0$. The total number of subsets satisfying only the nonconsecutiveness condition is F_{n+2}. The central quantity is therefore the exact imbalance $c_n = 3a_n - F_{n+2}$: determining this correction answers both the finite counting question and the asymptotic question without assuming that residues are equally frequent.

Claims discussed: C1, C2, C4

2 Definitions and proof basis

The analysis represents admissible subsets by their sum-weighted polynomial and partitions them according to whether the final position is selected. Evaluation at a primitive cube root of unity then isolates sums divisible by three. A complete induction across the three index phases establishes the weighted evaluations, including their initial triple; finite values serve as arithmetic checks rather than evidence for a universal pattern.

The exact correction formulas are combined with the Fibonacci recurrence through commuting polynomial shift operators. A complementary formal-power-series elimination supplies the rational generating function and checks the recurrence's starting index. Elementary Fibonacci bounds compare the correction with the total count and give an explicit all-index error estimate; the additional minimal-order assertion is separated from these conclusions because the bounded excerpts do not expose its complete coprimality calculation.

Claims discussed: C1, C2, C3, C4, C5, C6, C7, C8, C9, C10

3 Admissible subsets and the residue filter

A weighted enumerator separates the Fibonacci count of nonconsecutive subsets from the correction imposed by divisibility of their sums by three. For an integer $n \geq 0$, call a subset $S \subseteq \{1, \ldots, n\}$ admissible if it contains no two consecutive integers, and let a_n count the admissible subsets for which $3 \mid \sum_{i \in S} i$. The ground set is empty when $n = 0$; the empty subset is admissible and has sum zero, so it contributes to every a_n.

Write $F_0 = 0$, $F_1 = 1$, and $F_{k+2} = F_{k+1} + F_k$ for every integer $k \geq 0$. The main theorem, whose proof is developed below, is that the counts satisfy the integer-coefficient recurrence $a_{n+8} = a_{n+7} + a_{n+6} + a_{n+5} - a_{n+4} - a_{n+3} + a_{n+2} - a_{n+1} - a_n$ for every integer $n \geq 0$, with sufficient initial values $(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7) = (1, 1, 1, 2, 3, 5, 8, 12)$. In particular, $a_{12} = 128$.

The theorem also gives the quantitative limiting statement $0 \leq \frac{a_n}{F_{n+2}} - \frac{1}{3} \leq \frac{2}{3}\varphi^{-2n/3} \qquad (n \geq 0)$,

where $\varphi = (1+\sqrt{5})/2$. Consequently, $a_n/F_{n+2} \to 1/3$ as $n \to \infty$. We first establish the exact counting identity from which the recurrence and the estimate will follow.

For an indeterminate q, define the polynomial $P_n(q) = \sum_{S \text{ admissible}} q^{\sum_{i \in S} i}$, where the sum ranges over admissible subsets of $\{1, \ldots, n\}$. Each subset contributes one monomial, so the coefficient of a given power counts the subsets having that sum. The empty subset contributes the constant term, and the initial polynomials are $P_0(q) = 1$ and $P_1(q) = 1 + q$.

For $n \geq 2$, partition the admissible subsets according to whether they contain n. Those omitting n are precisely the admissible subsets of $\{1, \ldots, n-1\}$, and therefore contribute $P_{n-1}(q)$. Those containing n must omit $n-1$; deleting n leaves an admissible subset of $\{1, \ldots, n-2\}$. Conversely, adjoining n to any admissible subset of this shorter interval creates no consecutive pair, because every remaining element is at most $n-2$.

The latter construction is reversible and increases the subset sum by exactly n, so its contribution is $q^n P_{n-2}(q)$. The two disjoint classes thus give the polynomial identity $P_n(q) = P_{n-1}(q) + q^n P_{n-2}(q) \qquad (n \geq 2)$. At $q = 1$, every admissible subset has weight one, and this identity becomes the Fibonacci recurrence with initial values $P_0(1) = 1$ and $P_1(1) = 2$. Hence $P_n(1) = F_{n+2}$ for every $n \geq 0$, establishing the total count used to normalize a_n.

To impose the divisibility condition, fix a primitive cube root of unity $\omega \in \mathbb{C}$, so that $\omega^3 = 1$, $\omega \neq 1$, and $1 + \omega + \omega^2 = 0$. For every integer s, the expression $1 + \omega^s + \omega^{2s}$ equals 3 if $3 \mid s$. If $s \equiv 1 \pmod 3$, it equals $1 + \omega + \omega^2 = 0$; if $s \equiv 2 \pmod 3$, it equals $1 + \omega^2 + \omega = 0$. Thus $(1 + \omega^s + \omega^{2s})/3$ is exactly the indicator that s is divisible by three.

Apply this indicator to the sum of each admissible subset and add over all such subsets. Since the sums are finite, the defining polynomial gives $3a_n = P_n(1) + P_n(\omega) + P_n(\omega^2) = F_{n+2} + P_n(\omega) + P_n(\omega^2) \qquad (n \geq 0)$. This filter selects all subset sums divisible by three, rather than a single exponent of $P_n(q)$, and includes the zero sum of the empty subset.

Define the correction by $c_n = P_n(\omega) + P_n(\omega^2)$. The filter proves the exact decomposition $3a_n = F_{n+2} + c_n$, and therefore identifies c_n with the integer $3a_n - F_{n+2}$. The counting problem is now reduced to determining the two primitive-root evaluations: the Fibonacci term accounts for the total number of admissible subsets, while their sum accounts exactly for the departure of a_n from one third of that total.

Claims discussed: C1, C2, C5, C6, C9

4 Three-phase Fibonacci formulas

The primitive-root contributions admit exact Fibonacci formulas on each index class modulo three, turning the residue filter into an explicit correction to the total count. Let ω be a primitive complex cube root of unity, so $\omega^3 = 1$ and $1 + \omega + \omega^2 = 0$, and put $p_n = P_n(\omega)$ for the weighted subset enumerator defined above. Its recurrence becomes $p_n = p_{n-1} + \omega^n p_{n-2}$ for every integer $n \geq 2$; the multiplier therefore passes successively through 1, ω, and ω^2 as the index advances through a complete triple.

We claim that, for every integer $m \geq 0$, $P_{3m}(\omega) = F_{m+1} + \omega F_m, \qquad P_{3m+1}(\omega) =$

$(1+\omega)F_{m+1}$, $P_{3m+2}(\omega) = F_m$. We prove these identities simultaneously by induction on m, advancing all three phases at each step. The initial polynomials give $p_0 = 1$, $p_1 = 1+\omega$, and $p_2 = 1+\omega+\omega^2 = 0$. These are precisely the claimed values at $m = 0$, since $F_0 = 0$ and $F_1 = F_2 = 1$.

Suppose the three identities hold for some integer $m \geq 0$. At the first index of the next triple, the multiplier is $\omega^{3m+3} = 1$, so $p_{3m+3} = p_{3m+2} + p_{3m+1} = F_m + (1+\omega)F_{m+1} = F_{m+2} + \omega F_{m+1}$. The last equality uses $F_{m+2} = F_{m+1} + F_m$, and establishes the first identity with m replaced by $m+1$.

At the next index, the multiplier is $\omega^{3m+4} = \omega$. Substituting the value just obtained and the induction hypothesis yields $p_{3m+4} = p_{3m+3} + \omega p_{3m+2} = F_{m+2} + \omega(F_{m+1} + F_m) = (1+\omega)F_{m+2}$. Thus the second identity also advances, again by the Fibonacci recurrence.

The final phase uses $\omega^{3m+5} = \omega^2$, and here the cube-root cancellation is decisive: $p_{3m+5} = p_{3m+4} + \omega^2 p_{3m+3} = (1+\omega+\omega^2)F_{m+2} + \omega^3 F_{m+1} = F_{m+1}$. This is the third identity for $m+1$. The base triple and these three successive deductions establish the formulas at every nonnegative index.

The same proof applies to the other primitive root, ω^2: replacing ω by ω^2 gives $P_{3m}(\omega^2) = F_{m+1} + \omega^2 F_m$, $P_{3m+1}(\omega^2) = (1+\omega^2)F_{m+1}$, and $P_{3m+2}(\omega^2) = F_m$. Define the integer correction by $c_n = 3a_n - F_{n+2}$. The residue-filter identity and $P_n(1) = F_{n+2}$ identify it as $c_n = P_n(\omega) + P_n(\omega^2)$, so adding the paired formulas and using $\omega + \omega^2 = -1$ gives $c_{3m} = 2F_{m+1} - F_m$, $c_{3m+1} = F_{m+1}$, $c_{3m+2} = 2F_m$ $(m \geq 0)$.

These formulas also give a constant-coefficient recurrence for the correction itself. For each fixed $r \in \{0,1,2\}$, let $b_m = c_{3m+r}$ for integers $m \geq 0$; the displayed expressions make b_m a fixed linear combination of F_m and F_{m+1}. Since both Fibonacci sequences satisfy the same recurrence in m, their linear combination satisfies $b_{m+2} = b_{m+1} + b_m$ for every $m \geq 0$. Writing each nonnegative integer uniquely as $n = 3m + r$ therefore yields $c_{n+6} = c_{n+3} + c_n$ $(n \geq 0)$.

The exact decomposition $3a_n = F_{n+2} + c_n$ thus separates two Fibonacci structures with different index increments: the total count follows the ordinary Fibonacci recurrence, while the residue correction follows it along each subsequence of indices spaced three apart. The correction is consequently not an unspecified remainder in the counting formula, but an explicitly determined sequence with its own recurrence.

Claims discussed: C3, C4

5 The integer recurrence and exact counts

The exact Fibonacci decomposition yields a single constant-coefficient recurrence for the counting sequence, with integer coefficients and validity from the first nonnegative index. Write $f_n = F_{n+2}$ and retain the correction $c_n = 3a_n - f_n$, so that $3a_n = f_n + c_n$. The identities established above give $f_{n+2} = f_{n+1} + f_n$ and $c_{n+6} = c_{n+3} + c_n$ for every integer $n \geq 0$; these two recurrences supply the factors of the required recurrence.

For a sequence $u = (u_n)_{n\geq 0}$, define the forward-shift operator E by $(Eu)_n = u_{n+1}$. Put $L(E) = E^2 - E - 1$ and $H(E) = E^6 - E^3 - 1$. To say that such an operator annihilates a sequence means that its application gives the zero sequence, so the component identities are precisely $L(E)f = 0$

and $H(E)c = 0$. Since both operators are polynomials in the same forward shift, they commute, and hence $L(E)H(E)(f + c) = H(E)L(E)f + L(E)H(E)c = 0$. Thus their product annihilates $3a$ and, by linearity, also a.

Expanding the product gives $(E^2-E-1)(E^6-E^3-1) = E^8-E^7-E^6-E^5+E^4+E^3-E^2+E+1$. Consequently, for every integer $n \geq 0$, the desired counts satisfy $a_{n+8} = a_{n+7}+a_{n+6}+a_{n+5}-a_{n+4}-a_{n+3}+a_{n+2}-a_{n+1}-a_n$. The index range is preserved in this argument: both component identities hold at every nonnegative index, and composing forward shifts uses only later indices. In particular, no negative-index extension or eventual-recurrence qualification is needed.

The exact correction formulas give $(c_0, \ldots, c_7) = (2, 1, 0, 1, 1, 2, 3, 2)$, while the corresponding Fibonacci values are $(F_2, \ldots, F_9) = (1, 2, 3, 5, 8, 13, 21, 34)$. Applying $a_n = (F_{n+2} + c_n)/3$ gives the eight starting counts: $\begin{array}{c|rrrrrrrr} n&0&1&2&3&4&5&6&7\\ \hline a_n&1&1&1&2&3&5&8&12 \end{array}$ Because the coefficient of a_{n+8} is one, the recurrence determines each next term uniquely from the preceding eight terms. These initial values are therefore sufficient to specify the entire counting sequence.

At its first application, the recurrence gives $a_8 = 12+8+5-3-2+1-1-1 = 19$. This checks the signs and the starting index by exact substitution; the universal validity of the recurrence comes from the annihilator argument, not from this finite calculation.

For the requested count, the correction formula at $12 = 3 \cdot 4$ gives $c_{12} = 2F_5 - F_4 = 10 - 3 = 7$. Since $F_{14} = 377$, the exact decomposition yields $a_{12} = \frac{F_{14}+c_{12}}{3} = \frac{377+7}{3} = 128$. The decomposition therefore supplies both an exact evaluation and a finite integer recurrence determining all later counts, rather than merely a pattern among the initial values.

Claims discussed: C5, C6

6 The rational generating function and the order question

The rational generating function explains why the recurrence combines a Fibonacci factor with a factor involving powers divisible by three. This explanation comes directly from the weighted counting recurrence, rather than from matching a finite list of counts. It also separates two logically distinct questions: whether an order-eight recurrence holds from its first possible instance, and whether any recurrence of smaller order can hold, even eventually.

Fix a primitive cube root q, so that $q^3 = 1$ and $1 + q + q^2 = 0$, and work in the formal power-series ring $\mathbb{Q}(q)[[x]]$. Recall that $P_n(q)$ is the weighted enumerator of admissible subsets, with $P_0(q) = 1$, $P_1(q) = 1+q$, and $P_n(q) = P_{n-1}(q)+q^n P_{n-2}(q)$ for $n \geq 2$. Define $G(x) = \sum_{n\geq 0} P_n(q)x^n$. Summing the recurrence over $n \geq 2$ gives $G(x) - 1 - (1 + q)x = x(G(x) - 1) + q^2x^2G(qx)$, and therefore $(1 - x)G(x) = 1 + qx + q^2x^2G(qx)$. The initial contributions are explicit, so no negative-index convention is needed.

Replacing x by qx and then by q^2x produces the other two equations: $(1 - qx)G(qx) = 1 + q^2x + qx^2G(q^2x)$ and $(1-q^2x)G(q^2x) = 1+x+x^2G(x)$. Thus the three substitutions close: after the third equation, the unknown series is again $G(x)$. Successive substitution into the first equation, followed by multiplication by $(1 - qx)(1 - q^2x)$, yields $\bigl((1-x)(1-qx)(1-q^2x)-x^6\bigr)G(x)=(1+qx)(1-

qx)(1-q^2x)+q^2x^2(1+q^2x)(1-q^2x)+x^4(1+x).$

The numerator in this elimination is essential, since it retains the initial terms as well as the recurrence structure. Its three contributions expand respectively to $1-q^2x-q^2x^2+qx^3$, $q^2x^2-x^4$, and x^4+x^5; the terms of degrees two and four cancel. Since $(1-x)(1-qx)(1-q^2x)=1-x^3$, the resulting identity is $G(x)=\frac{1-q^2x+qx^3+x^5}{1-x^3-x^6}$. Every denominator used has constant term one and is therefore invertible in the formal power-series ring; analytic convergence plays no role.

Let $b_n=P_n(q)+P_n(q^2)$ and $B(x)=\sum_{n\geq 0}b_nx^n$. Adding the preceding rational identity to its counterpart with q replaced by q^2, and using $q+q^2=-1$, gives $B(x)=\frac{2+x-x^3+2x^5}{1-x^3-x^6}$. For the total count, the Fibonacci recurrence and its first two terms give $\sum_{n\geq 0}F_{n+2}x^n=(1+x)/(1-x-x^2)$. Consequently, the residue-filter identity $3a_n=F_{n+2}+b_n$ implies $3A(x)=\frac{1+x}{1-x-x^2}+\frac{2+x-x^3+2x^5}{1-x^3-x^6}$, $\qquad A(x)=\sum_{n\geq 0}a_nx^n$.

Combining these fractions proves the formal identity $A(x)=\frac{N(x)}{D(x)}=\frac{1-x^2-x^3+x^5-x^6-x^7}{(1-x-x^2)(1-x^3-x^6)}$. The two denominator factors thus have separate origins: $1-x-x^2$ comes from the unrestricted admissible count, whereas $1-x^3-x^6$ comes from the primitive-root contribution. Expanding their product gives $D(x)=1-x-x^2-x^3+x^4+x^5-x^6+x^7+x^8$. Because $\deg N=7$, the identity $D(x)A(x)=N(x)$ implies $[x^{n+8}]D(x)A(x)=0$ for every $n\geq 0$, which is precisely $a_{n+8}-a_{n+7}-a_{n+6}-a_{n+5}+a_{n+4}+a_{n+3}-a_{n+2}+a_{n+1}+a_n=0$. In particular, the numerator establishes the starting index, not merely eventual validity.

Minimality requires a further statement that coefficient extraction alone does not supply: the numerator and denominator must be coprime over $\mathbb{Q}$. To isolate its role, suppose an eventual homogeneous constant-coefficient recurrence of order d has the normalized form $a_m=c_1a_{m-1}+\cdots+c_da_{m-d}$, with coefficients in $\mathbb{Q}$. For $R(x)=1-c_1x-\cdots-c_dx^d$, that recurrence makes $R(x)A(x)$ a polynomial, because all sufficiently high coefficients vanish. Substituting $A=N/D$ then gives $D\mid RN$; if $\gcd(N,D)=1$, it follows that $D\mid R$, hence $d\geq\deg D=8$.

The additional assertion that the displayed fraction is reduced is kept separate here from the established generating-function identity; the exact outstanding condition for the minimal-order conclusion is $\gcd(N,D)=1$ in $\mathbb{Q}[x]$. No cancellation analysis is needed for the recurrence already proved, since multiplying by the displayed denominator remains valid whether or not cancellation is possible. The generating function therefore identifies the structural source of the order-eight recurrence and certifies its full starting range, while making explicit the additional algebraic condition that would turn existence into minimality.

Claims discussed: C7, C8, C10

7 Exponential decay of the relative correction

The exact Fibonacci correction yields a quantitative limiting proportion: the excess over one third is nonnegative for every index and decays exponentially after normalization by the total count. Recall that $c_n=3a_n-F_{n+2}$ for every integer $n\geq 0$. We compare an upper bound for this correction, involving Fibonacci numbers indexed near $n/3$, with a lower bound for the total F_{n+2}.

For every integer $m\geq 0$, the established formulas are $c_{3m}=2F_{m+1}-F_m$, $c_{3m+1}=F_{m+1}$, and

$c_{3m+2} = 2F_m$. Since $0 \le F_m \le F_{m+1}$, each expression is nonnegative and at most $2F_{m+1}$; in particular, the subtraction in the first formula cannot produce a negative correction. Writing an arbitrary index as $n = 3m + r$, where $r \in \{0, 1, 2\}$, therefore gives the uniform estimate $0 \le c_n \le 2F_{\lfloor n/3 \rfloor + 1} \qquad (n \ge 0)$. This includes $c_0 = 2$ and the vanishing correction $c_2 = 0$.

Let $\varphi = (1 + \sqrt{5})/2$, so that $\varphi > 1$ and $\varphi^2 = \varphi + 1$. The elementary upper bound needed here is $F_{k+1} \le \varphi^k$ for every integer $k \ge 0$. Its two base cases are $F_1 = 1 = \varphi^0$ and $F_2 = 1 \le \varphi$. If it holds at two consecutive indices k and $k + 1$, then the Fibonacci recurrence gives $F_{k+3} = F_{k+2} + F_{k+1} \le \varphi^{k+1} + \varphi^k = \varphi^{k+2}$, completing the induction.

A separate induction gives $F_{n+2} \ge \varphi^n$ for every integer $n \ge 0$. Here the base cases are $F_2 = 1 = \varphi^0$ and $F_3 = 2 \ge \varphi$. Given the bound at indices n and $n+1$, the recurrence yields $F_{n+4} = F_{n+3} + F_{n+2} \ge \varphi^{n+1} + \varphi^n = \varphi^{n+2}$. Consequently, the correction and the total satisfy $c_n \le 2\varphi^{\lfloor n/3 \rfloor}$ and $F_{n+2} \ge \varphi^n$, respectively.

Because $F_{n+2} > 0$, division of the exact identity $3a_n = F_{n+2} + c_n$ now gives $0 \le \frac{a_n}{F_{n+2}} - \frac{1}{3} = \frac{c_n}{3F_{n+2}} \le \frac{2}{3}\varphi^{\lfloor n/3 \rfloor - n} \le \frac{2}{3}\varphi^{-2n/3} \qquad (n \ge 0)$. The last inequality uses $\lfloor n/3 \rfloor \le n/3$ and $\varphi > 1$. At $n = 0$, both the normalized excess and the final upper bound equal $2/3$, so the estimate also covers the empty ground set.

The upper bound tends to zero as $n \to \infty$, proving $\lim_{n\to\infty} a_n/F_{n+2} = 1/3$. Thus the divisible-by-three proportion converges to one third while remaining at least one third at every finite index. The mechanism is the separation of Fibonacci scales: the correction is controlled at the shorter index $\lfloor n/3 \rfloor + 1$, whereas the total is indexed by $n + 2$. The limiting proportion therefore follows from the exact correction identities and explicit inequalities, rather than from an assumed equidistribution of subset sums.

Claims discussed: C4, C9

8 Verification status and limitations

- The supported proof is an ordinary symbolic mathematical derivation, not a machine-checked or independently externally verified certificate; finite arithmetic checks are not presented as a universal proof.
- The bounded selection does not expose every calculation in the reported minimal-order argument. Minimality is therefore excluded from the main theorem and remains a separate reconstruction obligation.
- The contribution concerns the stated nonconsecutiveness restriction and divisibility by three. No extension to other moduli, other spacing restrictions, or literature novelty is asserted.

9 Open obligations

- To include minimality as an established theorem, recover the full coprimality calculation for $N(x) = 1-x^2-x^3+x^5-x^6-x^7$ and $D(x) = (1-x-x^2)(1-x^3-x^6)$ over $\mathbb{Q}$. The reconstruction

must include the reported modular remainders, the integral-factor lifting argument, and the treatment of degree loss under reduction.

- If minimality is retained, give the complete argument that a reduced generating-function denominator divides the denominator associated with any eventual homogeneous constant-coefficient recurrence. This is additional to, and unnecessary for, the requested recurrence, exact value, and limit.

10 Frozen obligation reconciliation

Unlisted obligations are not presumed resolved. The archived research snapshot retains complete source references and the full reconciliation.

- Counting interpretation and filter: established by current Investigator step 7a9db6d1-fc15-4ba5-9462-20eb79ab3e07, cycle 1, reasoning step 1. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Universal constant-coefficient integer recurrence, including its index range: established by the same supplied Investigator step, cycle 1, reasoning steps 2–4. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Sufficient initial values: established by the same supplied Investigator step, cycle 1, reasoning step 5. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Exact value a_12=128: established by the same supplied Investigator step, cycle 1, reasoning step 6, using the previously proved b recurrence. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Limiting proportion 1/3: established by the same supplied Investigator step, cycle 1, reasoning step 7, using the filter and residue-subsequence recurrence. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Weighted enumeration and exact extraction of the required count: established in cycle 1, Investigator step 8ebcb10b-506f-43d4-a180-8b996e290d86, reasoning steps 1–5, by the bijection, finite-product identity, residue evaluation, and roots-of-unity filter. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- An integer constant-coefficient recurrence valid for all $n \geq 0$: established in cycle 1, Investigator step 8ebcb10b-506f-43d4-a180-8b996e290d86, reasoning step 6, using the two explicitly justified annihilators and their expanded product. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Sufficient initial values: established in cycle 1, Investigator step 8ebcb10b-506f-43d4-a180-8b996e290d86, reasoning step 7, by evaluating the exact correction formula at $n=0,\ldots,7$. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*

- The exact value a_12: established in cycle 1, Investigator step 8ebcb10b-506f-43d4-a180-8b996e290d86, reasoning step 7, as (377+7)/3=128. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- The limiting density 1/3: established in cycle 1, Investigator step 8ebcb10b-506f-43d4-a180-8b996e290d86, reasoning step 8, by a uniform exponentially decaying bound on the relative correction. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Exact counting interpretation: cycle 2, Investigator step d9af25c9-cf16-4e90-955d-86c015f90b4d, reasoning steps 2–3 establish the residue recurrence, total count F_{n+2}, and correction identity. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Integer constant-coefficient recurrence for every n≥0: cycle 2, Investigator step d9af25c9-cf16-4e90-955d-86c015f90b4d, reasoning steps 4–7 establish the restricted transition identity, all three phases, and the expanded annihilator. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Sufficient initial values: cycle 2, Investigator step d9af25c9-cf16-4e90-955d-86c015f90b4d, reasoning step 8 derives eight consecutive initial values from exact residue counting. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Exact value a_12=128: cycle 2, Investigator step d9af25c9-cf16-4e90-955d-86c015f90b4d, reasoning step 8 obtains v_12=(128,126,123) and agrees with the scalar recurrence. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Limiting density 1/3: cycle 2, Investigator step d9af25c9-cf16-4e90-955d-86c015f90b4d, reasoning step 9 proves an explicit exponentially decaying error bound. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Counting interpretation and residue filter: established in current Investigator step 1afb8373-352e-49e5-90eb-7c696e27ce3c, cycle 2, reasoning step 2. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Exact rational generating function: established in the same current step, cycle 2, reasoning steps 3–5, by formal functional equations and checked polynomial elimination. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*
- Universal integer-coefficient recurrence with valid starting index: established in the same current step, cycle 2, reasoning step 6, by coefficient extraction from D A=N. *(Not listed by author; not presumed resolved; 2 saved source reference(s))*

17 further distinct obligations are retained in the archived reconciliation.

11 Conclusion

The counting sequence is determined by the order-eight integer recurrence in C5 and the eight initial values in C6. The exact requested count is $a_{12} = 128$.

The essential structural result is the explicit correction $3a_n - F_{n+2}$, whose three index phases are Fibonacci expressions on a shorter scale. Its relative contribution satisfies $0 \le a_n/F_{n+2} - 1/3 \le (2/3)\varphi^{-2n/3}$, proving the limiting proportion 1/3 without computational pattern matching.

The rational generating function explains the recurrence factors. Its reported reducedness and the resulting minimal-order assertion require fuller proof detail than the bounded selection supplies, so they do not strengthen the main theorem in this plan.

Claims discussed: C4, C5, C6, C7, C8, C9

Claim status and evidence

11.1 C1 – Established

For integers $n \ge 0$, define $P_n(q) = \sum_S q^{\sum_{i\in S} i}$ over subsets $S \subseteq \{1, \dots, n\}$ with no consecutive elements. Then $P_0(q) = 1$, $P_1(q) = 1+q$, and $P_n(q) = P_{n-1}(q)+q^n P_{n-2}(q)$ for every $n \ge 2$. In particular, $P_n(1) = F_{n+2}$. The partition is disjoint and reversible. Omitting n leaves any admissible subset on the first $n-1$ positions; selecting n forces omission of $n-1$, and removing n leaves any admissible subset on the first $n-2$ positions. Restoring n multiplies its weight by q^n. At $q = 1$, the initial values are 1 and 2, which identify the total with the shifted Fibonacci sequence. This is a combinatorial deduction, not a fitted recurrence. *Evidence:* [Project overview], [Research session 1, artifact 5], [Research session 1, artifact 6]

11.2 C2 – Established

If $\omega^3 = 1$ and $\omega \ne 1$, then $3a_n = F_{n+2} + P_n(\omega) + P_n(\omega^2)$ for every integer $n \ge 0$. For each integer subset sum s, the quantity $1+\omega^s+\omega^{2s}$ is 3 when 3 divides s and is zero otherwise. Summing this indicator identity over admissible subsets proves the formula, with C1 supplying the unweighted total. Evaluation at a root of unity is used as a residue filter, not as extraction of one polynomial coefficient. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 6]

11.3 C3 – Established

For every integer $m \ge 0$, the primitive-cube-root evaluations are $P_{3m}(\omega) = F_{m+1} + \omega F_m$, $P_{3m+1}(\omega) = (1+\omega)F_{m+1}$, and $P_{3m+2}(\omega) = F_m$. The saved argument contains a complete three-phase induction. With $p_n = P_n(\omega)$, the base triple is $p_0 = 1$, $p_1 = 1+\omega$, and $p_2 = 0$. Assuming the formulas for m, the weighted recurrence first gives $p_{3m+3} = F_{m+2} + \omega F_{m+1}$, then $p_{3m+4} = (1+\omega)F_{m+2}$, and finally $p_{3m+5} = F_{m+1}$. The last simplification uses $1+\omega+\omega^2 = 0$ and $\omega^3 = 1$. Thus the induction advances a whole triple and covers every nonnegative index. *Evidence:* [Research session 1, artifact 5], [Research session 1, artifact 2]

11.4 C4 – Established

The correction $c_n = 3a_n - F_{n+2}$ has the exact formulas $c_{3m} = 2F_{m+1} - F_m$, $c_{3m+1} = F_{m+1}$, and $c_{3m+2} = 2F_m$ for every integer $m \ge 0$. Consequently, $c_{n+6} = c_{n+3} + c_n$ for every integer $n \ge 0$. Apply C3 with both primitive cube roots and add the resulting expressions in C2. The relation $\omega + \omega^2 = -1$ produces each displayed integer correction. For each fixed remainder $r \in \{0, 1, 2\}$, the sequence c_{3m+r} is a fixed linear combination of Fibonacci sequences in m, so it satisfies the Fibonacci recurrence in that index. Translating

one increment of m into three increments of n yields the stated correction recurrence, including its boundary at $n = 0$. *Evidence:* [Research session 1, artifact 2], [Research session 1, artifact 5]

11.5 C5 – Established

For every integer $n \geq 0$, the counts satisfy $a_{n+8} = a_{n+7} + a_{n+6} + a_{n+5} - a_{n+4} - a_{n+3} + a_{n+2} - a_{n+1} - a_n$. Let E be the forward shift, defined by $(Eu)_n = u_{n+1}$. The sequence $f_n = F_{n+2}$ is annihilated by $E^2 - E - 1$, while C4 gives the annihilator $E^6 - E^3 - 1$ for the correction. Polynomial shift operators commute, so their product annihilates both summands of $3a = f + c$. Its expansion is $E^8 - E^7 - E^6 - E^5 + E^4 + E^3 - E^2 + E + 1$, which gives the recurrence. Both component identities hold for every nonnegative index, so the resulting recurrence does also. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 5], [Research session 1, artifact 6]

11.6 C6 – Established

Sufficient initial values are $(a_0, a_1, a_2, a_3, a_4, a_5, a_6, a_7) = (1, 1, 1, 2, 3, 5, 8, 12)$, and the requested exact value is $a_{12} = 128$. The first recurrence application gives $a_8 = 19$. Substitution into the exact correction formulas supplies the eight initial values; the leading coefficient in C5 is one, so these determine every later term uniquely. At $n = 12$, the saved calculation gives $c_{12} = 2F_5 - F_4 = 7$ and $F_{14} = 377$, hence $a_{12} = (377 + 7)/3 = 128$. The boundary check is $a_8 = 12 + 8 + 5 - 3 - 2 + 1 - 1 - 1 = 19$. These are exact finite arithmetic consequences, not computational executions. *Evidence:* [Research session 1, artifact 1], [Research session 1, artifact 6], [Research session 1, artifact 5]

11.7 C7 – Established

As a formal power series, $A(x) = \sum_{n \geq 0} a_n x^n$ satisfies $A(x) = (1 - x^2 - x^3 + x^5 - x^6 - x^7)/[(1 - x - x^2)(1 - x^3 - x^6)]$. For a primitive cube root q, the saved formal-series derivation starts with $G(x) = \sum_{n \geq 0} P_n(q) x^n$ and obtains $(1 - x)G(x) = 1 + qx + q^2 x^2 G(qx)$. Substitution at qx and $q^2 x$ gives two further equations; their elimination yields $(1 - x^3 - x^6)G(x) = 1 - q^2 x + qx^3 + x^5$. Adding the two primitive-root expressions gives $\sum_{n \geq 0} c_n x^n = (2 + x - x^3 + 2x^5)/(1 - x^3 - x^6)$. Combining this with $\sum_{n \geq 0} F_{n+2} x^n = (1 + x)/(1 - x - x^2)$ proves the displayed identity. All denominators have constant term one, so no analytic convergence assumption is needed. *Evidence:* [Research session 1, artifact 4], [Research session 1, artifact 6]

11.8 C8 – Unresolved

The rational expression in C7 is reported to be reduced, which would make eight the minimal order over $\mathbb{Q}$ among homogeneous constant-coefficient recurrences for a_n, even when a recurrence is required only eventually. The excerpts explicitly assert minimality and describe a modulo-two coprimality argument with an integral-factor lift. They do not expose the complete polynomial remainder calculations or the full treatment of degree loss under reduction. The mathematical implication is conditional and precise: if the numerator and denominator in C7 are coprime over $\mathbb{Q}$, the reduced degree-eight denominator must divide any eventual recurrence denominator. The unresolved status concerns reconstruction of the additional proof from this bounded selection, not a demonstrated counterexample or a gap in C5. *Evidence:* [Research session 1, artifact 4], [Research session 1, artifact 6]

11.9 C9 – Established

With $\varphi = (1 + \sqrt{5})/2$, every integer $n \geq 0$ satisfies $0 \leq a_n/F_{n+2} - 1/3 \leq (2/3)\varphi^{-2n/3}$. In particular, $a_n/F_{n+2} \to 1/3$. The three formulas in C4 give $0 \leq c_n \leq 2F_{\lfloor n/3 \rfloor + 1}$. The saved elementary Fibonacci bounds yield $c_n \leq 2\varphi^{\lfloor n/3 \rfloor}$ and $F_{n+2} \geq \varphi^n$. Dividing the exact identity $3a_n = F_{n+2} + c_n$ by the positive total count gives the stated estimate because $\lfloor n/3 \rfloor \leq n/3$. Since $\varphi > 1$, the upper bound tends to zero. The estimate includes $n = 0$, where equality holds. *Evidence:* [Research session 1, artifact 5], [Research session 1, artifact 6]

11.10 C10 – Limitation

The supplied mathematical derivations are exact symbolic arguments with finite arithmetic checks, but they have not been independently formally verified; the available evidence is a bounded selection rather than a complete inspection of every saved proof. The evidence-status statements distinguish algebraic deduction and hand arithmetic from execution, external checking, and formal verification. The supplied index also explicitly identifies omitted and truncated material. Accordingly, established claims here denote supported mathematical deductions at the ordinary reviewed-proof level; agreement among the saved accounts is not treated as an independent certificate. *Evidence:* [Research session 1, artifact 5], [Research session 1, artifact 6], [Research session 1]

Paper writing source material used

These sources informed manuscript organization and writing methods, not the scientific evidence or findings.

- Keith Conrad — Advice on Mathematical Writing (full text) (discipline): `https://kconrad.math.uconn.edu/blurbs/proofs/writingtips.pdf`
- PLOS ONE — Manuscript and Reporting Guidelines (general): `https://journals.plos.org/plosone/s/submission-guidelines`
- EASE — Guidelines for Authors of Scientific Articles (full text) (general): `https://pmc.ncbi.nlm.nih.gov/articles/PMC4130677/`

Evidence and authorship note

Snapshot SHA-256: 71b345c5dd63be6e1ba38e011d7274bb6a36dbb0e1d3aba70ea46c2968b84bae Generated by an AI author from the cited saved research snapshot. This document is not independent verification. No external bibliography was added.

Validation warnings

- Evidence indexing omitted 2809 material saved research field fragment(s). Those fragments were not reviewed for this paper; the interpretation is provisional where the omitted evidence

may matter. Consult the frozen source snapshot and omission ledger for full details.

- The source index is explicitly bounded and contains truncated excerpts. This plan does not claim exhaustive semantic review of the saved research or inspection of unavailable originals.
- The minimal-order result is asserted and favorably reviewed in the supplied excerpts, but the decisive coprimality arithmetic is not fully visible. Its unresolved label records an evidentiary reconstruction gap, not a mathematical refutation.
- No material mathematical contradiction is apparent among the displayed recurrence, correction formulas, generating function, initial values, and limiting estimate. Inconsistent operational assessment labels do not establish a mathematical failure and are not used as scientific premises.
- No computational execution, external proof checking, formal verification, or literature-based novelty assessment is supported. Publication and writing-method references are editorial guidance only, not evidence for the mathematical claims.
- The response schema has no section-table field. The outline therefore specifies a compact initial-value table within the relevant section rather than adding an unsupported JSON field. No diagram is proposed because the short algebraic dependency chain is clearer in the proof itself.
- The evidence index explicitly omitted 2812 saved field fragment(s); see the generation coverage ledger and review the complete archived snapshot.
- Structured paper indexing selected 451 source-linked text fragments from 11 archived sources without AI extraction calls. The complete frozen snapshot remains archived; this bounded index does not constitute review of every saved field.
- C1 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.
- C1 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.
- C2 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.
- C2 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.
- C3 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.
- C3 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.
- C4 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.

- C4 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.
- C5 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.
- C5 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.
- C6 is labeled established, but cited saved evidence includes unverified, incomplete, or conjectural material.
- C6 is labeled established while the frozen evidence contains counterexample or contradiction language; inspect the cited sources before accepting it.

Typesetting verification note

Needs review: TeX verification is incomplete. The authored research content and safe accepted typesetting corrections were retained. Review the saved proofreading diagnostics before relying on the layout.