

Primary Antibodies (/us/en/home/life-science/antibodies/primary-antibodies.html) › CD68 Antibodies (/antibody/primary/target/cd68)

Invitrogen

CD68 Monoclonal Antibody (KP1), eBioscience™

✓ **Advanced Verification**
(/us/en/home/life-science/antibodies/invitrogen-antibody-validation.html)

21
Published Figures

72
References

[View all \(200\) CD68 antibodies \(/antibody/primary/target/cd68\)](/antibody/primary/target/cd68)

Promotions [View available promotion\(s\)](#) ▼

Catalog # 14-0688-82

Price (USD)

123.00

Special offer

Online exclusive

Ends: 31-May-2026

~~164.00~~

Save 41.00 (25%)

100 µg ▼

— 1 + **In stock**

Check your price

Request bulk or custom format

(/content/lifetech/global/en/home/global/form-quote-rfd.html?

productType=Antibody&primarySiteId=U81&c0688-82&productDescription=CD68

Monoclonal Antibody (KP1),
eBioscience™)

Add to cart

Antibody Testing Data (5)

Published Figures (21)

Advanced Verification (1)

Aspire™ (/us/en/home/products-and-services/for-free-cid=bid_asp_sbu_r02_co_cp1438_pj-antibodies


Application ▼

FIGURE: 1 / 27

CD68 Antibody (14-0688-82) in ICC/IF

Immunofluorescence analysis of CD68 was performed using 70% confluent log phase THP-1 cells differentiated into Macrophage (M0 phase). The cells were fixed with 4% paraformaldehyde for 10 minutes, permeabilized with 0.1% Triton™ X-100 for 15 minutes, and blocked with 1% BSA for 1 hour at room temperature. The cells were labeled





with CD68 Mouse Monoclonal Antibody (KP1) (Product # 14-0688-80 (/order/genome-database/details/antibody/14-0688-80)) at 5 µg/mL in 0.1% BSA, incubated at 4 degree Celsius overnight and then labeled with Goat anti-Mouse IgG (H+L) Superclonal™ Secondary Antibody, Alexa... View More



Product Details

Applications	Tested Dilution	Publications
--------------	-----------------	--------------

Western Blot (WB)	Assay-Dependent	View 2 publications
Immunohistochemistry (IHC) (P)	-	View 32 publications
ThermoFisher SCIENTIFIC (/us/en/home.html)		
Immunohistochemistry (Paraffin) (IHC (P))	1-5 µg/mL	View 8 publications
Immunohistochemistry (PFA fixed) (IHC (PFA))	-	View 1 publication
Immunohistochemistry (Frozen) (IHC (F))	1 µg/mL	View 3 publications
Immunocytochemistry (ICC/IF)	5 µg/mL	View 14 publications
Flow Cytometry (Flow)	Assay-Dependent	View 8 publications
ELISA (ELISA)	-	View 1 publication
Immunoprecipitation (IP)	Assay-Dependent	View 1 publication
Mass Spectrometry (MS)	-	View 1 publication
Miscellaneous PubMed (Misc)	-	View 1 publication
Product Specifications		

Species Reactivity	Human	
Published studies	Human, Mouse, Non-human primate	ThermoFisher SCIENTIFIC
(/us/en/home.html)		
Host/Isotype	Mouse / IgG1, kappa	
Class	Monoclonal	
Type	Antibody	
Clone	KP1	
Conjugate	Unconjugated	View additional formats
Form	Liquid	
Concentration	0.5 mg/mL	
Amount	100 µg	
Purification	Affinity chromatography	
Storage buffer	PBS, pH 7.2	
Contains	0.09% sodium azide	
Storage conditions	4°C	
Shipping conditions	Ambient (domestic); Wet ice (international)	
RRID ⓘ	AB_11151139	

Product Specific Information

Description: The KP1 antibody reacts with human CD68 which belongs to the sialomucin family and is closely related to the family of lysosomal-associated membrane proteins (lamps) and scavenger receptor. CD68 is predominantly an intracellular protein, found mainly in the late endosomal compartment but can be detected in smaller amounts on the surface of mainly myeloid-derived cells; monocytes and macrophages, dendritic cells (and in some langerhan cells), neutrophils, basophils, mast cells, myeloid progenitor cells, and a subset of CD34+ hematopoietic bone marrow progenitor cells. The function has not been fully elucidated but based on homology



[/order/catalog/product/A88068?cid=bid_pca_frg_r01_co_cp1600_pjt12250_col120901_0wm](https://order/catalog/product/A88068?cid=bid_pca_frg_r01_co_cp1600_pjt12250_col120901_0wm)

and structure, CD68 may play a role in antigen processing or presentation of the lysosomal membrane and enzymes. Reports have also shown that CD68 is expressed on activated T cells, and about 40% of peripheral blood B-lymphocytes and 50% of all B-ALL. The KP1 antibody has also been reported to stain fibroblasts which may be due to a conserved epitope rather than fibroblasts expressing CD68.

Applications Reported: This KP1 antibody has been reported for use in flow cytometric analysis, immunoprecipitation, western blotting, immunohistochemical staining of frozen tissue sections, and immunohistochemical staining of formalin-fixed paraffin embedded tissue sections.

Applications Tested: This KP1 antibody has been tested by immunohistochemistry with both low and high pH antigen retrieval on FFPE tissue. Either antigen retrieval method can be used. This antibody can be used at less than or equal to 1 µg/mL. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Purity: Greater than 90%, as determined by SDS-PAGE.

Aggregation: Less than 10%, as determined by HPLC.

Filtration: 0.2 µm post-manufacturing filtered.

Target Information

CD68 (Macrosialin) is a 110 kDa integral membrane glycoprotein predominantly expressed on the intracellular lysosomes of monocytes and macrophages and to a lesser extent by dendritic cells and peripheral blood granulocytes. Also, CD68 could play a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. CD68 is expressed by interdigitating reticulum cells in tonsil and some histiocytic lymphoma or histiocytosis, acute myeloid leukemia (AML), and granulocytic sarcoma. Elevated expression of CD68 has been demonstrated on CD34+ cells in various human malignancies, including several Acute Myeloid Leukemia studies.




Related Products for

Immunohistochemistry



- Secondary Antibodies
- Additional Formats
- Isotype Controls
- Mounting Media IHC
- Counter Stain




Catalog # A21121

Invitrogen
Goat anti-Mouse IgG1 Cross-
Adsorbed Secondary Antibody,
Alexa Fluor™ 488
(/antibody/product/A21121)

Price (USD)	Unit Size :
259.00	500 µg

Add To Cart



Catalog # A21240

Invitrogen
Goat anti-Mouse IgG1 Cross-
Adsorbed Secondary Antibody,
Alexa Fluor™ 647
(/antibody/product/A21240)

Price (USD)	Unit Size :
216.65	500 µg
Online Exclusive	
289.00	
Save 72.35 (25%)	

Add To Cart

Bioinformatics

Protein Aliases: 110kda transmembrane glycoprotein;
CD68; CD68 antigen; Gp110; macrophage antigen CD68;
Macrosialin; scavenger receptor class D, member 1;
unnamed protein product

Gene Aliases: CD68; GP110; LAMP4; SCARD1

UniProt ID: (Human) P34810

(<http://www.uniprot.org/uniprot/P34810>)

Entrez Gene ID: (Human) 968

(<http://www.ncbi.nlm.nih.gov/gene?term=968>)

Function(s)



()

Process(es)



()

Performance Guarantee

If an Invitrogen™ antibody doesn't perform as described on our website or datasheet, we'll replace the product at no cost to you, or provide you with a credit for a future purchase.*

Learn more

(<https://www.thermofisher.com/us/en/home/life-science/antibodies/antibody-performance-guarantee.html>)



We're here to help

Get expert recommendations for common problems or connect directly with an on staff expert for technical assistance related to applications, equipment and general product use.

Contact tech support



(/us/en/home.html)