

NMP Liver Composition of the perfusion solution*

Component	Manufacturer/Distributor	Volume (mL)
Oxygen carrier		
Red blood cells (hematocrit 60%)	---	840
Gelofusine 4%	B Braun, Melsungen, Germany	500
Albumin 20%	---	250
Total parenteral nutrition (TPN N14G30E)	Hospital pharmacy	20
Addamel (trace elements)	Fresenius Kabi, Bad Homburg vor der Höhe, Germany	10
Metronidazol (Flagyl) 5 mg/ml	Baxter BV, Utrecht, Netherlands	44
Sterile water	B Braun, Melsungen, Germany	335
Insulin (NovoRapid) 100 IU/ml	Novo Nordisk BV, Alphen aan den Rijn, Netherlands	1
Cernevit (Multi vitamins)	Baxter BV, Utrecht, Netherlands	2
Heparin (5000 IU/ml)	Leo Pharma, Amsterdam Netherlands	2
Cefazolin 1g /5mL	Baxter BV, Utrecht, Netherlands	2
Taurocholic acid sodium salt 0.1% (1mg/mL)	Hospital pharmacy	7,7
Sodium bicarbonate 8.4%	B Braun, Melsungen, Germany	35
KCl 1mmol/mL	B Braun, Melsungen, Germany	2
Glutathion (Tationil) 600mg/4mL	Teofarma, Pavia, Italy	14
Total volume		
RBC-solution		2064,7

* Composition was designed to provide a fluid with physiological colloid osmotic pressure and osmolality. Abbreviations: HBOC; hemoglobin-based oxygen carrier, RBC; red blood cells. * The HBOC used in this study contained bovine polymerized hemoglobin dissolved in Ringer's lactate solution. # The packed red blood cells were derived from the local blood bank, and were, according to national protocol, dissolved in SAGM (saline, adenine, glucose, mannitol) solution.