



Photograph your local culture, help Wikipedia and win!

Broyden's method

🌐 5 languages ▾

In numerical analysis, **Broyden's method** is a [quasi-Newton method](#) for [finding roots](#) in k variables. It was originally described by [C. G. Broyden](#) in 1965.^[1]

[Newton's method](#) for solving $\mathbf{f}(\mathbf{x}) = \mathbf{0}$ uses the [Jacobian matrix](#), \mathbf{J} , at every iteration. However, computing this Jacobian is a difficult and expensive operation. The idea behind Broyden's method is to compute the whole Jacobian at most only at the first iteration and to do rank-one updates at other iterations.

In 1979 Gay proved that when Broyden's method is applied to a linear system of size $n \times n$, it terminates in $2n$ steps,^[2] although like all quasi-Newton methods, it may not converge for nonlinear systems.

Description of the method [\[edit\]](#)

Solving single-variable equation [\[edit\]](#)

In the [secant method](#), we replace the first derivative f' at x_n with the [finite-difference](#) approximation:

$$f'(x_n) \simeq \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}},$$

and proceed similar to [Newton's method](#):

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

where n is the iteration index.

Solving a system of nonlinear equations [\[edit\]](#)

Consider a system of k nonlinear equations

$$\mathbf{f}(\mathbf{x}) = \mathbf{0},$$

where \mathbf{f} is a vector-valued function of vector \mathbf{x} :

$$\mathbf{x} = (x_1, x_2, x_3, \dots, x_k),$$

$$\mathbf{f}(\mathbf{x}) = (f_1(x_1, x_2, \dots, x_k), f_2(x_1, x_2, \dots, x_k), \dots, f_k(x_1, x_2, \dots, x_k)).$$

For such problems, Broyden gives a generalization of the one-dimensional Newton's method, replacing the derivative with the [Jacobian](#) \mathbf{J} . The Jacobian matrix is determined iteratively, based on the **secant equation** in the finite-difference approximation:

$$\mathbf{J}_n(\mathbf{x}_n - \mathbf{x}_{n-1}) \simeq \mathbf{f}(\mathbf{x}_n) - \mathbf{f}(\mathbf{x}_{n-1}),$$

where n is the iteration index. For clarity, let us define:

$$\begin{aligned}\mathbf{f}_n &= \mathbf{f}(\mathbf{x}_n), \\ \Delta \mathbf{x}_n &= \mathbf{x}_n - \mathbf{x}_{n-1}, \\ \Delta \mathbf{f}_n &= \mathbf{f}_n - \mathbf{f}_{n-1},\end{aligned}$$

so the above may be rewritten as

$$\mathbf{J}_n \Delta \mathbf{x}_n \simeq \Delta \mathbf{f}_n.$$

The above equation is [underdetermined](#) when k is greater than one. Broyden suggests using the current estimate of the Jacobian matrix \mathbf{J}_{n-1} and improving upon it by taking the solution to the secant equation that is a minimal modification to \mathbf{J}_{n-1} :

$$\mathbf{J}_n = \mathbf{J}_{n-1} + \frac{\Delta \mathbf{f}_n - \mathbf{J}_{n-1} \Delta \mathbf{x}_n}{\|\Delta \mathbf{x}_n\|^2} \Delta \mathbf{x}_n^T.$$

This minimizes the following [Frobenius norm](#):

$$\|\mathbf{J}_n - \mathbf{J}_{n-1}\|_F.$$

We may then proceed in the Newton direction:

$$\mathbf{x}_{n+1} = \mathbf{x}_n - \mathbf{J}_n^{-1} \mathbf{f}(\mathbf{x}_n).$$

Broyden also suggested using the [Sherman–Morrison formula](#) to update directly the inverse of the Jacobian matrix:

$$\mathbf{J}_n^{-1} = \mathbf{J}_{n-1}^{-1} + \frac{\Delta \mathbf{x}_n - \mathbf{J}_{n-1}^{-1} \Delta \mathbf{f}_n}{\Delta \mathbf{x}_n^T \mathbf{J}_{n-1}^{-1} \Delta \mathbf{f}_n} \Delta \mathbf{x}_n^T \mathbf{J}_{n-1}^{-1}.$$

This first method is commonly known as the "good Broyden's method".

A similar technique can be derived by using a slightly different modification to \mathbf{J}_{n-1} . This yields a second method, the so-called "bad Broyden's method" (but see^[3]):

$$\mathbf{J}_n^{-1} = \mathbf{J}_{n-1}^{-1} + \frac{\Delta \mathbf{x}_n - \mathbf{J}_{n-1}^{-1} \Delta \mathbf{f}_n}{\|\Delta \mathbf{f}_n\|^2} \Delta \mathbf{f}_n^T.$$

This minimizes a different Frobenius norm:

$$\|\mathbf{J}_n^{-1} - \mathbf{J}_{n-1}^{-1}\|_F.$$

Many other quasi-Newton schemes have been suggested in [optimization](#), where one seeks a maximum or minimum by finding the root of the first derivative ([gradient](#) in multiple dimensions). The Jacobian of the gradient is called [Hessian](#) and is symmetric, adding further constraints to its update.

The Broyden Class of Methods [\[edit \]](#)

In addition to the two methods described above, Broyden defined a whole class of related methods.^{[1]:578} In general, methods in the *Broyden class* are given in the form^{[4]:150}

$$\mathbf{J}_{k+1} = \mathbf{J}_k - \frac{\mathbf{J}_k \mathbf{s}_k \mathbf{s}_k^T \mathbf{J}_k}{\mathbf{s}_k^T \mathbf{J}_k \mathbf{s}_k} + \frac{\mathbf{y}_k \mathbf{y}_k^T}{\mathbf{y}_k^T \mathbf{s}_k} + \phi_k \left(\mathbf{s}_k^T \mathbf{J}_k \mathbf{s}_k \right) \mathbf{v}_k \mathbf{v}_k^T,$$

where $\mathbf{y}_k := \mathbf{f}(\mathbf{x}_{k+1}) - \mathbf{f}(\mathbf{x}_k)$, $\mathbf{s}_k := \mathbf{x}_{k+1} - \mathbf{x}_k$, and

$$\mathbf{v}_k = \left[\frac{\mathbf{y}_k}{\mathbf{y}_k^T \mathbf{s}_k} - \frac{\mathbf{J}_k \mathbf{s}_k}{\mathbf{s}_k^T \mathbf{J}_k \mathbf{s}_k} \right],$$

and $\phi_k \in \mathbb{R}$ for each $k = 1, 2, \dots$. The choice of ϕ_k determines the method.

Other methods in the Broyden class have been introduced by other authors.

- The [Davidon–Fletcher–Powell \(DFP\) method](#) is the only member of this class being published before the two methods defined by Broyden.^{[1]:582} For the DFP method, $\phi_k = 1$.^{[4]:150}
- Schubert's or sparse Broyden algorithm – a modification for sparse Jacobian matrices.^[5]
- Klement (2014) – uses fewer iterations to solve many equation systems.^{[6][7]}



See also [\[edit\]](#)

- [Secant method](#)
- [Newton's method](#)
- [Quasi-Newton method](#)
- [Newton's method in optimization](#)
- [Davidon–Fletcher–Powell formula](#)
- [Broyden–Fletcher–Goldfarb–Shanno \(BFGS\) method](#)

References [\[edit\]](#)

- [^] ^{[a](#)} ^{[b](#)} ^{[c](#)} Broyden, C. G. (October 1965). "A Class of Methods for Solving Nonlinear Simultaneous Equations"[↗]. *Mathematics of Computation*. American Mathematical Society. **19** (92): 577–593. doi:10.1090/S0025-5718-1965-0198670-6[↗]. JSTOR 2003941[↗].
- [^] Gay, D. M. (August 1979). "Some convergence properties of Broyden's method". *SIAM Journal on Numerical Analysis*. SIAM. **16** (4): 623–630. doi:10.1137/0716047[↗].
- [^] Kvaalen, Eric (November 1991). "A faster Broyden method". *BIT Numerical Mathematics*. SIAM. **31** (2): 369–372. doi:10.1007/BF01931297[↗].
- [^] ^{[a](#)} ^{[b](#)} Nocedal, Jorge; Wright, Stephen J. (2006). *Numerical Optimization* [↗]. Springer Series in Operations Research and Financial Engineering. Springer New York. doi:10.1007/978-0-387-40065-5[↗]. ISBN 978-0-387-30303-1.
- [^] Schubert, L. K. (1970-01-01). "Modification of a quasi-Newton method for nonlinear equations with a sparse Jacobian"[↗]. *Mathematics of Computation*. **24** (109): 27–30. doi:10.1090/S0025-5718-1970-0258276-9[↗]. ISSN 0025-5718[↗].
- [^] Klement, Jan (2014-11-23). "On Using Quasi-Newton Algorithms of the Broyden Class for Model-to-Test Correlation"[↗]. *Journal of Aerospace Technology and Management*. **6** (4): 407–414. doi:10.5028/jatm.v6i4.373[↗]. ISSN 2175-9146[↗].
- [^] "Broyden class methods – File Exchange – MATLAB Central"[↗]. *www.mathworks.com*. Retrieved 2016-02-04.

Further reading [\[edit\]](#)

-
- [Dennis, J. E.](#); [Schnabel, Robert B.](#) (1983). *Numerical Methods for Unconstrained Optimization and Nonlinear Equations*. Englewood Cliffs: Prentice Hall. pp. 168–193. [ISBN 0-13-627216-9](#).
 - Fletcher, R. (1987). *Practical Methods of Optimization*  (Second ed.). New York: John Wiley & Sons. pp. [44–79](#) . [ISBN 0-471-91547-5](#).

External links [[edit](#)]

- [Simple basic explanation: The story of the blind archer](#) 

Category: [Quasi-Newton methods](#)