Report on "Matrix Theory Reloaded: A BPS Road to Holography"

Blair, Lahnsteiner, Obers, Yan (arXiv-style preprint IFT-UAM/CSIC-24-139, NORDITA 2024-033)

9 November 2026

Scope

This report is based on a reading of the abstract, introduction, the summaries of the holographic and $T\bar{T}$ results, the flow-equation summary, and the outlook of `paper.tex`. I did not re-derive the calculations.

1 Principal achievement

The paper unifies two decoupling limits that were previously treated separately. Matrix p-brane theory (MpT) arises from a BPS decoupling limit that makes the target-space geometry non-Lorentzian. The authors show that the near-horizon limit of a Dp-brane bulk geometry is the same kind of limit, applied to the bulk. For $\mathrm{AdS}_5 \times S^5$, the near-horizon metric takes the M3T form with $(r/\ell)^2$ playing the role of the limit parameter ω. This recasts the AdS/CFT decoupling limit as a statement about asymptotic non-Lorentzian geometry and gives a systematic way to generate further holographic pairs, including ones with non-Lorentzian bulks.

The second achievement is a limit-free route to the bulk. The authors argue that the bulk geometry is an irrelevant, $T\bar{T}$-type deformation of the flat non-Lorentzian boundary geometry. They derive p-brane generalisations of $T\bar{T}$ with explicit flow equations. For free scalars the flow equation has a closed-form solution,

$$\mathcal{L}(\lambda) = \frac{1}{\lambda}\Big[1 - \sqrt{-\det\big(\eta_{\alpha\beta} + \lambda\, \partial_\alpha X^{A'} \partial_\beta X^{A'}\big)}\Big],$$

which is the Dirac–Nambu–Goto action of a gauge-fixed p-brane. Full flow equations including a gauge field are obtained for $p = 1, 2$. At $p = 1$ with vanishing field strength they reduce to the standard $T\bar{T}$ flow.

2 Principal drawback

The paper's most ambitious claims rest on geometric matching and on a conjecture, not on independent checks.

- **The DLCQ^n/DLCQ^m correspondence is a conjecture supported by consistency.** The examples reproduce known AdS/CFT limits and fit the classification of BPS limits. I did not find a case where an observable, such as a spectrum or a correlator, is computed on both the bulk and the field-theory side and compared. Without such a test, the claim that this captures holography in string theory remains a proposal.

- **The non-Lorentzian bulk geometries are not verified as solutions.** The new non-Lorentzian holographic backgrounds are obtained by introducing an extra rescaling of the length ℓ in the harmonic function by hand. The authors say they "simply view it as a trick". They also state that they have not verified that these backgrounds solve the equations of motion and torsional constraints of the MpT limit of supergravity, because that theory has not yet been written down explicitly.

- **The $T\bar{T}$-type flow equations are limited.** They are classical and bosonic, and restricted to abelian worldvolume theories. The full flow is obtained only for $p = 1, 2$. For $p = 3$ it is given only to linear order in λ, and for $p > 3$ only at zeroth order. The connection to the non-abelian DBI action, which is what matrix theory needs, is therefore not established.

3 Most promising follow-up

I compared three candidate directions on feasibility and impact.

1. **Check the non-Lorentzian bulk backgrounds against the $\omega \to \infty$ limit of type II supergravity.** Take one single-stack example, write the type II field equations in the ω-scaled variables (including the dilaton and RR rescalings the paper uses), expand in $1/\omega$, and test whether the leading terms vanish on the background. This is a finite symbolic computation that needs no new formalism at first, and it is the gap the authors identify themselves. Its outcome is informative either way. If the backgrounds pass, the non-Lorentzian holographic examples and the DLCQ framing gain real support. If they fail, the failing rescalings give a sharper criterion for which geometries are admissible. The main risk is that the expansion produces new constraints, such as the torsional constraints the authors mention, which a naive limit of the relativistic equations does not capture. Resolving that risk is essentially the formulation of MpT supergravity that the paper defers.

2. **Extend the p-brane $T\bar{T}$ flow to the non-abelian case at leading order.** This would have higher impact, since it would connect matrix theory to non-abelian DBI. Its feasibility is low, because the non-abelian D-brane action is itself known only partially, beyond the leading order in α'.

3. **Compute the isometry group of the non-Lorentzian bulk.** This is feasible and the authors flag it as a natural calculation. Its impact is more limited, since it does not by itself test whether the backgrounds are solutions or whether the correspondence holds.

Recommendation: option 1. It is the best balance of feasibility and impact. It is a concrete calculation that turns the most speculative part of the paper into a testable statement, and it directly supports the DLCQ conjecture the paper is built around.