Report on
Matrix Theory Reloaded: A BPS Road to Holography

(Blair, Lahnsteiner, Obers, Yan)

9 November 2026

Summary of the paper. The paper revisits the decoupling limit that produces matrix theory on Dp-branes, treating it as a *BPS* limit ($\omega \to \infty$, with the RR (p+1)-form tuned to cancel the brane tension) of the whole of type II string theory. The result is "matrix p-brane theory" (MpT), which lives on a non-Lorentzian, p-brane Newton–Cartan target space. The paper makes three claims: (i) applying this limit at asymptotic infinity to a Dp-brane supergravity solution induces exactly the Maldacena/IMSY near-horizon limit, and the near-horizon geometry in turn approaches flat MpT geometry asymptotically; (ii) applying two such limits to off-aligned or intersecting branes (e.g. M0T on the D4 geometry, or D1–D5) generates non-Lorentzian bulk geometries and recovers recent non-Lorentzian holography proposals. In M-theory every BPS limit is a DLCQ, which leads to the conjecture that holography is generally a DLCQ^n/DLCQ^m correspondence ($m > n$), with AdS$_5$/CFT$_4$ as DLCQ^0/DLCQ^1; (iii) undoing the limit is U-dual to the $T\bar{T}$ deformation of the non-relativistic string. The authors use this to argue that the near-horizon bulk is a background-dependent "p-brane $T\bar{T}$" deformation of flat MpT space, and they derive classical flow equations: a closed form for any p that takes free scalars to Nambu–Goto, and closed forms for $p = 1, 2$ that take scalars plus Maxwell to DBI.

1. Principal achievement

The main achievement is **conceptual unification**. The paper places the matrix theory limit, the near-horizon limit of AdS/CFT, non-Lorentzian holography and the $T\bar{T}$ deformation in one systematic framework. The key technical observation is that one asymptotic BPS limit, with $L^{7-p} \propto \omega^{(p-3)/2}$ and $R = \omega^{-1/2} r$, turns the harmonic function into $H = 1 + \omega^2(\ell/r)^{7-p}$. This is a geometric account of why the decoupling of the open string sector at infinity *is* the near-horizon limit in the bulk. It also explains why the field theory side should be thought of as coupling to a non-Lorentzian 10D geometry rather than to Minkowski space.

The real gain over the classic Maldacena/IMSY treatment is that the same rule can be *iterated*. Composing BPS limits, constrained by U-duality and the DLCQ interpretation, gives a generating procedure for holographic pairs. It reproduces the Lambert et al. non-Lorentzian dualities (e.g. the M0T limit of the D4 near-horizon geometry, dual to quantum mechanics on instanton moduli space) and organises AdS$_3$/CFT$_2$ from D1–D5 as DLCQ^0/DLCQ^2. There is also a concrete, self-contained by-product: the flow equation

$$\partial_\lambda \mathcal{L} = \frac{1}{2\lambda^2}\Big\{\mathrm{tr}(\mathbb{1} - \lambda\mathcal{T}) - (p-1)\big[\det(\mathbb{1} - \lambda\mathcal{T})\big]^{1/(p-1)} - 2\Big\},$$

which takes D free scalars to the static-gauge Dirac–Nambu–Goto action in any dimension. It reduces to $T\bar{T}$ for $p = 1$ and to the Gross et al. quantum mechanical deformation for $p = 0$. The paper also identifies a simple membrane flow for $p = 2$. These results correct the expectation, built on earlier flows such as Ferko et al.'s, that a λ-independent quadratic flow suffices for multiple scalars.

2. Principal drawback

The main weakness is that the **holographic claims are kinematic rather than dynamical**. Everything in Sections 3–5 consists of reparametrisations and limits of *classical* brane solutions and probe actions. For the standard ($n = 0$) case this mostly rewrites known limits in ω language. For the genuinely new non-Lorentzian cases, no independent evidence of a duality is given:

- The non-Lorentzian near-horizon backgrounds are not shown to solve the MpT supergravity equations or torsion constraints. The authors acknowledge this in the Outlook, and those equations are not formulated for general p.
- Neither the (super)isometry algebras of the bulk geometries nor the symmetries of the proposed boundary theories are computed and matched.
- No observable (spectrum, index, correlator, entropy) is compared across any proposed pair.
- Several of the geometries need an *ad hoc* rescaling of ℓ, effectively $N \to \infty$ with ω, or smearing. The physical meaning of these sources is left open.

As a result, the DLCQ^n/DLCQ^m conjecture is close to a classification rule: it says which limits commute and how many DLCQs each side has, but it does not make a falsifiable prediction beyond examples already proposed elsewhere. The analysis is also confined to the Poincaré patch, and for $p > 3$ it is valid only up to strong-coupling caveats. The $T\bar{T}$ part has a parallel limitation. Its flows are classical, bosonic and abelian. They are built to reach DBI and are therefore not unique, since terms that vanish along the DBI trajectory can be added. The $p = 3$ gauge-field flow is known only to $O(\lambda)$, and for $p \geq 2$ there is no factorisation or spectral argument that would make the deformation well defined quantum mechanically.

3. Most worthwhile follow-up (feasibility vs. impact)

Proposal: make the simplest DLCQ^1/DLCQ^2 example quantitative, namely the M0T limit of the D4-brane near-horizon geometry, dual to the null reduction of the 6D $(2, 0)$ theory (equivalently, quantum mechanics on instanton moduli space).

Why this example is feasible. M0T is DLCQ M-theory, so this background uplifts to $\mathrm{AdS}_7 \times S^4$ with a null identification along a boundary direction. The bulk equations are then simply 11D supergravity, with null reduction, and the paper's need for a general MpT supergravity disappears. On the field theory side, protected quantities are under good control: 5D SYM instanton partition functions, $(2, 0)$ superconformal indices, and the instanton moduli space quantum mechanics. Concrete steps:

1. Check explicitly that the 10D non-Lorentzian background of Section 3.2 (the near-horizon MpT fields at $p = 0$, $q = 4$) solves the null-reduced 11D supergravity equations, including the torsion constraints.
2. Compute its super-isometry algebra, which should be the subalgebra of $\mathfrak{osp}(8^*|4)$ commuting with the null translation, a Schrödinger-type superalgebra. Match it with the symmetry algebra of the non-relativistic boundary theory.
3. Compute the linearised Kaluza–Klein spectrum at fixed D0 charge (null momentum k) and compare it with protected operators or BPS states of the boundary theory at instanton number k, for example via the k-instanton index.
4. Use the non-Lorentzian 10D description to revisit the known pathologies of null-compactified AdS, such as the small null circle and the breakdown of the Lorentzian sugra description. This is where the paper's framework should earn its keep.

Why the impact is high. A positive result would be the first dynamical test of a top-down non-Lorentzian holographic duality. It would also turn the DLCQ^n/DLCQ^m proposal from a classification into a computational tool, with an obvious extension to the D3-in-M(-1)T and M2/ABJM analogues. A negative result, such as a symmetry or spectral mismatch, would be equally informative about where the conjecture fails. The competing directions are less balanced.

Quantising the $p = 2$ membrane flow, or applying it to ABJM, has high impact but low feasibility. Supersymmetric or non-abelian extensions of the flows are feasible but of more incremental impact.