

Additive Genetic Models in Mixed Populations

Facundo Muñoz

2026-09-10 breedR version: 0.13.0

Contents

Full diallel trial with founders from two different base populations E and J

```
## Setup
library(breedR)

## Loading required package: sp
library(ggplot2)
set.seed(123)

## Simulation parameters
n.founders <- c(E = 9, J = 9)
sigma2 <- c(E = 3, J = 2, resid = 1)

founders <- data.frame(id = c(paste0('E', 1:n.founders['E']),
                             paste0('J', 1:n.founders['J'])),
                      pop.idx = c(rep(1, n.founders['E']),
                                rep(2, n.founders['J'])),
                      BV = c(rnorm(n.founders['E'], sd = sqrt(sigma2['E'])),
                            rnorm(n.founders['J'], sd = sqrt(sigma2['J']))))

n.obs <- sum(n.founders)*150
obs.parents.idx <- matrix(sample(nrow(founders), 2*n.obs, replace = TRUE), ncol = 2)

## The 'family' is independent of the order of the parents
## While the population only takes into account the origin
## e.g. cross2fam(c('J3', 'E1')) gives 'E1:J3'
## while cross2pop(c('J3', 'E1')) gives 'EJ'
cross2fam <- function(x) paste(founders$id[sort(x)], collapse = ':')
cross2pop <- function(x) paste(names(n.founders)[founders$pop.idx[sort(x)]], collapse = '')

## Mendelian sampling term
msp <- function(x) {
  ss <- sigma2[founders$pop.idx[x]]
  s2 <- (ss[1] + ss[2])/4
  rnorm(1, sd = sqrt(s2))
}

dat <- data.frame(
  id = sum(n.founders) + seq.int(n.obs),
```

```

dad = obs.parents.idx[, 1],
mum = obs.parents.idx[, 2],
fam = as.factor(apply(obs.parents.idx, 1, cross2fam)),
sp = as.factor(apply(obs.parents.idx, 1, cross2pop)),
bv = apply(obs.parents.idx, 1,
           function(x) mean(founders$BV[x]) + msp(x)),
resid = rnorm(n.obs, sd = sqrt(sigma2['resid'])))

dat <- transform(dat,
                 y = bv + resid)

## Printing simulated setting
print(table(dat[, c('mum', 'dad')]), zero.print = "")

```

```

##      dad
## mum  1  2  3  4  5  6  7  8  9 10 11 12 13 14 15 16 17 18
##  1   8 11 12  7  7 14  8 10  9 14 11 13  7  8  8  7 11  7
##  2  12  8  5 12  5  9  5 10  7 13  7  5  5  7  8 14 11  7
##  3   6 12  5  7  8 10  6 13  5  5  7  6  6  9  9  7  2 13
##  4   6  8  8  6 11  8 11  9 11 10  9  5  8  6  9  9 10 11
##  5   7  7  8  4 13  8 14  6  6  5  6 11  7  5  3  6  8 11
##  6  10  5  9  7  6  6  5  7 14  9  5  9  4  6  2  7  9  7
##  7   8 10  7 14 14 10  8 11  5  8  9  5  9 15  6  7  7  8
##  8  10  7 11  5  5  3  3 15 10 10  7  9  7  7  9  4  8 11
##  9   6  5  9  2  3 11 11  7  7  5 11 10  8  8  9  9  5  4
## 10   6  2 10  9  7  4 11 13  6  9  7  7  6 14 14  7 13 10
## 11   4  3  9 10  7 10  5 14  9  9 12  7  8  5  9  8 11  7
## 12   9  5 10  5 10  6  6  6  8  9 11  7 10  8 10  8  6 10
## 13   3 14  8  8  5  5  9  9  5  6 10 10  7  6  8  8  8  5
## 14   5  9  8  4 12 10  8  9  8 17 12 13  4  7  8 11  9  8
## 15   6 11  6 12 14  7 12  5 17 12  6  4  8  9 12 10  3 13
## 16  13 10  9  8  9  8  8 10  9 11 10  8  4  7  6  9 10  5
## 17   7  5  7  8 13 11 10  6  6 14  9  9 14 11  9  4  9  8
## 18   9  9  8 11  9  4 12 14  9  4 11 15  5 10 10  8 12 10

```

```
str(dat)
```

```

## 'data.frame':    2700 obs. of  8 variables:
## $ id      : num  19 20 21 22 23 24 25 26 27 28 ...
## $ dad     : int  11 7 12 15 10 13 7 9 9 10 ...
## $ mum     : int  7 11 5 16 17 17 8 3 8 14 ...
## $ fam     : Factor w/ 171 levels "E1:E1","E1:E2",...: 98 98 74 163 134 155 95 42 107 131 ...
## $ sp      : Factor w/ 3 levels "EE","EJ","JJ": 2 2 2 3 3 3 1 1 1 3 ...
## $ bv      : num  2.797 0.395 0.19 1.284 0.862 ...
## $ resid   : num  -0.721 -2.585 0.271 -0.142 1.104 ...
## $ y       : num  2.076 -2.19 0.461 1.142 1.966 ...

```

There will be two independent additive-genetic variance models for the J and E populations. Each variance component will be estimated using the *pure* offspring only.

```

## Build a pedigree for the whole mixed population
## and get the kinship matrix A
ped <- build_pedigree(1:3, data = dat)
A <- pedigreeemm::getA(ped)

```

```
## 'as(<dtTMatrix>, "dtCMatrix")' is deprecated.
```

```
## Use 'as(., "CsparseMatrix")' instead.
## See help("Deprecated") and help("Matrix-deprecated").

## Build the full incidence matrix
Z <- as(dat$id, 'indMatrix')

## Give the index vector of additive-genetic random effects
## that belong to one subpopulation;
## 'E', 'J' (founders) or 'EE', 'EJ' or 'JJ' (offspring).
idx_pop <- function(x) {
  if (nchar(x) == 1) grep(x, founders$id)
  else
    match(dat$id[dat$sp == x], as.data.frame(ped)$self)
}
```

0.1 Method 1: Hybrids as an independent population

This is the easiest way to go. It works as a first approximation, but has several shortcomings.

It needs to estimate one *virtual* variance for the hybrid population which is not linked to any genetic variance in the *real world*. Moreover, we don't use the hybrid observations to learn about the two variances that really matter: σ^2_E and σ^2_J

The advantage is that it predicts the Breeding Values of the hybrid offspring. However, the accuracy may be limited by the violation of the *single population* hypothesis of the model.

```
## Avoid estimating BLUPS for which we don't have information
## Otherwise, the run takes much longer (5 hs vs 6 min in this example)

## A[idx_pop('EE'), idx_pop('JJ')] # This is null: populations are independent
Z_EE <- Z[, idx_pop('EE')]
Z_EJ <- Z[, idx_pop('EJ')]
Z_JJ <- Z[, idx_pop('JJ')]

A_EE <- A[idx_pop('EE'), idx_pop('EE')]
A_EJ <- A[idx_pop('EJ'), idx_pop('EJ')]
A_JJ <- A[idx_pop('JJ'), idx_pop('JJ')]

## Now fit a model with three additive-genetic components,
## by means of generic effects (as only one 'genetic' is allowed in breedR)

res1 <- remlf90(y ~ sp,
  generic = list(
    E = list(incidence = Z_EE,
              covariance = A_EE),
    J = list(incidence = Z_JJ,
              covariance = A_JJ),
    H = list(incidence = Z_EJ,
              covariance = A_EJ)),
  data = dat
)


## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.

summary(res1)
```

```
## Formula: y ~ 0 + sp
##   Data: dat
##   AIC  BIC logLik
## 9937 9961 -4965
##
## Parameters of special components:
##
##
## Variance components:
##           Estimated variances   S.E.
## E                2.7194 0.7185
## J                2.5068 0.7144
## H                2.7841 0.6992
## Residual          0.8589 0.3495
##
## Fixed effects:
##           value   s.e.
## sp.EE 0.23745 0.5527
## sp.EJ 0.20826 0.3955
## sp.JJ 0.21205 0.5306

PBV <- as.matrix(cbind(Z_EE, Z_JJ, Z_EJ)) %*%
  do.call('rbind', lapply(ranef(res1), function(x) cbind(PBV = x, se = attr(x, 'se'))))

ggplot(cbind(dat, PBV), aes(bv, PBV)) +
  geom_point() +
  geom_abline(intercept = 0, slope = 1, col = 'darkgray')
```



0.2 Method 2: GCA/SCA model for hybrids

Another approach is to model hybrids with GCAs and SCAs, with additional variance parameters. The additive component of the genetic variances for the E and J populations is half the population variance. But these parameters will also gather dominance and epistasis effects.

However, we don't take advantage of the known relationship between the additive components by treating them as independent parameters. Furthermore, we don't use the relationship between hybrids and pures to

learn about the original genetic variances. Nor even within hybrids, as we treat SCA as an unstructured effect, while there are both half and full siblings.

```
## We only want to apply 'dad', 'mum' and 'sca' effects to hybrids,
## and make it zero for non-hybrids. We do so by pre-multiplying by a
## diagonal indicator matrix
Ind <- diag(dat$sp == 'EJ')

## Build a specific incidence matrices for generic random effects
Z_dad <- Ind %*% as(dat$dad, 'indMatrix')
Z_mum <- Ind %*% as(dat$mum, 'indMatrix')
Z_sca <- Ind %*% as(as.numeric(dat$fam), 'indMatrix')

## The structure variances are diagonal
D <- diag(sum(n(founders)))

res2 <- remlf90(y ~ sp,
               generic = list(
                 E = list(incidence = Z_EE,
                          covariance = A_EE),
                 J = list(incidence = Z_JJ,
                          covariance = A_JJ),
                 dad = list(incidence = Z_dad,
                           covariance = D),
                 mum = list(incidence = Z_mum,
                           covariance = D),
                 sca = list(incidence = Z_sca,
                           covariance = diag(nlevels(dat$fam)))),
               data = transform(dat)
)
```

```
## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
```


```
summary(res2)
```

```
## Formula: y ~ 0 + sp
##   Data: transform(dat)
##   AIC   BIC logLik
## 10010 10045 -4999
##
## Parameters of special components:
##
## Variance components:
##           Estimated variances      S.E.
## E           0.9373800 0.2014200
## J           0.7679600 0.1731200
## dad         0.6773600 0.2457700
## mum         0.7146600 0.2585500
## sca         0.0002231 0.0002437
## Residual    2.0314000 0.0657310
##
## Fixed effects:
##           value   s.e.
## sp.EE 0.23742 0.3285
```

```
## sp.EJ 0.21363 0.2809
## sp.JJ 0.20965 0.2979

PBV <- as.matrix(cbind(Z_EE, Z_JJ, Z_dad, Z_mum, Z_sca)) %*%
  do.call('rbind', lapply(ranef(res2), function(x) cbind(PBV = x, se = attr(x, 'se'))))

ggplot(cbind(dat, PBV), aes(bv, PBV)) +
  geom_point() +
  geom_abline(intercept = 0, slope = 1, col = 'darkgray')
```



This approach does not work very well. Much of the variation has not been adequately accounted for, and ended up in the residuals. May be there is some misspecification in the model.

0.3 Method 3: grid search

The difficulty of the problem resides that we cannot estimate two variance parameters at the same time. If we split the matrix A in blocks corresponding to each subpopulation, we can write the covariance matrix for the mixture as

$$\begin{aligned} \Sigma &= \begin{bmatrix} \sigma_E^2 A_E & \mathbf{0} & \sigma_E^2 A_{E:EE} & \frac{3\sigma_E^2 + \sigma_J^2}{4} A_{E:H} & \mathbf{0} \\ \mathbf{0} & \sigma_J^2 A_J & \mathbf{0} & \frac{\sigma_E^2 + 3\sigma_J^2}{4} A_{J:H} & \sigma_J^2 A_{J:JJ} \\ \sigma_E^2 A_{E:EE} & \mathbf{0} & \sigma_E^2 A_{EE} & \frac{3\sigma_E^2 + \sigma_J^2}{4} A_{EH} & \mathbf{0} \\ \frac{3\sigma_E^2 + \sigma_J^2}{4} A_{E:H} & \frac{\sigma_E^2 + 3\sigma_J^2}{4} A_{J:H} & \frac{3\sigma_E^2 + \sigma_J^2}{4} A_{HE} & \frac{\sigma_E^2 + \sigma_J^2}{2} A_{HH} & \frac{\sigma_E^2 + 3\sigma_J^2}{4} A_{HJ} \\ \mathbf{0} & \sigma_J^2 A_{J:JJ} & \mathbf{0} & \frac{\sigma_E^2 + 3\sigma_J^2}{4} A_{JH} & \sigma_J^2 A_{JJ} \end{bmatrix} \\ &= \sigma_E^2 \begin{bmatrix} A_E & \mathbf{0} & A_{E:EE} & \frac{3+\lambda}{4} A_{E:H} & \mathbf{0} \\ \mathbf{0} & \lambda A_J & \mathbf{0} & \frac{1+3\lambda}{4} A_{J:H} & \lambda A_{J:JJ} \\ A_{E:EE} & \mathbf{0} & A_{EE} & \frac{3+\lambda}{4} A_{EH} & \mathbf{0} \\ \frac{3+\lambda}{4} A_{E:H} & \frac{1+3\lambda}{4} A_{J:H} & \frac{3+\lambda}{4} A_{HE} & \frac{1+\lambda}{2} A_{HH} & \frac{1+3\lambda}{4} A_{HJ} \\ \mathbf{0} & \lambda A_{J:JJ} & \mathbf{0} & \frac{1+3\lambda}{4} A_{JH} & \lambda A_{JJ} \end{bmatrix}, \end{aligned} \quad (1)$$

where $\lambda = \frac{\sigma_J^2}{\sigma_E^2}$.

However, if we are not interested in evaluating the parents, we can disregard the first two block rows and columns from the matrix.

Now, fit the model for several values of λ and maximize the likelihood.

```

## Setup parallel computing
# library(doParallel)
# cl <- makeCluster(2)
# registerDoParallel()
# on.exit(stopCluster(cl))

## Introduce the corresponding scaling factors
## in the relationship matrix
scale_A <- function(x) {

  ## The pure E subpopulations remains the same
  S <- A
  E.idx <- c(idx_pop('E'), idx_pop('EE'))

  ## The pure J subpopulations get multiplied by lambda
  J.idx <- c(idx_pop('J'), idx_pop('JJ'))
  S[J.idx, J.idx] <- A[J.idx, J.idx] * x

  ## The hybrids related wuth pure E get a factor of (3+lambda)/4
  S[idx_pop('EJ'), E.idx] <- A[idx_pop('EJ'), E.idx] * (3+x)/4
  S[E.idx, idx_pop('EJ')] <- A[E.idx, idx_pop('EJ')] * (3+x)/4

  ## The hybrids related wuth pure J get a factor of (1+3*lambda)/4
  S[idx_pop('EJ'), J.idx] <- A[idx_pop('EJ'), J.idx] * (1+3*x)/4
  S[J.idx, idx_pop('EJ')] <- A[J.idx, idx_pop('EJ')] * (1+3*x)/4

  ## Finally, the hybrids related with other hybrids get a factor of (1+lambda)/2
  S[idx_pop('EJ'), idx_pop('EJ')] <- A[idx_pop('EJ'), idx_pop('EJ')] * (1+x)/2

  return(S)
}

## Conditional likelihood given lambda
cond_lik <- function(x) {
  require(breedR)
  ## Conditional structure matrix
  S <- scale_A(x)

  ## Temporarily, let's use only the pure pops
  idx <- c(idx_pop('EE'), idx_pop('JJ'))

  suppressWarnings(
    res <- remlf90(y ~ sp,
      generic = list(
        E = list(incidence = Z[dat$sp != 'EJ', idx],
          covariance = S[idx, idx])),
        data = dat[dat$sp != 'EJ', ]
      )
    )
  logLik(res)
}


lambda <- seq(.3, 1.5, length.out = 5)

```

```
lik <- sapply(lambda, cond_lik) # (sequential)

## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
##
## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
##
## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
##
## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
##
## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.
##
# lik <- foreach(x = seq.int(lambda), .combine = c) %dopar% cond_lik(lambda[x])

ggplot(data.frame(lambda, lik), aes(lambda, lik)) +
  geom_line()
```



The profile is flat near its peak. With nine founders per population λ is weakly identified, and the best grid point shifts from one simulated data set to the next; the simulated value is $\lambda = 2/3$. The estimate is only as fine as the grid, which extends past 1 so that a peak above 1 would show up inside it rather than at the edge. But for some reason, when I include the hybrids, for lambdas below about 0.58 and all the variance is accounted for residual variance.

There must be something wrong somewhere. May be the simulation of the Breeding Values is not correct?

Analogously, if I include only the hybrid subpopulation, it works quite well, identifying the base variance.

```
## Take lambda maximizing the likelihood
lambda0 <- lambda[which.max(lik)]

S <- scale_A(lambda0)
```

```

## Temporarily, let's use only the pure pops
idx <- c(idx_pop('EE'), idx_pop('JJ'))

# ## Remove the founders, which I don't want to evaluate
# idx <- -(1:sum(n.founders))

res3 <- remlf90(y ~ sp,
               generic = list(
                 E = list(incidence = Z[dat$sp != 'EJ', idx],
                           covariance = S[idx, idx])),
               data = dat[dat$sp != 'EJ', ])

## Using default initial variances given by default_initial_variance()
## See ?breedR.getOption.

summary(res3)

```

```

## Formula: y ~ 0 + sp
##   Data: dat[dat$sp != "EJ", ]
##   AIC   BIC logLik
##  5072 5082  -2534
##
## Parameters of special components:
##
##
## Variance components:
##           Estimated variances   S.E.
## E                2.642 0.9102
## Residual          0.913 0.4380
##
## Fixed effects:
##           value   s.e.
## sp.EE 0.23745 0.5449
## sp.EJ 0.00000 0.0000
## sp.JJ 0.21201 0.5168

```

The best grid point is $\lambda = 0.9$, giving an estimated additive-genetic variance for the J population of 2.38, against a simulated value of 2.

```

PBV <- as.matrix(Z[dat$sp != 'EJ', idx]) %*%
  do.call('rbind', lapply(ranef(res3), function(x) cbind(PBV = x, se = attr(x, 'se'))))

ggplot(cbind(dat[dat$sp != 'EJ', ], PBV), aes(bv, PBV)) +
  geom_point() +
  geom_abline(intercept = 0, slope = 1, col = 'darkgray')

```

