


Radioactivity and dose in radioactive media


Eirik Malinen



The valley of stability




Nuclear binding energy



Radioactivity and radiation dose


- About 255 stable nuclei
- Nuclides in nature: 340
- Decay modes in radioactive nuclei
 - β^- : neutron rich
 - β^+ : neutron deficient
 - EC : neutron deficient near valley
 - α : heavy nuclei
 - Fission : very heavy nuclei

α -disintegration




- α -particle typically have 2-7 MeV energy
- ~ 4 cm range in air; 50-80 μm in water
- Becomes He-atom after slowing down

α -decay scheme




Ra-223 - therapy

- Radiopharmaceutical for bone cancer




- Ra-223 and daughters emit over 95% of the decay energy in the form of alpha radiation

β^- -disintegration



- Spectrum of electrons emitted: $T=[0, T_{\beta, \text{max}}]$
- Remember: $n \rightarrow p + e^- + \bar{\nu}_e$


β^- -decay scheme



UiO Department of Physics
University of Oslo

Oslo University Hospital

β^+ -disintegration




- Spectrum of positrons emitted
- Here: $p + 1.022 \text{ MeV} \rightarrow n + e^+ + \nu_e$

UiO Department of Physics
University of Oslo

Oslo University Hospital

β^+ -decay scheme




UiO Department of Physics
University of Oslo

Oslo University Hospital

Fluorine-18

- Irradiation of stable oxygen-18 isotope with high energy protons (18 MeV)




- The target is either normal or enriched water

UiO Department of Physics
University of Oslo

Oslo University Hospital


Fluorine-18 - PET



UiO Department of Physics
University of Oslo

Oslo University Hospital

β^-/β^+ decay spectrum




- β^- spectrum shifted due to electrostatic attraction with nucleus
- Rule of thumb: $\bar{T} = 1/3 T_{\max}$

UiO Department of Physics
University of Oslo

Oslo University Hospital

Disintegration by electron capture




- Same daughter nuclei as β^+ disintegration

UiO Department of Physics
University of Oslo

Oslo University Hospital

EC / β^+ -decay scheme




UiO Department of Physics
University of Oslo

Oslo University Hospital

Internal conversion

- γ -ray emission with energy $h\nu$ often accompanies decay
- Alternative: impart kinetic energy to atomic electron
- Energy: $T = h\nu - E_b$

Internal conversion cont'd



Radioactivity and radiation dose

- Radioactivity important in
 - Radiation protection – external and internal exposure
 - Brachytherapy
 - Nuclear medicine – PET / SPECT, Radionuclide / Radioimmunotherapy

Radioactivity

- Number of decay events dN occurring during dt is proportional to number of atoms N :

$$\frac{dN}{dt} \propto -N$$

- Decay constant λ introduced:

$$\frac{dN}{dt} = -\lambda N$$

- Simple differential equation with solution:

$$N = N_0 e^{-\lambda t}$$

Radioactivity

- λ is characteristic of each nuclide; decay probability per unit time per atom

- Rate of decay; activity is given by:

$$\left| \frac{dN}{dt} \right| = \left| \frac{d}{dt} (N_0 e^{-\lambda t}) \right| = \lambda N$$

- If nucleus has several modes of disintegration:

$$\lambda = \lambda_A + \lambda_B + \dots = \sum_i \lambda_i$$

Radioactivity cont'd

- The total activity is then

$$\lambda N = N \sum_i \lambda_i$$

- Partial activity for i'th mode of disintegration:

$$\lambda_i N = \lambda_i N_0 e^{-\lambda t}$$

- Activity given in units of Becquerel [Bq] = s⁻¹

- Earlier: 1 Ci = 3.7 × 10¹⁰ Bq

Radioactivity cont'd

- Time needed to decay to 1/e of original number of nuclei:

$$\frac{N}{N_0} = \frac{1}{e} = e^{-\lambda \tau} \Rightarrow \tau = \frac{1}{\lambda}$$

- τ : mean life, average lifetime of a nucleus.
- Probability of nucleus not having decayed:


$$p(t) = C \exp^{-\lambda t} \Rightarrow \int_0^{\infty} p(t) dt = 1 \Rightarrow C = \frac{1}{\lambda}$$

- Mean lifetime: $\tau = \int_0^{\infty} t p(t) dt = \frac{1}{\lambda} \int_0^{\infty} t e^{-\lambda t} dt = \frac{1}{\lambda}$

Radioactivity cont'd

- Half life $t_{1/2}$ defined as

$$\frac{N}{N_0} = \frac{1}{2} = e^{-\lambda t_{1/2}} \Rightarrow t_{1/2} = \frac{\ln(2)}{\lambda}$$



Parent-daughter relationships

- Initial population of parent nuclei $N_{P,0}$
- Decays into a radioactive daughter by λ_P
- Daughter activity:

$$\frac{dN_D}{dt} = \lambda_P N_P - \lambda_D N_D = \lambda_P N_{P,0} e^{-\lambda_P t} - \lambda_D N_D$$

- It can be shown that:


$$N_D = N_{P,0} \frac{\lambda_P}{\lambda_D - \lambda_P} (e^{-\lambda_P t} - e^{-\lambda_D t})$$

Parent-daughter relationships cont'd

- Daughter will have zero activity at $t=0$ and $t=\infty$. Maximum of N_D reached at:

$$\frac{dN_D}{dt} = 0 \Rightarrow -\lambda_P e^{-\lambda_P t_m} + \lambda_D e^{-\lambda_D t_m} = 0$$

$$\Rightarrow t_m = \frac{\ln(\lambda_D / \lambda_P)}{\lambda_D - \lambda_P}$$



Parent-daughter relationships cont'd


- Ratio of activities:

$$\begin{aligned} \frac{\lambda_D N_D}{\lambda_P N_P} &= \frac{\lambda_D N_{P,0} \frac{\lambda_P}{\lambda_D - \lambda_P} (e^{-\lambda_P t} - e^{-\lambda_D t})}{\lambda_P N_{P,0} e^{-\lambda_P t}} \\ &= \frac{\lambda_D}{\lambda_D - \lambda_P} (1 - e^{-(\lambda_D - \lambda_P)t}) \end{aligned}$$


- For $\lambda_D < \lambda_P$ the ratio increases with time
- For $\lambda_D > \lambda_P$ at $t \gg t_m$, the ratio is $\lambda_D / (\lambda_D - \lambda_P)$

Parent-daughter relationships cont'd

- For $\lambda_D \gg \lambda_P$ the activity ratio is unity
- Example



Parent-daughter relationships cont'd



UiO Department of Physics
University of Oslo

Oslo University Hospital

Absorbed dose in radioactive media

- Dose from α -particles: $D = n \times T$
- Dose from β 's: $D \approx n \times \bar{T}$
- Dose from γ 's; two limiting cases:
 1. In a *small* radioactive object V , CPE is obtained at least a distance $d = \langle t \rangle$ (or a few MFPs) from the boundary of V . The dose is given by KERMA.
 2. In a large radioactive object ($\gg 1/\mu$), the dose will be given by the sum of the kinetic energies emitted.

UiO Department of Physics
University of Oslo


Oslo University Hospital

Absorbed dose in radioactive media

- At intermediate radioactive objects, we use the absorbed fraction

$$AF = \frac{\gamma\text{-ray energy absorbed}}{\gamma\text{-ray energy emitted}}$$

$$= \frac{\epsilon_{dv,V}}{R_{dv,V}}$$




- Assuming exponential attenuation over r

$$AF = \frac{1}{4\pi} \iint (1 - e^{-\mu r}) \sin \theta d\theta d\varphi$$

UiO Department of Physics
University of Oslo

Oslo University Hospital

Absorbed fraction



UiO Department of Physics
University of Oslo

Oslo University Hospital