Lesson 7: Linear First-Order and Bernoulli Equations

Francisco Blanco-Silva

University of South Carolina

September 12, 2013

WHAT DO WE KNOW?

► The concepts of differential equation and initial value problem

$$F(x, y, y', \dots, y^{(n)}) = 0$$

- ► The concept of order of a differential equation.
- The concepts of general solution, particular solution and singular solution.
- ► Slope fields
- Approximations to solutions via Euler's Method and Improved Euler's Method

- ► Separable equations $y' = H_1(x)H_2(y)$
- ► Homogeneous First-Order Equations y' = H(y/x)

DEFINITION

LINEAR FIRST-ORDER EQUATIONS

The linear first-order equation has the form

$$\frac{dy}{dx} + P(x)y = Q(x)$$

DEFINITION

The linear first-order equation has the form

$$\frac{dy}{dx} + P(x)y = Q(x)$$

To solve it, we use an integrating factor (we disregard constant of integration now)

$$\rho(x) = e^{\int P(x) \, dx}$$

DEFINITION

The linear first-order equation has the form

$$\frac{dy}{dx} + P(x)y = Q(x)$$

To solve it, we use an integrating factor (we disregard constant of integration now)

$$\rho(x) = e^{\int P(x) \, dx}$$

The solution of the linear first-order equation can then be written in implicit form as

$$\rho(x)y = C + \int \rho(x)Q(x) \, dx$$

DEFINITION

The linear first-order equation has the form

$$\frac{dy}{dx} + P(x)y = Q(x)$$

To solve it, we use an integrating factor (we disregard constant of integration now)

$$\rho(x) = e^{\int P(x) \, dx}$$

The solution of the linear first-order equation can then be written in implicit form as

$$\rho(x)y = C + \int \rho(x)Q(x) \, dx$$

Why?

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

The key of the formula for the solution of these equations is the product $\rho(x)y$.

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

$$\frac{d}{dx}(\rho(x)y) = \frac{d}{dx}\rho(x)y + \rho(x)\frac{dy}{dx}$$

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

$$\frac{d}{dx}(\rho(x)y) = \frac{d}{dx}\rho(x)y + \rho(x)\frac{dy}{dx}$$
$$= P(x)\rho(x)y + \rho(x)\frac{dy}{dx}$$

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

$$\frac{d}{dx}(\rho(x)y) = \frac{d}{dx}\rho(x)y + \rho(x)\frac{dy}{dx}$$

$$= P(x)\rho(x)y + \rho(x)\frac{dy}{dx}$$

$$= \rho(x)\underbrace{\left(P(x)y + \frac{dy}{dx}\right)}_{\text{What is this?}}$$

PROOF

Why does this formula work? We start with the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Let us examine first the integration factor $\rho(x) = e^{\int P(x) dx}$. What is its derivative?

$$\frac{d}{dx}(\rho(x)) = P(x)e^{\int P(x) dx} = P(x)\rho(x)$$

$$\frac{d}{dx}(\rho(x)y) = \frac{d}{dx}\rho(x)y + \rho(x)\frac{dy}{dx}$$

$$= P(x)\rho(x)y + \rho(x)\frac{dy}{dx}$$

$$= \rho(x)\underbrace{\left(P(x)y + \frac{dy}{dx}\right)}_{\text{What is this?}} = \rho(x)Q(x)$$

PROOF

We have just discovered that

$$\frac{d}{dx}(\rho(x)y) = \rho(x)Q(x).$$

PROOF

We have just discovered that

$$\frac{d}{dx}(\rho(x)y) = \rho(x)Q(x).$$

Integrate both sides.

$$\rho(x)y = \int \rho(x)Q(x) \, dx$$

PROOF

We have just discovered that

$$\frac{d}{dx}(\rho(x)y) = \rho(x)Q(x).$$

Integrate both sides.

$$\rho(x)y = \int \rho(x)Q(x) \, dx + C$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

$$P(x) = -1 \qquad Q(x) = e^{-x}$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

Let us identify P, Q, and compute ρ :

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

We need only one more ingredient:

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

Let us identify P, Q, and compute ρ :

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

We need only one more ingredient:

$$\int \rho(x)Q(x) dx = \int e^{-x}e^{-x} dx = \int e^{-2x} dx = -\frac{1}{2}e^{-2x}$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

Let us identify P, Q, and compute ρ :

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

We need only one more ingredient:

$$\int \rho(x)Q(x)\,dx = \int e^{-x}e^{-x}\,dx = \int e^{-2x}\,dx = -\frac{1}{2}e^{-2x}$$

The solution is then

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

Let us identify P, Q, and compute ρ :

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

We need only one more ingredient:

$$\int \rho(x)Q(x) dx = \int e^{-x}e^{-x} dx = \int e^{-2x} dx = -\frac{1}{2}e^{-2x}$$

The solution is then

$$\rho(x) y = C + \int \rho(x)Q(x) dx$$
$$e^{-x}y = C - \frac{1}{2}e^{-2x}$$

EXAMPLES

Find a general solution

$$\frac{dy}{dx} - y = e^{-x}$$

Let us identify P, Q, and compute ρ :

$$P(x) = -1$$
 $Q(x) = e^{-x}$ $\int P(x) dx = -x$ $\rho(x) = e^{-x}$

We need only one more ingredient:

$$\int \rho(x)Q(x) dx = \int e^{-x}e^{-x} dx = \int e^{-2x} dx = -\frac{1}{2}e^{-2x}$$

The solution is then

$$\rho(x) y = C + \int \rho(x)Q(x) dx$$
$$e^{-x}y = C - \frac{1}{2}e^{-2x}$$
$$y = Ce^{x} - \frac{1}{2}e^{-x}$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

Let us identify *P*, *Q*. But first, we need to re-write the equation:

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

Let us identify *P*, *Q*. But first, we need to re-write the equation:

$$\frac{dy}{dx} = \frac{6x - 3xy}{x^2 + 1} = \frac{6x}{x^2 + 1} - \frac{3x}{x^2 + 1}y$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

Let us identify *P*, *Q*. But first, we need to re-write the equation:

$$\frac{dy}{dx} = \frac{6x - 3xy}{x^2 + 1} = \frac{6x}{x^2 + 1} - \frac{3x}{x^2 + 1}y$$
$$\frac{dy}{dx} + \underbrace{\frac{3x}{x^2 + 1}}_{P(x)} y = \underbrace{\frac{6x}{x^2 + 1}}_{Q(x)}$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

Let us identify *P*, *Q*. But first, we need to re-write the equation:

$$\frac{dy}{dx} = \frac{6x - 3xy}{x^2 + 1} = \frac{6x}{x^2 + 1} - \frac{3x}{x^2 + 1}y$$
$$\frac{dy}{dx} + \underbrace{\frac{3x}{x^2 + 1}}_{P(x)} y = \underbrace{\frac{6x}{x^2 + 1}}_{Q(x)}$$

Let us compute the integrating factor:

$$\int P(x) dx = \int \frac{3x}{x^2 + 1} dx = \frac{3}{2} \ln|x^2 + 1| = \frac{3}{2} \ln(x^2 + 1)$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

Let us identify *P*, *Q*. But first, we need to re-write the equation:

$$\frac{dy}{dx} = \frac{6x - 3xy}{x^2 + 1} = \frac{6x}{x^2 + 1} - \frac{3x}{x^2 + 1}y$$
$$\frac{dy}{dx} + \underbrace{\frac{3x}{x^2 + 1}}_{P(x)} y = \underbrace{\frac{6x}{x^2 + 1}}_{Q(x)}$$

Let us compute the integrating factor:

$$\int P(x) dx = \int \frac{3x}{x^2 + 1} dx = \frac{3}{2} \ln|x^2 + 1| = \frac{3}{2} \ln(x^2 + 1)$$
$$\rho(x) = e^{\frac{3}{2} \ln(x^2 + 1)} = (x^2 + 1)^{3/2}$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x)\,dx$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x)\,dx = \int (x^2+1)^{3/2} \frac{6x}{x^2+1}\,dx = \int 6x(x^2+1)^{1/2}\,dx$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x) dx = \int (x^2 + 1)^{3/2} \frac{6x}{x^2 + 1} dx = \int 6x(x^2 + 1)^{1/2} dx$$
$$= \underbrace{\int 3u^{1/2} du}_{u=x^2 + 1} = 2u^{3/2} = (x^2 + 1)^{3/2}$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x) dx = \int (x^2 + 1)^{3/2} \frac{6x}{x^2 + 1} dx = \int 6x(x^2 + 1)^{1/2} dx$$
$$= \underbrace{\int 3u^{1/2} du}_{u=x^2 + 1} = 2u^{3/2} = (x^2 + 1)^{3/2}$$

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x) dx = \int (x^2 + 1)^{3/2} \frac{6x}{x^2 + 1} dx = \int 6x(x^2 + 1)^{1/2} dx$$
$$= \underbrace{\int 3u^{1/2} du}_{u=x^2 + 1} = 2u^{3/2} = (x^2 + 1)^{3/2}$$

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$
$$(x^2 + 1)^{3/2} y = (x^2 + 1)^{3/2} + C$$

EXAMPLES

Find a general solution

$$(x^2+1)\frac{dy}{dx} = 6x - 3xy$$

The final ingredient:

$$\int \rho(x)Q(x) dx = \int (x^2 + 1)^{3/2} \frac{6x}{x^2 + 1} dx = \int 6x(x^2 + 1)^{1/2} dx$$
$$= \underbrace{\int 3u^{1/2} du}_{u=x^2 + 1} = 2u^{3/2} = (x^2 + 1)^{3/2}$$

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$
$$(x^2 + 1)^{3/2} y = (x^2 + 1)^{3/2} + C$$
$$y = 1 + C(x^2 + 1)^{-3/2}$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x)$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x) \qquad \qquad \frac{dy}{dx} \underbrace{-(1+x)}_{P(x)} y = \underbrace{1+x}_{Q(x)}$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x) \qquad \qquad \frac{dy}{dx} \underbrace{-(1+x)}_{P(x)} y = \underbrace{1+x}_{Q(x)}$$

$$\int P(x) \, dx = -\int (1+x) \, dx = -x - \frac{1}{2}x^2$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x)$$

$$\frac{dy}{dx} \underbrace{-(1+x)}_{P(x)} y = \underbrace{1+x}_{Q(x)}$$

$$\int P(x) dx = -\int (1+x) dx = -x - \frac{1}{2}x^2 \quad \rho(x) = e^{-x-x^2/2}$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x)$$

$$\frac{dy}{dx} \underbrace{-(1+x)}_{P(x)} y = \underbrace{1+x}_{Q(x)}$$

$$\int P(x) dx = -\int (1+x) dx = -x - \frac{1}{2}x^2 \quad \rho(x) = e^{-x-x^2/2}$$

$$\int \rho(x)Q(x)\,dx = \int (1+x)e^{-x-x^2/2}\,dx$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\frac{dy}{dx} = (1+x) + y(1+x) \qquad \frac{dy}{dx} \underbrace{-(1+x)}_{P(x)} y = \underbrace{1+x}_{Q(x)}$$

$$\int P(x) dx = -\int (1+x) dx = -x - \frac{1}{2}x^2 \quad \rho(x) = e^{-x-x^2/2}$$

$$\int \rho(x)Q(x) dx = \int (1+x)e^{-x-x^2/2} dx = \underbrace{\int e^{-u} du}_{u=x+x^2/2} = -e^{-u} = -e^{-x-x^2/2}$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\rho(x) y = C + \int \rho(x) Q(x) dx$$
$$e^{-x - x^2/2} y = C - e^{-x - x^2/2}$$

EXAMPLES

Find a general solution

$$y' = 1 + x + y + xy$$

$$\rho(x) y = C + \int \rho(x)Q(x) dx$$

$$e^{-x-x^2/2}y = C - e^{-x-x^2/2}$$

$$y = Ce^{x+x^2/2} - 1$$

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = y^{1-n}$.

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = y^{1-n}$.

$$y = v^{1/(1-n)}$$

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = y^{1-n}$.

$$y = v^{1/(1-n)}$$
 $\frac{dy}{dx} = \frac{1}{1-n} v^{n/(1-n)} \frac{dv}{dx}$

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = v^{1-n}$.

$$y = v^{1/(1-n)}$$
 $\frac{dy}{dx} = \frac{1}{1-n} v^{n/(1-n)} \frac{dv}{dx}$

We use this substitution in the original equation, to get rid of the y's:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

Bernoulli Equation

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = v^{1-n}$.

$$y = v^{1/(1-n)}$$
 $\frac{dy}{dx} = \frac{1}{1-n} v^{n/(1-n)} \frac{dv}{dx}$

We use this substitution in the original equation, to get rid of the y's:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^{n}$$

$$\frac{1}{1-n}v^{n/(1-n)}\frac{dv}{dx} + \overline{P}(x)v^{1/(1-n)} = \overline{Q}(x)v^{n/(1-n)}$$

DEFINITION

The Bernoulli equation is very similar to the linear equation:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^n$$

We can turn it into one of those, by performing a substitution: $v = y^{1-n}$.

$$y = v^{1/(1-n)}$$
 $\frac{dy}{dx} = \frac{1}{1-n} v^{n/(1-n)} \frac{dv}{dx}$

We use this substitution in the original equation, to get rid of the y's:

$$\frac{dy}{dx} + \overline{P}(x)y = \overline{Q}(x)y^{n}$$

$$\frac{1}{1-n}v^{n/(1-n)}\frac{dv}{dx} + \overline{P}(x)v^{1/(1-n)} = \overline{Q}(x)v^{n/(1-n)}$$

$$\frac{dv}{dx} + \underbrace{(1-n)\overline{P}(x)}_{P(x)}v = \underbrace{(1-n)\overline{Q}(x)}_{Q(x)}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is
$$\overline{P}(x) = 6/x$$
, $\overline{Q}(x) = 3$, $n = 4/3$.

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$.

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is
$$\overline{P}(x) = 6/x$$
, $\overline{Q}(x) = 3$, $n = 4/3$.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3} \qquad \qquad \frac{dy}{dx} = -3v^{-4}\frac{dv}{dx}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3} \qquad \qquad \frac{dy}{dx} = -3v^{-4}\frac{dv}{dx}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3} \qquad \qquad \frac{dy}{dx} = -3v^{-4}\frac{dv}{dx}$$

We get then

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3} \qquad \frac{dy}{dx} = -3v^{-4}\frac{dv}{dx}$$

We get then

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3} \qquad -3v^{-4}\frac{dv}{dx} + \frac{6}{x}v^{-3} = 3v^{-4}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

Rewrite the equation to realize it is a Bernoulli, and find \overline{P} , \overline{Q} , n:

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3}$$

It is $\overline{P}(x) = 6/x$, $\overline{Q}(x) = 3$, n = 4/3.

We need to apply the substitution $v = y^{1-4/3} = y^{-1/3}$. Rather than using the formula, it pays off to perform all the computations from scratch.

$$y = v^{-3} \qquad \frac{dy}{dx} = -3v^{-4}\frac{dv}{dx}$$

We get then

$$\frac{dy}{dx} + \frac{6}{x}y = 3y^{4/3} \qquad -3v^{-4}\frac{dv}{dx} + \frac{6}{x}v^{-3} = 3v^{-4} \qquad \frac{dv}{dx} - \frac{2}{x}v = -1$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

$$P(x) = -\frac{2}{x}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

$$P(x) = -\frac{2}{x} \qquad Q(x) = -1$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

$$P(x) = -\frac{2}{x}$$
 $Q(x) = -1$ $\int P(x) dx = -2 \ln|x|$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

$$P(x) = -\frac{2}{x}$$
 $Q(x) = -1$ $\int P(x) dx = -2 \ln|x|$ $\rho(x) = x^{-2}$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

$$P(x) = -\frac{2}{x} \qquad Q(x) = -1 \qquad \int P(x) \, dx = -2 \ln|x| \qquad \rho(x) = x^{-2}$$
$$\int \rho(x) Q(x) \, dx = \int -x^{-2} \, dx = x^{-1}$$

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

Let us compute all the ingredients of the formula:

$$P(x) = -\frac{2}{x} \qquad Q(x) = -1 \qquad \int P(x) \, dx = -2 \ln|x| \qquad \rho(x) = x^{-2}$$
$$\int \rho(x) Q(x) \, dx = \int -x^{-2} \, dx = x^{-1}$$

Therefore, the solution of this equation is

$$x^{-2}v = C + x^{-1}$$

Bernoulli Equation

EXAMPLES

Find a general solution

$$x\frac{dy}{dx} + 6y = 3xy^{4/3}$$

We need to solve now the linear first-order equation

$$\frac{dv}{dx} - \frac{2}{x}v = -1$$

Let us compute all the ingredients of the formula:

$$P(x) = -\frac{2}{x} \qquad Q(x) = -1 \qquad \int P(x) \, dx = -2 \ln|x| \qquad \rho(x) = x^{-2}$$
$$\int \rho(x) Q(x) \, dx = \int -x^{-2} \, dx = x^{-1}$$

Therefore, the solution of this equation is

$$x^{-2}v = C + x^{-1}$$
 $x^{-2}y^{-3} = C + x^{-1}$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

$$\frac{dy}{dx} + \underbrace{\frac{x}{1+x^2}}_{\overline{P}(x)} y = \underbrace{\frac{x^2}{1+x^2}}_{\overline{Q}(x)} y^2$$

n = 2. We use the substitution $v = y^{1-2} = y^{-1}$.

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

$$\frac{dy}{dx} + \underbrace{\frac{x}{1+x^2}}_{\overline{P}(x)} y = \underbrace{\frac{x^2}{1+x^2}}_{\overline{Q}(x)} y^2$$

n = 2. We use the substitution $v = y^{1-2} = y^{-1}$.

$$y = v^{-1} \qquad \frac{dy}{dx} = -v^{-2} \frac{dy}{dx}$$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

$$\frac{dy}{dx} + \underbrace{\frac{x}{1+x^2}}_{\overline{P}(x)} y = \underbrace{\frac{x^2}{1+x^2}}_{\overline{Q}(x)} y^2$$

n = 2. We use the substitution $v = y^{1-2} = y^{-1}$.

$$y = v^{-1} \qquad \qquad \frac{dy}{dx} = -v^{-2} \frac{dv}{dx}$$

We get the linear equation

$$\frac{dy}{dx} + \frac{x}{1+x^2}y = \frac{x^2}{1+x^2}y^2$$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

$$\frac{dy}{dx} + \underbrace{\frac{x}{1+x^2}}_{\overline{P}(x)} y = \underbrace{\frac{x^2}{1+x^2}}_{\overline{Q}(x)} y^2$$

n = 2. We use the substitution $v = y^{1-2} = y^{-1}$.

$$y = v^{-1} \qquad \qquad \frac{dy}{dx} = -v^{-2}\frac{dv}{dx}$$

We get the linear equation

$$\frac{dy}{dx} + \frac{x}{1+x^2}y = \frac{x^2}{1+x^2}y^2$$
$$-v^{-2}\frac{dv}{dx} + \frac{x}{1+x^2}v^{-1} = \frac{x^2}{1+x^2}v^{-2}$$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

$$\frac{dy}{dx} + \underbrace{\frac{x}{1+x^2}}_{\overline{P}(x)} y = \underbrace{\frac{x^2}{1+x^2}}_{\overline{Q}(x)} y^2$$

n = 2. We use the substitution $v = y^{1-2} = y^{-1}$.

$$y = v^{-1} \qquad \qquad \frac{dy}{dx} = -v^{-2}\frac{dv}{dx}$$

We get the linear equation

$$\frac{dy}{dx} + \frac{x}{1+x^2}y = \frac{x^2}{1+x^2}y^2 \qquad \frac{dv}{dx} - \frac{x}{1+x^2}v = -\frac{x^2}{1+x^2}$$
$$-v^{-2}\frac{dv}{dx} + \frac{x}{1+x^2}v^{-1} = \frac{x^2}{1+x^2}v^{-2}$$

EXAMPLES

Find a general solution

$$(1+x^2)y' + xy = x^2y^2$$

Let us solve the linear equation:

$$\frac{dv}{dx} - \frac{x}{1+x^2}v = -\frac{x^2}{1+x^2}$$

In this equation, $P(x) = -x/(1+x^2)$, $Q(x) = -x^2/(1+x^2)$. We solve it in the usual way:

$$\int P(x) dx = -\int \frac{x}{1+x^2} dx = -\frac{1}{2} \ln|1+x^2| = \ln(1+x^2)^{-1/2}$$

$$\rho(x) = (1+x^2)^{-1/2}$$

$$\int \rho(x)Q(x)\,dx = -\int \frac{x^2}{(1+x^2)^{3/2}}\,dx \leftarrow \text{can you find this integral?}$$