


## Lesson 2: Intercepts, Change and Average Rate of Change

Francisco Blanco-Silva

University of South Carolina



# WHAT DO WE KNOW?

## THE GENERAL PROGRAM

- ▶ Background Material (Basic Algebra)
- ▶ Functions
  - ▶ Definition

## WARM-UP

### A SOMEWHAT ADVANCED EXAMPLE

#### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

# WARM-UP

## A SOMEWHAT ADVANCED EXAMPLE

### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

We always start indicating independent variable and dependent variable.

# WARM-UP

## A SOMEWHAT ADVANCED EXAMPLE

### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

We always start indicating independent variable and dependent variable.

Independent variable:  $t$ , years after 2000 (let's keep it simple!)

Dependent variable:  $W$ , in millions of tons of waste.

## WARM-UP

### A SOMEWHAT ADVANCED EXAMPLE

#### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

We always start indicating independent variable and dependent variable.

Independent variable:  $t$ , years after 2000 (let's keep it simple!)

Dependent variable:  $W$ , in millions of tons of waste.

We are searching then for a function of the kind  $W = f(t) = \underbrace{a + mt}_{\text{a line!}}$

## WARM-UP

### A SOMEWHAT ADVANCED EXAMPLE

#### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

We always start indicating independent variable and dependent variable.

Independent variable:  $t$ , years after 2000 (let's keep it simple!)

Dependent variable:  $W$ , in millions of tons of waste.

We are searching then for a function of the kind  $W = f(t) = \underbrace{a + mt}_{\text{a line!}}$

$$a = f(0) = 238.3 \qquad m = \frac{251.3 - 238.3}{6 - 0} = \frac{13}{6} \approx 2.167$$

## WARM-UP

### A SOMEWHAT ADVANCED EXAMPLE

#### Example

The solid waste  $W$  generated each year in the cities of the US is increasing. The solid waste generated (in millions of tons) was 238.3 in 2000, and 251.3 in 2006.

Find a formula for this function by using the equation of the line through these two points.

We always start indicating independent variable and dependent variable.

Independent variable:  $t$ , years after 2000 (let's keep it simple!)

Dependent variable:  $W$ , in millions of tons of waste.

We are searching then for a function of the kind  $W = f(t) = \underbrace{a + mt}_{\text{a line!}}$

$$a = f(0) = 238.3 \qquad m = \frac{251.3 - 238.3}{6 - 0} = \frac{13}{6} \approx 2.167$$

Solution:

$$W = f(t) = 238.3 + \frac{13}{6}t$$



## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.
- ▶ What is the population predicted to be in 2020?
- ▶ When is the population expected to reach 55,000?

## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?
- ▶ When is the population expected to reach 55,000?

## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

# WARM-UP

## ANOTHER SOMEWHAT ADVANCED EXAMPLE

### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

$$f(t) = 55000$$

# WARM-UP

## ANOTHER SOMEWHAT ADVANCED EXAMPLE

### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

$$f(t) = 55000$$

$$30700 + 850t = 55000$$

## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

$$f(t) = 55000$$

$$30700 + 850t = 55000$$

$$850t = 55000 - 30700 = 24300$$

## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

$$f(t) = 55000$$

$$30700 + 850t = 55000$$

$$850t = 55000 - 30700 = 24300$$

$$t = \frac{24300}{850} \approx 28.59 \text{ years}$$

## WARM-UP

### ANOTHER SOMEWHAT ADVANCED EXAMPLE

#### Example

A city's population was 30,700 in the year 2000, and is growing by 850 people a year.

- ▶ Give a formula for the city's population,  $P$ , as a function of the number of years,  $t$ , since 2000.

$$P = f(t) = 30700 + 850t$$

- ▶ What is the population predicted to be in 2020?

$$f(20) = 30700 + 850 \times 20 = 47700 \text{ people}$$

- ▶ When is the population expected to reach 55,000?

$$f(t) = 55000$$

$$30700 + 850t = 55000$$

$$850t = 55000 - 30700 = 24300$$

$$t = \frac{24300}{850} \approx 28.59 \text{ years}$$

Solution: The population of this city will reach 55000 sometime during 2028.



# EXTRACTING BASIC INFORMATION FROM FUNCTIONS

INTERCEPTS, CHANGE, AVERAGE RATE OF CHANGE

In the previous examples, we have learned of the significance of the intersection of a function  $y = f(x)$  with the axes:

# EXTRACTING BASIC INFORMATION FROM FUNCTIONS

## INTERCEPTS, CHANGE, AVERAGE RATE OF CHANGE

In the previous examples, we have learned of the significance of the intersection of a function  $y = f(x)$  with the axes:

- ▶ The  $y$ -intersect indicates the **initial value** of the function.

*We obtain  $y$ -intersects by evaluating  $f(0)$ .*

# EXTRACTING BASIC INFORMATION FROM FUNCTIONS

## INTERCEPTS, CHANGE, AVERAGE RATE OF CHANGE

In the previous examples, we have learned of the significance of the intersection of a function  $y = f(x)$  with the axes:

- ▶ The  $y$ -intersect indicates the **initial value** of the function.

*We obtain  $y$ -intersects by evaluating  $f(0)$ .*

- ▶ The  $x$ -intersects (there might be many!) indicate when the function is zero.

*We obtain  $x$ -intersects by solving the equation  $f(x) = 0$*

## EXTRACTING BASIC INFORMATION FROM FUNCTIONS

### INTERCEPTS, CHANGE, AVERAGE RATE OF CHANGE

In the previous examples, we have learned of the significance of the intersection of a function  $y = f(x)$  with the axes:

- ▶ The  $y$ -intersect indicates the **initial value** of the function.

*We obtain  $y$ -intersects by evaluating  $f(0)$ .*

- ▶ The  $x$ -intersects (there might be many!) indicate when the function is zero.

*We obtain  $x$ -intersects by solving the equation  $f(x) = 0$*

Other relevant information we can ask:

- ▶ The **change** of  $f$  between  $x = a$  and  $x = b$

$$\Delta y = f(b) - f(a) \text{ (units = dep. variable)}$$

# EXTRACTING BASIC INFORMATION FROM FUNCTIONS

## INTERCEPTS, CHANGE, AVERAGE RATE OF CHANGE

In the previous examples, we have learned of the significance of the intersection of a function  $y = f(x)$  with the axes:

- ▶ The  $y$ -intersect indicates the **initial value** of the function.

*We obtain  $y$ -intersects by evaluating  $f(0)$ .*

- ▶ The  $x$ -intersects (there might be many!) indicate when the function is zero.

*We obtain  $x$ -intersects by solving the equation  $f(x) = 0$*

Other relevant information we can ask:

- ▶ The **change** of  $f$  between  $x = a$  and  $x = b$

$$\Delta y = f(b) - f(a) \text{ (units = dep. variable)}$$

- ▶ The **average rate of change** of  $f$  between  $x = a$  and  $x = b$ :

$$\frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a} \text{ (units = dep. variable/ind. variable)}$$

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?
- ▶ What was the average rate of change in height from 1960 to 1968?
- ▶ What was the change in height from 1992 to 2000?
- ▶ What was the average rate of change in height from 1992 to 2000?

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?  
 $\Delta f = f(b) - f(a)$
- ▶ What was the average rate of change in height from 1960 to 1968?
- ▶ What was the change in height from 1992 to 2000?
- ▶ What was the average rate of change in height from 1992 to 2000?

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?  
 $\Delta f = f(b) - f(a) = 213 - 185 = 28$  inches
- ▶ What was the average rate of change in height from 1960 to 1968?
- ▶ What was the change in height from 1992 to 2000?
- ▶ What was the average rate of change in height from 1992 to 2000?



## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?

$$\Delta f = f(b) - f(a) = 213 - 185 = 28 \text{ inches}$$

- ▶ What was the average rate of change in height from 1960 to 1968?

$$\text{ARC} = \frac{f(b) - f(a)}{b - a}$$

- ▶ What was the change in height from 1992 to 2000?

- ▶ What was the average rate of change in height from 1992 to 2000?

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?

$$\Delta f = f(b) - f(a) = 213 - 185 = 28 \text{ inches}$$

- ▶ What was the average rate of change in height from 1960 to 1968?

$$\text{ARC} = \frac{f(b) - f(a)}{b - a} = \frac{213 - 185}{1968 - 1960} = \frac{28}{8} = 3.5 \text{ inches/year}$$

- ▶ What was the change in height from 1992 to 2000?

- ▶ What was the average rate of change in height from 1992 to 2000?

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?

$$\Delta f = f(b) - f(a) = 213 - 185 = 28 \text{ inches}$$

- ▶ What was the average rate of change in height from 1960 to 1968?

$$\text{ARC} = \frac{f(b) - f(a)}{b - a} = \frac{213 - 185}{1968 - 1960} = \frac{28}{8} = 3.5 \text{ inches/year}$$

- ▶ What was the change in height from 1992 to 2000?

$$232 - 228 = 4 \text{ inches}$$

- ▶ What was the average rate of change in height from 1992 to 2000?

## EXAMPLES

## Example (Winning Height in Men's Olympic Pole Vault)

Year	1960	1964	1968	...	1992	1996	2000
Height (in)	185	201	213	...	228	233	232

- ▶ What was the change in height from 1960 to 1968?

$$\Delta f = f(b) - f(a) = 213 - 185 = 28 \text{ inches}$$

- ▶ What was the average rate of change in height from 1960 to 1968?

$$\text{ARC} = \frac{f(b) - f(a)}{b - a} = \frac{213 - 185}{1968 - 1960} = \frac{28}{8} = 3.5 \text{ inches/year}$$

- ▶ What was the change in height from 1992 to 2000?

$$232 - 228 = 4 \text{ inches}$$

- ▶ What was the average rate of change in height from 1992 to 2000?

$$\frac{232 - 228}{2000 - 1992} = \frac{4}{8} = 0.5 \text{ inches/year}$$

## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

## EXAMPLES

## Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

The thickness of pelican eggs decreases by an average of 0.0008 mm for every additional part per million of PCB in the environment.

## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

**The thickness of pelican eggs** decreases by an average of 0.0008 mm for every additional part per million of PCB in the environment.



## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

The thickness of pelican eggs **decreases by an average of** 0.0008 mm for every additional part per million of PCB in the environment.

## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

The thickness of pelican eggs decreases by an average of **0.0008 mm** for every additional part per million of PCB in the environment.

## EXAMPLES

### Example

High levels of PCB in the environment affect pelicans' eggs. As the concentration of PCB in environment increases, the thickness of eggshells decreases, making the eggs more likely to break.

PCB (parts per million)	87	147	204	289	356	452
thickness (mm)	0.44	0.39	0.28	0.23	0.22	0.14

What is the ARC of the thickness from 87 ppm to 452 ppm? Explain with words what that means.

$$\text{ARC} = \frac{0.14 - 0.44}{452 - 87} = \frac{-0.3}{365} \approx -0.0008 \text{ mm/ppm}$$

The thickness of pelican eggs decreases by an average of 0.0008 mm **for every additional part per million of PCB in the environment.**

# EXAMPLES

## Example (Tobacco production in the US)

Year	1996	1997	1998	1999	2000	2001	2002	2003
Production (million pounds)	1517	1787	1480	1293	1053	991	879	831

- ▶ What is the average rate of change in tobacco production between 1996 and 2003? Interpret your answer.
- ▶ During this seven-year period, is there any interval during which the average rate of change was positive? If so, when?

## EXAMPLES

## Example (Tobacco production in the US)

Year	1996	1997	1998	1999	2000	2001	2002	2003
Production (million pounds)	1517	1787	1480	1293	1053	991	879	831

- ▶ What is the average rate of change in tobacco production between 1996 and 2003? Interpret your answer.

$$\text{ARC} = \frac{831 - 1517}{2003 - 1996} = \frac{-686}{7} = -98 \text{ millions of pounds per year}$$

- ▶ During this seven-year period, is there any interval during which the average rate of change was positive? If so, when?

## EXAMPLES

## Example (Tobacco production in the US)

Year	1996	1997	1998	1999	2000	2001	2002	2003
Production (million pounds)	1517	1787	1480	1293	1053	991	879	831

- What is the average rate of change in tobacco production between 1996 and 2003? Interpret your answer.

$$\text{ARC} = \frac{831 - 1517}{2003 - 1996} = \frac{-686}{7} = -98 \text{ millions of pounds per year}$$

The production of tobacco in US decreased by an average of 98 millions of pounds every year between 1996 and 2003.

- During this seven-year period, is there any interval during which the average rate of change was positive? If so, when?

## EXAMPLES

## Example (Tobacco production in the US)

Year	1996	1997	1998	1999	2000	2001	2002	2003
Production (million pounds)	1517	1787	1480	1293	1053	991	879	831

- What is the average rate of change in tobacco production between 1996 and 2003? Interpret your answer.

$$\text{ARC} = \frac{831 - 1517}{2003 - 1996} = \frac{-686}{7} = -98 \text{ millions of pounds per year}$$

The production of tobacco in US decreased by an average of 98 millions of pounds every year between 1996 and 2003.

- During this seven-year period, is there any interval during which the average rate of change was positive? If so, when?  
Only between 1996 and 1997.