


Lesson 12: Homogeneous Second-Order Linear Equations with Constant Coefficients

Francisco Blanco-Silva

University of South Carolina



WHAT DO WE KNOW?

- ▶ The concepts of **differential equation** and **initial value problem**
- ▶ The concept of **order** of a differential equation.
- ▶ The concepts of **general solution**, **particular solution** and **singular solution**.
- ▶ **Slope fields**
- ▶ Approximations to solutions via **Euler's Method** and **Improved Euler's Method**
- ▶ **First-Order Differential Equations**
 - ▶ Separable equations
 - ▶ Homogeneous First-Order Equations
 - ▶ Linear First-Order Equations
 - ▶ Bernoulli Equations
 - ▶ General Substitution Methods
 - ▶ Exact Equations
- ▶ **Second-Order Differential Equations**
 - ▶ Reducible Equations
 - ▶ Linear Equations (Intro)

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

Can you guess (intuitively) what a solution of this equation looks like?

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

Can you guess (intuitively) what a solution of this equation looks like?

It looks like something of the form $y_1 = e^{rx}$ could do the trick. If that is the case, we may find the value of r by substitution:

$$y_1 = e^{rx}$$

$$y_1' = re^{rx}$$

$$y_1'' = r^2 e^{rx}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

Can you guess (intuitively) what a solution of this equation looks like?

It looks like something of the form $y_1 = e^{rx}$ could do the trick. If that is the case, we may find the value of r by substitution:

$$y_1 = e^{rx}$$

$$y_1' = re^{rx}$$

$$y_1'' = r^2 e^{rx}$$

Let's try:

$$ay_1'' + by_1' + cy_1 = 0$$

$$ar^2 e^{rx} + bre^{rx} + ce^{rx} = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

Can you guess (intuitively) what a solution of this equation looks like?

It looks like something of the form $y_1 = e^{rx}$ could do the trick. If that is the case, we may find the value of r by substitution:

$$y_1 = e^{rx}$$

$$y_1' = re^{rx}$$

$$y_1'' = r^2 e^{rx}$$

Let's try:

$$ay_1'' + by_1' + cy_1 = 0$$

$$ar^2 e^{rx} + bre^{rx} + ce^{rx} = 0$$

$$e^{rx}(ar^2 + br + c) = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

MOTIVATION

A homogeneous second-order linear equation with constant coefficients has the form:

$$ay'' + by' + cy = 0 \quad (a \neq 0)$$

Can you guess (intuitively) what a solution of this equation looks like?

It looks like something of the form $y_1 = e^{rx}$ could do the trick. If that is the case, we may find the value of r by substitution:

$$y_1 = e^{rx}$$

$$y_1' = re^{rx}$$

$$y_1'' = r^2 e^{rx}$$

Let's try:

$$ay_1'' + by_1' + cy_1 = 0$$

$$ar^2 e^{rx} + bre^{rx} + ce^{rx} = 0$$

$$e^{rx}(ar^2 + br + c) = 0$$

$$ar^2 + br + c = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

DEFINITION

Definition

Given a homogeneous second-order linear differential equation with constant coefficients $ay'' + by' + cy = 0$, we say that the quadratic equation $ar^2 + br + c = 0$ is the **characteristic equation**.

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

DEFINITION

Definition

Given a homogeneous second-order linear differential equation with constant coefficients $ay'' + by' + cy = 0$, we say that the quadratic equation $ar^2 + br + c = 0$ is the **characteristic equation**.

The roots of the characteristic equation give away the solutions of the differential equation. We have three cases:

- Both roots are real and different: $r_1 \neq r_2$

$$r^2 - 5r + 6 = 0 \quad \{r = 2, r = 3\}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

DEFINITION

Definition

Given a homogeneous second-order linear differential equation with constant coefficients $ay'' + by' + cy = 0$, we say that the quadratic equation $ar^2 + br + c = 0$ is the **characteristic equation**.

The roots of the characteristic equation give away the solutions of the differential equation. We have three cases:

- ▶ Both roots are real and different: $r_1 \neq r_2$

$$r^2 - 5r + 6 = 0 \quad \{r = 2, r = 3\}$$

- ▶ Both roots are real and equal: $r = r_1 = r_2$.

$$r^2 - 2r + 1 = 0 \quad \{r = 1\}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

DEFINITION

Definition

Given a homogeneous second-order linear differential equation with constant coefficients $ay'' + by' + cy = 0$, we say that the quadratic equation $ar^2 + br + c = 0$ is the **characteristic equation**.

The roots of the characteristic equation give away the solutions of the differential equation. We have three cases:

- ▶ Both roots are real and different: $r_1 \neq r_2$

$$r^2 - 5r + 6 = 0 \quad \{r = 2, r = 3\}$$

- ▶ Both roots are real and equal: $r = r_1 = r_2$.

$$r^2 - 2r + 1 = 0 \quad \{r = 1\}$$

- ▶ Both roots are complex: $r = \alpha \pm i\beta$ with $\beta > 0$.

$$r^2 + 1 = 0 \quad \{r = \pm i\}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

FIRST CASE: $r_1 \neq r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{r_1 x}$$

$$y_2 = e^{r_2 x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

FIRST CASE: $r_1 \neq r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{r_1 x}$$

$$y_2 = e^{r_2 x}$$

Note that the Wronskian is never zero:

$$y_1' = r_1 e^{r_1 x}$$

$$y_2' = r_2 e^{r_2 x}.$$

$$W(e^{r_1 x}, e^{r_2 x}) = \begin{vmatrix} e^{r_1 x} & e^{r_2 x} \\ r_1 e^{r_1 x} & r_2 e^{r_2 x} \end{vmatrix}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

FIRST CASE: $r_1 \neq r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{r_1 x}$$

$$y_2 = e^{r_2 x}$$

Note that the Wronskian is never zero:

$$y_1' = r_1 e^{r_1 x}$$

$$y_2' = r_2 e^{r_2 x}.$$

$$W(e^{r_1 x}, e^{r_2 x}) = \begin{vmatrix} e^{r_1 x} & e^{r_2 x} \\ r_1 e^{r_1 x} & r_2 e^{r_2 x} \end{vmatrix} = r_2 e^{r_1 x} e^{r_2 x} - r_1 e^{r_1 x} e^{r_2 x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

FIRST CASE: $r_1 \neq r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{r_1 x}$$

$$y_2 = e^{r_2 x}$$

Note that the Wronskian is never zero:

$$y_1' = r_1 e^{r_1 x}$$

$$y_2' = r_2 e^{r_2 x}.$$

$$W(e^{r_1 x}, e^{r_2 x}) = \begin{vmatrix} e^{r_1 x} & e^{r_2 x} \\ r_1 e^{r_1 x} & r_2 e^{r_2 x} \end{vmatrix} = r_2 e^{r_1 x} e^{r_2 x} - r_1 e^{r_1 x} e^{r_2 x} = \underbrace{(r_2 - r_1)}_{\neq 0} e^{(r_1 + r_2)x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

FIRST CASE: $r_1 \neq r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{r_1 x}$$

$$y_2 = e^{r_2 x}$$

Note that the Wronskian is never zero:

$$y_1' = r_1 e^{r_1 x}$$

$$y_2' = r_2 e^{r_2 x}.$$

$$W(e^{r_1 x}, e^{r_2 x}) = \begin{vmatrix} e^{r_1 x} & e^{r_2 x} \\ r_1 e^{r_1 x} & r_2 e^{r_2 x} \end{vmatrix} = r_2 e^{r_1 x} e^{r_2 x} - r_1 e^{r_1 x} e^{r_2 x} = \underbrace{(r_2 - r_1)}_{\neq 0} e^{(r_1 + r_2)x}$$

This means that a general solution to the differential equation has the form

$$y = A e^{r_1 x} + B e^{r_2 x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

SECOND CASE: $r = r_1 = r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{rx}$$

$$y_2 = xe^{rx}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

SECOND CASE: $r = r_1 = r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{rx}$$

$$y_2 = xe^{rx}$$

The Wronskian is also non-zero:

$$y_1' = re^{rx}$$

$$y_2' = e^{rx} + rxe^{rx} = (1 + rx)e^{rx}$$

$$W(e^{rx}, xe^{rx}) = \begin{vmatrix} e^{rx} & xe^{rx} \\ re^{rx} & (1 + rx)e^{rx} \end{vmatrix}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

SECOND CASE: $r = r_1 = r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{rx}$$

$$y_2 = xe^{rx}$$

The Wronskian is also non-zero:

$$y_1' = re^{rx}$$

$$y_2' = e^{rx} + rxe^{rx} = (1 + rx)e^{rx}$$

$$W(e^{rx}, xe^{rx}) = \begin{vmatrix} e^{rx} & xe^{rx} \\ re^{rx} & (1 + rx)e^{rx} \end{vmatrix} = (1 + rx)e^{2rx} - rxe^{2rx}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

SECOND CASE: $r = r_1 = r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{rx}$$

$$y_2 = xe^{rx}$$

The Wronskian is also non-zero:

$$y_1' = re^{rx}$$

$$y_2' = e^{rx} + rxe^{rx} = (1 + rx)e^{rx}$$

$$W(e^{rx}, xe^{rx}) = \begin{vmatrix} e^{rx} & xe^{rx} \\ re^{rx} & (1 + rx)e^{rx} \end{vmatrix} = (1 + rx)e^{2rx} - rxe^{2rx} = e^{2rx}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

SECOND CASE: $r = r_1 = r_2$ REAL ROOTS

In this case, the two possible solutions are

$$y_1 = e^{rx}$$

$$y_2 = xe^{rx}$$

The Wronskian is also non-zero:

$$y_1' = re^{rx}$$

$$y_2' = e^{rx} + rxe^{rx} = (1 + rx)e^{rx}$$

$$W(e^{rx}, xe^{rx}) = \begin{vmatrix} e^{rx} & xe^{rx} \\ re^{rx} & (1 + rx)e^{rx} \end{vmatrix} = (1 + rx)e^{2rx} - rxe^{2rx} = e^{2rx}$$

This means that a general solution to the differential equation has the form

$$y = Ae^{rx} + Bxe^{rx} = (A + Bx)e^{rx}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

The Wronskian is not zero:

$$y_1' = \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) \quad y_2' = \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x)$$

$$W(y_1, y_2) = \begin{vmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ e^{\alpha x} (\alpha \cos(\beta x) - \beta \sin(\beta x)) & e^{\alpha x} (\alpha \sin(\beta x) + \beta \cos(\beta x)) \end{vmatrix}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

The Wronskian is not zero:

$$y_1' = \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) \quad y_2' = \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x)$$

$$\begin{aligned} W(y_1, y_2) &= \begin{vmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ e^{\alpha x} (\alpha \cos(\beta x) - \beta \sin(\beta x)) & e^{\alpha x} (\alpha \sin(\beta x) + \beta \cos(\beta x)) \end{vmatrix} \\ &= e^{2\alpha x} \cos(\beta x) (\alpha \sin(\beta x) + \beta \cos(\beta x)) - e^{2\alpha x} \sin(\beta x) (\alpha \cos(\beta x) - \beta \sin(\beta x)) \end{aligned}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

The Wronskian is not zero:

$$y_1' = \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) \quad y_2' = \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x)$$

$$\begin{aligned} W(y_1, y_2) &= \begin{vmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ e^{\alpha x} (\alpha \cos(\beta x) - \beta \sin(\beta x)) & e^{\alpha x} (\alpha \sin(\beta x) + \beta \cos(\beta x)) \end{vmatrix} \\ &= e^{2\alpha x} \cos(\beta x) (\alpha \sin(\beta x) + \beta \cos(\beta x)) - e^{2\alpha x} \sin(\beta x) (\alpha \cos(\beta x) - \beta \sin(\beta x)) \\ &= \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \cos^2(\beta x) - \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \sin^2(\beta x) \end{aligned}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

The Wronskian is not zero:

$$y_1' = \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) \quad y_2' = \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x)$$

$$\begin{aligned} W(y_1, y_2) &= \begin{vmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ e^{\alpha x} (\alpha \cos(\beta x) - \beta \sin(\beta x)) & e^{\alpha x} (\alpha \sin(\beta x) + \beta \cos(\beta x)) \end{vmatrix} \\ &= e^{2\alpha x} \cos(\beta x) (\alpha \sin(\beta x) + \beta \cos(\beta x)) - e^{2\alpha x} \sin(\beta x) (\alpha \cos(\beta x) - \beta \sin(\beta x)) \\ &= \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \cos^2(\beta x) - \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \sin^2(\beta x) \\ &= \beta e^{2\alpha x} (\cos^2(\beta x) + \sin^2(\beta x)) = \beta e^{2\alpha x} \end{aligned}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

THIRD CASE: $r = \alpha \pm i\beta$ COMPLEX ROOTS ($\beta > 0$)

In this case, the two possible solutions are

$$y_1 = e^{\alpha x} \cos(\beta x)$$

$$y_2 = e^{\alpha x} \sin(\beta x)$$

The Wronskian is not zero:

$$y_1' = \alpha e^{\alpha x} \cos(\beta x) - \beta e^{\alpha x} \sin(\beta x) \quad y_2' = \alpha e^{\alpha x} \sin(\beta x) + \beta e^{\alpha x} \cos(\beta x)$$

$$\begin{aligned} W(y_1, y_2) &= \begin{vmatrix} e^{\alpha x} \cos(\beta x) & e^{\alpha x} \sin(\beta x) \\ e^{\alpha x} (\alpha \cos(\beta x) - \beta \sin(\beta x)) & e^{\alpha x} (\alpha \sin(\beta x) + \beta \cos(\beta x)) \end{vmatrix} \\ &= e^{2\alpha x} \cos(\beta x) (\alpha \sin(\beta x) + \beta \cos(\beta x)) - e^{2\alpha x} \sin(\beta x) (\alpha \cos(\beta x) - \beta \sin(\beta x)) \\ &= \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \cos^2(\beta x) - \alpha e^{2\alpha x} \sin(\beta x) \cos(\beta x) + \beta e^{2\alpha x} \sin^2(\beta x) \\ &= \beta e^{2\alpha x} (\cos^2(\beta x) + \sin^2(\beta x)) = \beta e^{2\alpha x} \end{aligned}$$

This means that a general solution to the differential equation has the form

$$y = Ae^{\alpha x} \sin(\beta x) + Be^{\alpha x} \cos(\beta x)$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 5r + 6 = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 5r + 6 = 0 \quad r = \frac{5 \pm \sqrt{25 - 4 \cdot 6}}{2} = \frac{5 \pm 1}{2} = \{2, 3\}$$

The general solution (and its derivative) is then

$$y = Ae^{2x} + Be^{3x}$$

$$y' = 2Ae^{2x} + 3Be^{3x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 5r + 6 = 0 \quad r = \frac{5 \pm \sqrt{25 - 4 \cdot 6}}{2} = \frac{5 \pm 1}{2} = \{2, 3\}$$

The general solution (and its derivative) is then

$$y = Ae^{2x} + Be^{3x}$$

$$y' = 2Ae^{2x} + 3Be^{3x}$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A + B \\ 2 = y'(0) = 2A + 3B \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 5r + 6 = 0 \quad r = \frac{5 \pm \sqrt{25 - 4 \cdot 6}}{2} = \frac{5 \pm 1}{2} = \{2, 3\}$$

The general solution (and its derivative) is then

$$y = Ae^{2x} + Be^{3x}$$

$$y' = 2Ae^{2x} + 3Be^{3x}$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A + B \\ 2 = y'(0) = 2A + 3B \end{cases} \quad \begin{cases} A = 1 \\ B = 0 \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 5y' + 6y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 5r + 6 = 0 \quad r = \frac{5 \pm \sqrt{25 - 4 \cdot 6}}{2} = \frac{5 \pm 1}{2} = \{2, 3\}$$

The general solution (and its derivative) is then

$$y = Ae^{2x} + Be^{3x}$$

$$y' = 2Ae^{2x} + 3Be^{3x}$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A + B \\ 2 = y'(0) = 2A + 3B \end{cases}$$

$$\begin{cases} A = 1 \\ B = 0 \end{cases}$$

$$y = e^{2x}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 2r + 1 = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 2r + 1 = 0 \quad r = \frac{2 \pm \sqrt{4 - 4}}{2} = 1$$

The general solution (and its derivative) is then

$$y = (A + Bx)e^x \quad y' = Be^x + (A + Bx)e^x = (A + B + Bx)e^x$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 2r + 1 = 0 \quad r = \frac{2 \pm \sqrt{4 - 4}}{2} = 1$$

The general solution (and its derivative) is then

$$y = (A + Bx)e^x \quad y' = Be^x + (A + Bx)e^x = (A + B + Bx)e^x$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = A + B \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 2r + 1 = 0 \quad r = \frac{2 \pm \sqrt{4 - 4}}{2} = 1$$

The general solution (and its derivative) is then

$$y = (A + Bx)e^x \quad y' = Be^x + (A + Bx)e^x = (A + B + Bx)e^x$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = A + B \end{cases} \quad \begin{cases} A = 1 \\ B = 1 \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' - 2y' + y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 - 2r + 1 = 0 \quad r = \frac{2 \pm \sqrt{4 - 4}}{2} = 1$$

The general solution (and its derivative) is then

$$y = (A + Bx)e^x \quad y' = Be^x + (A + Bx)e^x = (A + B + Bx)e^x$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = A + B \end{cases} \quad \begin{cases} A = 1 \\ B = 1 \end{cases} \quad y = (1 + x)e^x$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' = -y$$

$$y(0) = 1$$

$$y'(0) = 2$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' = -y$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 1 = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' = -y$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 1 = 0 \quad r = \pm i \quad (\alpha = 0, \beta = 1)$$

The general solution (and its derivative) is then

$$y = Ae^{0 \cdot x} \cos(1 \cdot x) + Be^{0 \cdot x} \sin(1 \cdot x) = A \cos x + B \sin x, \quad y' = -A \sin x + B \cos x$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' = -y$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 1 = 0 \quad r = \pm i \quad (\alpha = 0, \beta = 1)$$

The general solution (and its derivative) is then

$$y = Ae^{0 \cdot x} \cos(1 \cdot x) + Be^{0 \cdot x} \sin(1 \cdot x) = A \cos x + B \sin x, \quad y' = -A \sin x + B \cos x$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = B \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' = -y$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 1 = 0 \quad r = \pm i \quad (\alpha = 0, \beta = 1)$$

The general solution (and its derivative) is then

$$y = Ae^{0 \cdot x} \cos(1 \cdot x) + Be^{0 \cdot x} \sin(1 \cdot x) = A \cos x + B \sin x, \quad y' = -A \sin x + B \cos x$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = B \end{cases} \quad y = \cos x + 2 \sin x$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 2r + 5 = 0$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 2r + 5 = 0 \quad r = \frac{-2 \pm \sqrt{4 - 4 \cdot 5}}{2} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = \underbrace{-1 \pm 2i}_{\alpha + \beta i}$$

The general solution (and its derivative) is then

$$y = Ae^{-x} \cos(2x) + Be^{-x} \sin(2x) = e^{-x} (A \cos(2x) + B \sin(2x))$$

$$y' = -Ae^{-x} \cos(2x) - 2Ae^{-x} \sin(2x) - Be^{-x} \sin(2x) + 2Be^{-x} \cos(2x)$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 2r + 5 = 0 \quad r = \frac{-2 \pm \sqrt{4 - 4 \cdot 5}}{2} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = \underbrace{-1 \pm 2i}_{\alpha + \beta i}$$

The general solution (and its derivative) is then

$$y = Ae^{-x} \cos(2x) + Be^{-x} \sin(2x) = e^{-x} (A \cos(2x) + B \sin(2x))$$

$$y' = -Ae^{-x} \cos(2x) - 2Ae^{-x} \sin(2x) - Be^{-x} \sin(2x) + 2Be^{-x} \cos(2x)$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = -A + 2B \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 2r + 5 = 0 \quad r = \frac{-2 \pm \sqrt{4 - 4 \cdot 5}}{2} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = \underbrace{-1 \pm 2i}_{\alpha + \beta i}$$

The general solution (and its derivative) is then

$$y = Ae^{-x} \cos(2x) + Be^{-x} \sin(2x) = e^{-x} (A \cos(2x) + B \sin(2x))$$

$$y' = -Ae^{-x} \cos(2x) - 2Ae^{-x} \sin(2x) - Be^{-x} \sin(2x) + 2Be^{-x} \cos(2x)$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \\ 2 = y'(0) = -A + 2B \end{cases} \quad \begin{cases} A = 1 \\ B = \frac{3}{2} \end{cases}$$

HOMOGENEOUS SECOND-ORDER LINEAR EQUATIONS WITH CONSTANT COEFFICIENTS

EXAMPLES

Find a particular solution to the initial value problem

$$y'' + 2y' + 5y = 0$$

$$y(0) = 1$$

$$y'(0) = 2$$

First, we seek a general solution of the differential equation. We form the characteristic equation, and solve it to find the roots:

$$r^2 + 2r + 5 = 0 \quad r = \frac{-2 \pm \sqrt{4 - 4 \cdot 5}}{2} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = \underbrace{-1 \pm 2i}_{\alpha + \beta i}$$

The general solution (and its derivative) is then

$$y = Ae^{-x} \cos(2x) + Be^{-x} \sin(2x) = e^{-x} (A \cos(2x) + B \sin(2x))$$

$$y' = -Ae^{-x} \cos(2x) - 2Ae^{-x} \sin(2x) - Be^{-x} \sin(2x) + 2Be^{-x} \cos(2x)$$

We need to find the value of the constants A, B that solve the IVP:

$$\begin{cases} 1 = y(0) = A \end{cases}$$

$$\begin{cases} 2 = y'(0) = -A + 2B \end{cases}$$

$$\begin{cases} A = 1 \\ B = \frac{3}{2} \end{cases}$$

$$y = e^{-x} \left(\cos(2x) + \frac{3}{2} \sin(2x) \right)$$