


# Lesson 20: Laplace Transform of Derivatives. Transformation of Initial Value Problems

Francisco Blanco-Silva

University of South Carolina



# WHAT DO WE KNOW?

- ▶ The concepts of **differential equation** and **initial value problem**
- ▶ The concept of **order** of a differential equation.
- ▶ The concepts of **general solution**, **particular solution** and **singular solution**.
- ▶ **Slope fields**
- ▶ Approximations to solutions via **Euler's Method** and **Improved Euler's Method**
- ▶ **First-Order Differential Equations**
  - ▶ Separable equations
  - ▶ Homogeneous First-Order Equations
  - ▶ Linear First-Order Equations
  - ▶ Bernoulli Equations
  - ▶ General Substitution Methods
  - ▶ Exact Equations
- ▶ **Second-Order Differential Equations**
  - ▶ Reducible Equations
  - ▶ General Linear Equations (Intro)
  - ▶ Linear Equations with Constant Coefficients
    - ▶ Characteristic Equation
    - ▶ Variation of Parameters
    - ▶ Undetermined Coefficients

# WHAT DO WE KNOW?

## LAPLACE TRANSFORMS

$f(x)$	$\mathcal{L}\{f\} = \int_0^\infty e^{-sx} f(x) dx$	$f(x)$	$\mathcal{L}\{f\} = \int_0^\infty e^{-sx} f(x) dx$
1	$\frac{1}{s} \quad s > 0$	$cf(x) \pm g(x)$	$cF(s) \pm G(s) \quad s > \max(a, b)$
$x^p$	$\frac{\Gamma(p+1)}{s^{p+1}} \quad s > 0$	$x^n f(x)$	$(-1)^n F^{(n)} \quad s > a$
$e^{\alpha x}$	$\frac{1}{s - \alpha} \quad s > \alpha$	$e^{\alpha x} f(x)$	$F(s - \alpha) \quad s > a + \alpha$
$\sin \beta x$	$\frac{\beta}{s^2 + \beta^2} \quad s > 0$	$\frac{f(x)}{x}$	$\int_s^\infty F(\sigma) d\sigma \quad s > a$
$\cos \beta x$	$\frac{s}{s^2 + \beta^2} \quad s > 0$	$f \star g$	$F(s)G(s) \quad s > \max(a, b)$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$F(s) = \frac{1}{(s+1)^2}$$

# WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$F(s) = \frac{1}{(s+1)^2}$$

The easiest way to go about this one is to interpret the function as a  $-1$ -shift of the function  $s^{-2}$  (which is the Laplace transform of  $x$ ). It is then

$$\mathcal{L}^{-1}\left\{\frac{1}{(s+1)^2}\right\} = xe^{-x}$$

## WARM-UP

### EXAMPLES

Compute the inverse Laplace transform

$$F(s) = \frac{1}{(s+1)^2}$$

The easiest way to go about this one is to interpret the function as a  $-1$ -shift of the function  $s^{-2}$  (which is the Laplace transform of  $x$ ). It is then

$$\mathcal{L}^{-1}\left\{\frac{1}{(s+1)^2}\right\} = xe^{-x}$$

Another possibility is, of course, via convolution:

$$\mathcal{L}^{-1}\left\{\frac{1}{(s+1)^2}\right\} = \mathcal{L}^{-1}\left\{\frac{1}{s+1} \cdot \frac{1}{s+1}\right\} = \int_0^x e^{-x+t} e^{-t} dt = e^{-x} \int_0^x dt = xe^{-x}$$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2 - s}{(s^2 + 4)^2}$$

# WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2 - s}{(s^2 + 4)^2}$$

There is no shift in this expression, but the square in the denominator suggests a derivative of the Laplace transform of a sine.



## WARM-UP

### EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

There is no shift in this expression, but the square in the denominator suggests a derivative of the Laplace transform of a sine. Note that, if  $f(x) = \sin 2x$ , then

$$F(s) = \frac{2}{s^2+4},$$

$$F'(s) = -\frac{4s}{(s^2+4)^2}$$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

There is no shift in this expression, but the square in the denominator suggests a derivative of the Laplace transform of a sine. Note that, if  $f(x) = \sin 2x$ , then

$$F(s) = \frac{2}{s^2+4}, \quad F'(s) = -\frac{4s}{(s^2+4)^2}$$

In our case, we have

$$G(s) = \frac{2-s}{(s^2+4)^2} = \frac{2}{(s^2+4)^2} - \frac{s}{(s^2+4)^2}$$

## WARM-UP

## EXAMPLES

**Compute the inverse Laplace transform**

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

There is no shift in this expression, but the square in the denominator suggests a derivative of the Laplace transform of a sine. Note that, if  $f(x) = \sin 2x$ , then

$$F(s) = \frac{2}{s^2+4}, \quad F'(s) = -\frac{4s}{(s^2+4)^2}$$

In our case, we have

$$\begin{aligned} G(s) &= \frac{2-s}{(s^2+4)^2} = \frac{2}{(s^2+4)^2} - \frac{s}{(s^2+4)^2} \\ &= \frac{1}{2} \cdot \underbrace{\frac{2}{s^2+4} \cdot \frac{2}{s^2+4}}_{\sin 2x * \sin 2x} + \frac{1}{4} \cdot \underbrace{\frac{-4s}{(s^2+4)^2}}_{-x \sin 2x} \end{aligned}$$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\int_0^x \sin 2(x-t) \sin 2t \, dt$$

# WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\int_0^x \sin 2(x-t) \sin 2t \, dt = \int_0^x (\sin 2x \cos 2t - \cos 2x \sin 2t) \sin 2t \, dt$$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\begin{aligned}\int_0^x \sin 2(x-t) \sin 2t \, dt &= \int_0^x (\sin 2x \cos 2t - \cos 2x \sin 2t) \sin 2t \, dt \\ &= \sin 2x \int_0^x \cos 2t \sin 2t \, dt - \cos 2x \int_0^x \sin^2 2t \, dt\end{aligned}$$

# WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\begin{aligned} \int_0^x \sin 2(x-t) \sin 2t \, dt &= \int_0^x (\sin 2x \cos 2t - \cos 2x \sin 2t) \sin 2t \, dt \\ &= \sin 2x \int_0^x \cos 2t \sin 2t \, dt - \cos 2x \int_0^x \sin^2 2t \, dt \\ &= \frac{1}{2} \sin 2x \int_0^x \sin 4t \, dt - \frac{1}{8} \cos 2x (4x - \sin 4x) \end{aligned}$$

## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\begin{aligned}\int_0^x \sin 2(x-t) \sin 2t \, dt &= \int_0^x (\sin 2x \cos 2t - \cos 2x \sin 2t) \sin 2t \, dt \\&= \sin 2x \int_0^x \cos 2t \sin 2t \, dt - \cos 2x \int_0^x \sin^2 2t \, dt \\&= \frac{1}{2} \sin 2x \int_0^x \sin 4t \, dt - \frac{1}{8} \cos 2x (4x - \sin 4x) \\&= \frac{1}{8} \sin 2x (1 - \cos(4x)) - \frac{1}{8} \cos 2x (4x - \sin 4x)\end{aligned}$$



## WARM-UP

## EXAMPLES

Compute the inverse Laplace transform

$$G(s) = \frac{2-s}{(s^2+4)^2}$$

It only remains to compute the convolution of  $\sin 2x$  with itself:

$$\begin{aligned}\int_0^x \sin 2(x-t) \sin 2t \, dt &= \int_0^x (\sin 2x \cos 2t - \cos 2x \sin 2t) \sin 2t \, dt \\&= \sin 2x \int_0^x \cos 2t \sin 2t \, dt - \cos 2x \int_0^x \sin^2 2t \, dt \\&= \frac{1}{2} \sin 2x \int_0^x \sin 4t \, dt - \frac{1}{8} \cos 2x (4x - \sin 4x) \\&= \frac{1}{8} \sin 2x (1 - \cos(4x)) - \frac{1}{8} \cos 2x (4x - \sin 4x)\end{aligned}$$

The solution of this problem is then

$$g(x) = \frac{1}{16} \left( \sin 2x (1 - \cos 4x) - \cos 2x (4x - \sin 4x) \right) - \frac{1}{4} x \sin 2x$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

*Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .*

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

*Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .*

We prove it in the usual way:

$$\mathcal{L}\{f'\} = \int_0^{\infty} e^{-sx} f'(x) dx$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

*Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .*

We prove it in the usual way:

$$\mathcal{L}\{f'\} = \int_0^{\infty} e^{-sx} f'(x) dx = \lim_{A \rightarrow \infty} \int_0^A \underbrace{e^{-sx}}_u \underbrace{f'(x) dx}_{dv}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .

We prove it in the usual way:

$$\begin{aligned}\mathcal{L}\{f'\} &= \int_0^{\infty} e^{-sx} f'(x) dx = \lim_{A \rightarrow \infty} \int_0^A \underbrace{e^{-sx}}_u \underbrace{f'(x)}_{dv} dx \\ &= \lim_{A \rightarrow \infty} \left[ f(x)e^{-sx} \Big|_0^A + s \int_0^A f(x)e^{-sx} dx \right]\end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .

We prove it in the usual way:

$$\begin{aligned}\mathcal{L}\{f'\} &= \int_0^\infty e^{-sx} f'(x) dx = \lim_{A \rightarrow \infty} \int_0^A \underbrace{e^{-sx}}_u \underbrace{f'(x)}_{dv} dx \\ &= \lim_{A \rightarrow \infty} \left[ \underbrace{f(x)e^{-sx}}_{\text{blue}} \Big|_0^A + s \int_0^A f(x) e^{-sx} dx \right]\end{aligned}$$

Note that

$$\lim_{A \rightarrow \infty} \left( f(x) e^{-sx} \Big|_0^A \right) = \lim_{A \rightarrow \infty} \left( f(A) e^{-sA} - f(0) \right)$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .

We prove it in the usual way:

$$\begin{aligned}\mathcal{L}\{f'\} &= \int_0^\infty e^{-sx} f'(x) dx = \lim_{A \rightarrow \infty} \int_0^A \underbrace{e^{-sx}}_u \underbrace{f'(x)}_{dv} dx \\ &= \lim_{A \rightarrow \infty} \left[ f(x)e^{-sx} \Big|_0^A + s \int_0^A f(x)e^{-sx} dx \right]\end{aligned}$$

Note that

$$\begin{aligned}\lim_{A \rightarrow \infty} \left( f(x)e^{-sx} \Big|_0^A \right) &= \lim_{A \rightarrow \infty} \left( f(A)e^{-sA} - f(0) \right) \\ &\leq \lim_{A \rightarrow \infty} Ke^{aA}e^{-sA} - f(0)\end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .

We prove it in the usual way:

$$\begin{aligned}\mathcal{L}\{f'\} &= \int_0^\infty e^{-sx} f'(x) dx = \lim_{A \rightarrow \infty} \int_0^A \underbrace{e^{-sx}}_u \underbrace{f'(x)}_{dv} dx \\ &= \lim_{A \rightarrow \infty} \left[ f(x)e^{-sx} \Big|_0^A + s \int_0^A f(x)e^{-sx} dx \right]\end{aligned}$$

Note that

$$\begin{aligned}\lim_{A \rightarrow \infty} \left( f(x)e^{-sx} \Big|_0^A \right) &= \lim_{A \rightarrow \infty} \left( f(A)e^{-sA} - f(0) \right) \\ &\leq \lim_{A \rightarrow \infty} Ke^{aA}e^{-sA} - f(0) = \lim_{A \rightarrow \infty} Ke^{(a-s)A} - f(0)\end{aligned}$$



# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

*Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .*

It must then be

$$\lim_{A \rightarrow \infty} \left( f(x)e^{-sx} \Big|_0^A \right) = -f(0)$$

only if  $a - s < 0$ . The limit diverges otherwise.

# LAPLACE TRANSFORM OF DERIVATIVES

## Theorem

*Suppose that both  $f$  and  $f'$  are both continuous functions in  $(0, \infty)$ . Suppose further that there exists constants  $K, a, M$  such that  $|f(x)| \leq Ke^{at}$  for  $x > M$ . Then  $\mathcal{L}\{f'\} = s\mathcal{L}\{f\} - f(0)$  for  $s > a$ .*

It must then be

$$\lim_{A \rightarrow \infty} \left( f(x)e^{-sx} \Big|_0^A \right) = -f(0)$$

only if  $a - s < 0$ . The limit diverges otherwise.

In summary: If  $s > a$ ,

$$\mathcal{L}\{f'\} = -f(0) + s \underbrace{\int_0^\infty e^{-sx} f(x) dx}_{\mathcal{L}\{f\}} = s\mathcal{L}\{f\} - f(0)$$

# LAPLACE TRANSFORM OF DERIVATIVES

We may generalize this result:

## Theorem

*If  $f, f', f'', \dots, f^{(n-1)}$  are all continuous functions in  $(0, \infty)$ , and they are good enough, then*

$$\mathcal{L}\{f''\} = s^2 \mathcal{L}\{f\} - sf(0) - f'(0)$$

$$\mathcal{L}\{f'''\} = s^3 \mathcal{L}\{f\} - s^2 f(0) - sf'(0) - f''(0)$$

$$\vdots$$

$$\mathcal{L}\{f^{(n)}\} = s^n \mathcal{L}\{f\} - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{(n-1)}(0)$$

# LAPLACE TRANSFORM OF DERIVATIVES

## EXAMPLES

Use the transform of  $\sin \beta x$  and the transform of its derivative to deduct the formula

$$\mathcal{L}\{\cos \beta x\} = \frac{s}{s^2 + \beta^2}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## EXAMPLES

Use the transform of  $\sin \beta x$  and the transform of its derivative to deduct the formula

$$\mathcal{L}\{\cos \beta x\} = \frac{s}{s^2 + \beta^2}$$

The trick here is to realize that  $\beta \cos \beta x$  is the derivative of  $f(x) = \sin \beta x$ . In that case,

$$\mathcal{L}\{\cos \beta x\} = \frac{1}{\beta} \mathcal{L}\{\beta \cos \beta x\} = \frac{1}{\beta} \mathcal{L}\{f'(x)\}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## EXAMPLES

Use the transform of  $\sin \beta x$  and the transform of its derivative to deduct the formula

$$\mathcal{L}\{\cos \beta x\} = \frac{s}{s^2 + \beta^2}$$

The trick here is to realize that  $\beta \cos \beta x$  is the derivative of  $f(x) = \sin \beta x$ . In that case,

$$\begin{aligned}\mathcal{L}\{\cos \beta x\} &= \frac{1}{\beta} \mathcal{L}\{\beta \cos \beta x\} = \frac{1}{\beta} \mathcal{L}\{f'(x)\} \\ &= \frac{1}{\beta} \left( s \mathcal{L}\{f(x)\} - f(0) \right)\end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## EXAMPLES

Use the transform of  $\sin \beta x$  and the transform of its derivative to deduct the formula

$$\mathcal{L}\{\cos \beta x\} = \frac{s}{s^2 + \beta^2}$$

The trick here is to realize that  $\beta \cos \beta x$  is the derivative of  $f(x) = \sin \beta x$ . In that case,

$$\begin{aligned}\mathcal{L}\{\cos \beta x\} &= \frac{1}{\beta} \mathcal{L}\{\beta \cos \beta x\} = \frac{1}{\beta} \mathcal{L}\{f'(x)\} \\ &= \frac{1}{\beta} \left( s \mathcal{L}\{f(x)\} - f(0) \right) \\ &= \frac{1}{\beta} \left( s \frac{\beta}{s^2 + \beta^2} - \sin 0 \right)\end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## EXAMPLES

Use the transform of  $\sin \beta x$  and the transform of its derivative to deduct the formula

$$\mathcal{L}\{\cos \beta x\} = \frac{s}{s^2 + \beta^2}$$

The trick here is to realize that  $\beta \cos \beta x$  is the derivative of  $f(x) = \sin \beta x$ . In that case,

$$\begin{aligned}\mathcal{L}\{\cos \beta x\} &= \frac{1}{\beta} \mathcal{L}\{\beta \cos \beta x\} = \frac{1}{\beta} \mathcal{L}\{f'(x)\} \\ &= \frac{1}{\beta} \left( s \mathcal{L}\{f(x)\} - f(0) \right) \\ &= \frac{1}{\beta} \left( s \frac{\beta}{s^2 + \beta^2} - \sin 0 \right) \\ &= \frac{s}{s^2 + \beta^2}\end{aligned}$$



# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

$$\mathcal{L}\{y'' + 2y' + y\} = \mathcal{L}\{e^{3x}\}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

$$\begin{aligned}\mathcal{L}\{y'' + 2y' + y\} &= \mathcal{L}\{e^{3x}\} \\ \mathcal{L}\{y''\} + 2\mathcal{L}\{y'\} + \mathcal{L}\{y\} &= \frac{1}{s-3} \quad (s > 3)\end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

$$\mathcal{L}\{y'' + 2y' + y\} = \mathcal{L}\{e^{3x}\}$$

$$\mathcal{L}\{y''\} + 2\mathcal{L}\{y'\} + \mathcal{L}\{y\} = \frac{1}{s-3} \quad (s > 3)$$

$$\left(s^2\mathcal{L}\{y\} - sy(0) - y'(0)\right) + 2\left(s\mathcal{L}\{y\} - y(0)\right) + \mathcal{L}\{y\} = \frac{1}{s-3}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

$$\mathcal{L}\{y'' + 2y' + y\} = \mathcal{L}\{e^{3x}\}$$

$$\mathcal{L}\{y''\} + 2\mathcal{L}\{y'\} + \mathcal{L}\{y\} = \frac{1}{s-3} \quad (s > 3)$$

$$\left(s^2\mathcal{L}\{y\} - sy(0) - y'(0)\right) + 2\left(s\mathcal{L}\{y\} - y(0)\right) + \mathcal{L}\{y\} = \frac{1}{s-3}$$

$$(s^2 + 2s + 1)\mathcal{L}\{y\} = \frac{1}{s-3}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

Note how we may use the Laplace Transform to compute particular solutions of an IVP:

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

The Laplace transform of the left hand-side must be equal to the Laplace transform of the right hand-side. Our target in this step is to compute the Laplace transform of a general solution  $y$ ,  $\mathcal{L}\{y\}$ :

$$\mathcal{L}\{y'' + 2y' + y\} = \mathcal{L}\{e^{3x}\}$$

$$\mathcal{L}\{y''\} + 2\mathcal{L}\{y'\} + \mathcal{L}\{y\} = \frac{1}{s-3} \quad (s > 3)$$

$$\left(s^2\mathcal{L}\{y\} - sy(0) - y'(0)\right) + 2\left(s\mathcal{L}\{y\} - y(0)\right) + \mathcal{L}\{y\} = \frac{1}{s-3}$$

$$(s^2 + 2s + 1)\mathcal{L}\{y\} = \frac{1}{s-3}$$

$$\mathcal{L}\{y\} = \frac{1}{(s-3)(s^2 + 2s + 1)} \quad (s > 3)$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

In the second stage, we compute the inverse Laplace transform of this function:

$$y = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s^2+2s+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s+1)^2} \right\}$$



# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

In the second stage, we compute the inverse Laplace transform of this function:

$$\begin{aligned} y &= \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s^2+2s+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{A}{s-3} + \frac{B}{s+1} + \frac{C}{(s+1)^2} \right\} \end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

In the second stage, we compute the inverse Laplace transform of this function:

$$\begin{aligned} y &= \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s^2+2s+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{A}{s-3} + \frac{B}{s+1} + \frac{C}{(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{1}{16} \cdot \frac{1}{s-3} - \frac{1}{16} \cdot \frac{1}{s+1} - \frac{1}{4} \cdot \frac{1}{(s+1)^2} \right\} \end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

In the second stage, we compute the inverse Laplace transform of this function:

$$\begin{aligned} y &= \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s^2+2s+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{A}{s-3} + \frac{B}{s+1} + \frac{C}{(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{1}{16} \cdot \frac{1}{s-3} - \frac{1}{16} \cdot \frac{1}{s+1} - \frac{1}{4} \cdot \frac{1}{(s+1)^2} \right\} \\ &= \frac{1}{16} \mathcal{L}^{-1} \left\{ \frac{1}{s-3} \right\} - \frac{1}{16} \mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} - \frac{1}{4} \mathcal{L}^{-1} \left\{ \frac{1}{(s+1)^2} \right\} \end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y'' + 2y' + y = e^{3x}, \quad y(0) = y'(0) = 0$$

In the second stage, we compute the inverse Laplace transform of this function:

$$\begin{aligned} y &= \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s^2+2s+1)} \right\} = \mathcal{L}^{-1} \left\{ \frac{1}{(s-3)(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{A}{s-3} + \frac{B}{s+1} + \frac{C}{(s+1)^2} \right\} \\ &= \mathcal{L}^{-1} \left\{ \frac{1}{16} \cdot \frac{1}{s-3} - \frac{1}{16} \cdot \frac{1}{s+1} - \frac{1}{4} \cdot \frac{1}{(s+1)^2} \right\} \\ &= \frac{1}{16} \mathcal{L}^{-1} \left\{ \frac{1}{s-3} \right\} - \frac{1}{16} \mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} - \frac{1}{4} \mathcal{L}^{-1} \left\{ \frac{1}{(s+1)^2} \right\} \\ &= \frac{1}{16} e^{3x} - \frac{1}{16} e^{-x} - \frac{1}{4} x e^{-x} \end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

$$\mathcal{L}\{y'''\} - 2\mathcal{L}\{y''\} = \frac{4s}{(s^2 + 4)^2} \quad (s > 0)$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

$$\mathcal{L}\{y'''\} - 2\mathcal{L}\{y''\} = \frac{4s}{(s^2 + 4)^2} \quad (s > 0)$$

$$\left(s^3 \mathcal{L}\{y\} - s^2 y(0) - s y'(0) - y''(0)\right) - 2\left(s^2 \mathcal{L}\{y\} - s y(0) - y'(0)\right) = \frac{4s}{(s^2 + 4)^2}$$



# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

$$\mathcal{L}\{y'''\} - 2\mathcal{L}\{y''\} = \frac{4s}{(s^2 + 4)^2} \quad (s > 0)$$

$$\left(s^3 \mathcal{L}\{y\} - s^2 y(0) - s y'(0) - y''(0)\right) - 2\left(s^2 \mathcal{L}\{y\} - s y(0) - y'(0)\right) = \frac{4s}{(s^2 + 4)^2}$$

$$(s^3 - 2s^2) \mathcal{L}\{y\} - s + 2 = \frac{4s}{(s^2 + 4)^2}$$

## LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

## Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

$$\mathcal{L}\{y'''\} - 2\mathcal{L}\{y''\} = \frac{4s}{(s^2 + 4)^2} \quad (s > 0)$$

$$\left(s^3 \mathcal{L}\{y\} - s^2 y(0) - s y'(0) - y''(0)\right) - 2\left(s^2 \mathcal{L}\{y\} - s y(0) - y'(0)\right) = \frac{4s}{(s^2 + 4)^2}$$

$$(s^3 - 2s^2) \mathcal{L}\{y\} - s + 2 = \frac{4s}{(s^2 + 4)^2}$$

$$\mathcal{L}\{y\} = \frac{4s}{(s^2 + 4)^2(s^3 - 2s^2)} + \frac{2 - s}{s^3 - 2s^2}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We proceed as before, finding first  $\mathcal{L}\{y\}$

$$\mathcal{L}\{y''' - 2y''\} = \mathcal{L}\{x \sin 2x\}$$

$$\mathcal{L}\{y'''\} - 2\mathcal{L}\{y''\} = \frac{4s}{(s^2 + 4)^2} \quad (s > 0)$$

$$\left(s^3 \mathcal{L}\{y\} - s^2 y(0) - s y'(0) - y''(0)\right) - 2\left(s^2 \mathcal{L}\{y\} - s y(0) - y'(0)\right) = \frac{4s}{(s^2 + 4)^2}$$

$$(s^3 - 2s^2) \mathcal{L}\{y\} - s + 2 = \frac{4s}{(s^2 + 4)^2}$$

$$\mathcal{L}\{y\} = \frac{4s}{(s^2 + 4)^2(s^3 - 2s^2)} + \frac{2 - s}{s^3 - 2s^2}$$

$$\mathcal{L}\{y\} = \frac{2 - s}{s^2(s - 2)} + \frac{4}{s(s - 2)(s^2 + 4)^2} = -\frac{1}{s^2} + \frac{4}{s(s - 2)(s^2 + 4)^2}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We now compute the inverse Laplace transform:

$$y = -\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} + \mathcal{L}^{-1}\left\{\frac{A}{s} + \frac{B}{s-2} + \frac{Cs+D}{s^2+4} + \frac{Es+F}{(s^2+4)^2}\right\}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We now compute the inverse Laplace transform:

$$\begin{aligned} y &= -\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} + \mathcal{L}^{-1}\left\{\frac{A}{s} + \frac{B}{s-2} + \frac{Cs+D}{s^2+4} + \frac{Es+F}{(s^2+4)^2}\right\} \\ &= -\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} - \frac{1}{8}\mathcal{L}^{-1}\left\{\frac{1}{s}\right\} - \frac{1}{32}\mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} \\ &\quad + \frac{3}{32}\mathcal{L}^{-1}\left\{\frac{s}{s^2+4}\right\} - \frac{1}{32}\mathcal{L}^{-1}\left\{\frac{2}{s^2+4}\right\} + \frac{1}{4}\mathcal{L}^{-1}\left\{\frac{s-2}{(s^2+4)^2}\right\} \end{aligned}$$

# LAPLACE TRANSFORM OF DERIVATIVES

## TRANSFORMATION OF INITIAL VALUE PROBLEMS

### Example

$$y''' - 2y'' = x \sin 2x, \quad y(0) = 0, y'(0) = 1, y''(0) = 0$$

We now compute the inverse Laplace transform:

$$\begin{aligned} y &= -\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} + \mathcal{L}^{-1}\left\{\frac{A}{s} + \frac{B}{s-2} + \frac{Cs+D}{s^2+4} + \frac{Es+F}{(s^2+4)^2}\right\} \\ &= -\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} - \frac{1}{8}\mathcal{L}^{-1}\left\{\frac{1}{s}\right\} - \frac{1}{32}\mathcal{L}^{-1}\left\{\frac{1}{s-2}\right\} \\ &\quad + \frac{3}{32}\mathcal{L}^{-1}\left\{\frac{s}{s^2+4}\right\} - \frac{1}{32}\mathcal{L}^{-1}\left\{\frac{2}{s^2+4}\right\} + \frac{1}{4}\mathcal{L}^{-1}\left\{\frac{s-2}{(s^2+4)^2}\right\} \\ &= -x - \frac{1}{8} - \frac{1}{32}e^{2x} + \frac{3}{32}\cos 2x - \frac{1}{32}\sin 2x \\ &\quad - \frac{1}{64}\left(\sin 2x(1 - \cos 4x) - \cos 2x(4x - \sin 4x)\right) + \frac{1}{16}x \sin 2x \end{aligned}$$