| Name:   |  |
|---------|--|
| VIP ID: |  |

- Write your name and your VIP ID in the space provided above.
- The test has five (5) pages, including this one.
- You must show sufficient work to justify all answers unless otherwise stated in the problem. Correct answers with inconsistent work may not be given credit.
- Credit for each problem is given in parentheses at the right of the problem number.
- No books, or notes may be used on this test.
- An approved calculator may be used on this test.

| Page  | Max. points | Your points |
|-------|-------------|-------------|
| 2     | 20          |             |
| 3     | 20          |             |
| 4     | 40          |             |
| 5     | 20          |             |
| Total | 100         |             |


**Problem 1** (10 pts). The function  $f(x) = x^4 - 5x^3 + 11x$  has a critical point at x = 1. Identify what kind of critical point it is.

- $\bigcirc f(x)$  has a local maximum at x = 1.
- $\bigcirc f(x)$  has a local minimum at x = 1.
- $\bigcirc x = 1$  is neither maximum nor minimum of f(x).

**Problem 2** (10 pts). Find intervals of increase/decrease and concavity of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 9.$$

**Problem 3** (10 pts). Concerning the graph of the function below, which of the following statements is true?



- $\bigcirc$  The derivative is zero at two values of x, both being local maxima.
- $\bigcirc$  The derivative is zero at two values of x, one is a local maximum, while the other is a local minimum.
- $\bigcirc$  The derivative is zero at two values of x, one is a local maximum on the interval, while the other is neither a local maximum nor a minimum.
- $\bigcirc$  The derivative is zero at two values of x, one is a local minimum on the interval, while the other is neither a local maximum nor a minimum.
- $\bigcirc$  The derivative is zero only at one value of x, where it is a local minimum.

**Problem 4** (10 pts). Find the absolute maximum and the absolute minimum values of the function  $f(x) = 2x^3 - 9x^2$  over the interval  $-1 \le x \le 6$ .

**Problem 5** (10 pts). Suppose P(t) is the number of individuals infected by a disease t days after it was first detected. Interpret P'(50) = 200.

**Problem 6** (10 pts). A baseball team plays in a stadium that holds 50,000 spectators. With the ticket price at \$8 the average attendance has been 19,000. When the price dropped to \$4, the average attendance rose to 25,000. Find a linear demand function D(q), where q is the number of spectators.

**Problem 7** (10 pts). The total cost of producing q units of good is given by C(q) = 7.3q + 56000.

- (a) What is the total cost of producing 4200 units?
- (b) What is the cost of the 4201st item?

**Problem 8** (10 pts). The cost function of a product (in dollars) is given by C(q) = 23.56 + 0.04q, where q is the number of units sold. What is the average cost of selling 200 units?

**Problem 9** (10 pts). A manufacturer has been selling 1300 flat screen TVs a week at \$3600 each. A market survey indicates that for each \$100 rebate offered to a buyer, the number of TVs sold increase by 100 per week. The weekly cost function is given by C(q) = 780000 + 1200q. What should be the rebate offered in order to maximize profit?

**Problem 10** (10 pts). At a price of \$30 per ticket, a musical theater group can fill every seat in the theater, which has a capacity of 2800. For every \$2.5 decrease in price, the number of people buying tickets increases by 175. What ticket price maximizes revenue?