

We are going to define two operations that can be performed on vector fields. Each operation resembles differentiation, but one produces a vector field, whereas the other produces a scalar field.

Definition. If $\mathbf{F} = \langle P, Q, R \rangle$ is a vector field on \mathbb{R}^3 and the partial derivatives of P , Q and R all exist, then the *curl* of \mathbf{F} is the vector field on \mathbb{R}^3 defined by

$$\operatorname{curl} \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \mathbf{k}.$$

The *divergence* of \mathbf{F} is the function of three variables defined by

$$\operatorname{div} \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}.$$

Problem 1 (5 pts). Assume that f is a function of three variables that satisfies

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}, \quad \frac{\partial^2 f}{\partial x \partial z} = \frac{\partial^2 f}{\partial z \partial x}, \quad \frac{\partial^2 f}{\partial z \partial y} = \frac{\partial^2 f}{\partial y \partial z}.$$

Prove that in this case, $\operatorname{curl}(\nabla f) = \mathbf{0}$.

Problem 2 (5 pts). Note that the previous problem implies that conservative vector fields will have zero curl. Determine, using this trick, whether $\mathbf{F} = \langle y^2 z^3, 2xyz^3, 3xy^2 z^2 \rangle$ is a conservative vector field or not.

Problem 3 (15 pts). For the vector field $\mathbf{F} = \cos(xz)\mathbf{j} - \sin(xy)\mathbf{k}$, compute:

- (a) $\operatorname{curl} \mathbf{F}$
- (b) $\operatorname{div} \mathbf{F}$
- (c) $\operatorname{div} \operatorname{curl} \mathbf{F}$

Use the back of this page for this problem