


Lesson 13: Using the Derivative I

Francisco Blanco-Silva

University of South Carolina



WHAT DO WE KNOW?

THE GENERAL PROGRAM

► Functions

- x - and y -**intercepts** ($f(x) = 0, f(0)$)
- **Change** from $x = a$ to $x = b$

$$\Delta y = f(b) - f(a)$$

- **Average Rate of Change** from $x = a$ to $x = b$

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

- **Relative Change** from $x = a$ to $x = b$

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

- **Instantaneous Rate of Change** at $x = a$

$$f'(a)$$

- **Relative Rate of Change** at $x = a$

$$\frac{f'(a)}{f(a)}$$

► Linear Functions:

$$f(x) = b + mx$$

► Exponential Functions

$$P_0 a^t = P_0(1 + r)^t = P_0 e^{kt}$$

► Power Functions

$$kx^p$$

► Polynomials

$$a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**.

We denote them by $f''(x)$.

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$g(x) = \ln(3x^2 - 4)$$

$$h(x) = (x^5 - x^{1/3})^2$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13 \qquad f'(x) = 2\pi x - 3e$$

$$g(x) = \ln(3x^2 - 4)$$

$$h(x) = (x^5 - x^{1/3})^2$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$h(x) = (x^5 - x^{1/3})^2$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$h(x) = (x^5 - x^{1/3})^2$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$g''(x) = \frac{6(3x^2 - 4) - 6x \cdot 6x}{(3x^2 - 4)^2}$$

$$h(x) = (x^5 - x^{1/3})^2$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$g''(x) = \frac{6(3x^2 - 4) - 6x \cdot 6x}{(3x^2 - 4)^2}$$

$$h(x) = (x^5 - x^{1/3})^2$$

$$h'(x) = 2(x^5 - x^{1/3})(5x^4 - \frac{1}{3}x^{-2/3})$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$g''(x) = \frac{6(3x^2 - 4) - 6x \cdot 6x}{(3x^2 - 4)^2}$$

$$h(x) = (x^5 - x^{1/3})^2$$

$$\begin{aligned} h'(x) &= 2(x^5 - x^{1/3})(5x^4 - \tfrac{1}{3}x^{-2/3}) \\ &= (2x^5 - 2x^{1/3})(5x^4 - \tfrac{1}{3}x^{-2/3}) \end{aligned}$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$g''(x) = \frac{6(3x^2 - 4) - 6x \cdot 6x}{(3x^2 - 4)^2}$$

$$h(x) = (x^5 - x^{1/3})^2$$

$$\begin{aligned} h'(x) &= 2(x^5 - x^{1/3})(5x^4 - \tfrac{1}{3}x^{-2/3}) \\ &= (2x^5 - 2x^{1/3})(5x^4 - \tfrac{1}{3}x^{-2/3}) \\ &= 10x^9 - \tfrac{32}{3}x^{13/3} + \tfrac{2}{3}x^{-1/3} \end{aligned}$$

USING THE DERIVATIVE

Taking derivatives of a derivative yields what we call a **second derivative**. We denote them by $f''(x)$. Let us compute some:

Compute the second derivative of the following functions:

$$f(x) = \pi x^2 - 3ex + 13$$

$$f'(x) = 2\pi x - 3e$$

$$f''(x) = 2\pi$$

$$g(x) = \ln(3x^2 - 4)$$

$$g'(x) = \frac{6x}{3x^2 - 4}$$

$$g''(x) = \frac{6(3x^2 - 4) - 6x \cdot 6x}{(3x^2 - 4)^2}$$

$$h(x) = (x^5 - x^{1/3})^2$$

$$h'(x) = 2(x^5 - x^{1/3})(5x^4 - \frac{1}{3}x^{-2/3})$$

$$= (2x^5 - 2x^{1/3})(5x^4 - \frac{1}{3}x^{-2/3})$$

$$= 10x^9 - \frac{32}{3}x^{13/3} + \frac{2}{3}x^{-1/3}$$

$$h''(x) = 90x^8 - \frac{416}{9}x^{10/3} - \frac{2}{9}x^{-4/3}$$

USING THE DERIVATIVE

HOW DOES THE SIGN OF DERIVATIVES AFFECT THE SHAPE OF A FUNCTION?

Because the derivative is linked to the values of slopes of tangent lines, we have the following results:

If the derivative is...	then, the function is...
$f'(x) > 0$	increasing at x
$f'(x) < 0$	decreasing at x
$f'(x) = 0$	<i>stationary</i> at x

USING THE DERIVATIVE

HOW DOES THE SIGN OF DERIVATIVES AFFECT THE SHAPE OF A FUNCTION?

Because the derivative is linked to the values of slopes of tangent lines, we have the following results:

If the derivative is...	then, the function is...
$f'(x) > 0$	increasing at x
$f'(x) < 0$	decreasing at x
$f'(x) = 0$	<i>stationary</i> at x

We say that f has a **critical point** at x if $f'(x) = 0$, or $f'(x)$ is not defined.

USING THE DERIVATIVE

HOW DOES THE SIGN OF DERIVATIVES AFFECT THE SHAPE OF A FUNCTION?

Because the derivative is linked to the values of slopes of tangent lines, we have the following results:

If the derivative is...	then, the function is...
$f'(x) > 0$	increasing at x
$f'(x) < 0$	decreasing at x
$f'(x) = 0$	<i>stationary</i> at x

We say that f has a **critical point** at x if $f'(x) = 0$, or $f'(x)$ is not defined.

The sign of the second derivative also offers a clear interpretation in terms of graph shape.

If the second derivative is...	then, the function is...
$f''(x) > 0$	concave upwards at x (∪)
$f''(x) < 0$	concave downwards at x (∩)
$f''(x) = 0$	the concavity might change at x

USING THE DERIVATIVE

HOW DOES THE SIGN OF DERIVATIVES AFFECT THE SHAPE OF A FUNCTION?

Because the derivative is linked to the values of slopes of tangent lines, we have the following results:

If the derivative is...	then, the function is...
$f'(x) > 0$	increasing at x
$f'(x) < 0$	decreasing at x
$f'(x) = 0$	<i>stationary</i> at x

We say that f has a **critical point** at x if $f'(x) = 0$, or $f'(x)$ is not defined.

The sign of the second derivative also offers a clear interpretation in terms of graph shape.

If the second derivative is...	then, the function is...
$f''(x) > 0$	concave upwards at x (∪)
$f''(x) < 0$	concave downwards at x (∩)
$f''(x) = 0$	the concavity might change at x

We say that f has an **inflection point** at x if the concavity of f changes at x .

USING THE DERIVATIVE

MAXIMA AND MINIMA

Definition

We say that a critical point $x = a$ is

- ▶ a (local) **minimum** of f , if $f(a) \leq f(x)$ for nearby values of x .
- ▶ a (local) **maximum** of f , if $f(a) \geq f(x)$ for nearby values of x .

USING THE DERIVATIVE

MAXIMA AND MINIMA

Definition

We say that a critical point $x = a$ is

- ▶ a (local) **minimum** of f , if $f(a) \leq f(x)$ for nearby values of x .
- ▶ a (local) **maximum** of f , if $f(a) \geq f(x)$ for nearby values of x .

We have two tests to verify whether a critical point is a maximum or a minimum

USING THE DERIVATIVE

MAXIMA AND MINIMA

Definition

We say that a critical point $x = a$ is

- ▶ a (local) **minimum** of f , if $f(a) \leq f(x)$ for nearby values of x .
- ▶ a (local) **maximum** of f , if $f(a) \geq f(x)$ for nearby values of x .

We have two tests to verify whether a critical point is a maximum or a minimum

The First Derivative Test

- ▶ If $f'(c) = 0$, $f'(x) < 0$ for $x < c$, and $f'(x) > 0$ for $x > c$, then $x = c$ is a local minimum.
- ▶ If $f'(c) = 0$, $f'(x) > 0$ for $x < c$, and $f'(x) < 0$ for $x > c$, then $x = c$ is a local maximum.

USING THE DERIVATIVE

MAXIMA AND MINIMA

Definition

We say that a critical point $x = a$ is

- ▶ a (local) **minimum** of f , if $f(a) \leq f(x)$ for nearby values of x .
- ▶ a (local) **maximum** of f , if $f(a) \geq f(x)$ for nearby values of x .

We have two tests to verify whether a critical point is a maximum or a minimum

The First Derivative Test

- ▶ If $f'(c) = 0$, $f'(x) < 0$ for $x < c$, and $f'(x) > 0$ for $x > c$, then $x = c$ is a local minimum.
- ▶ If $f'(c) = 0$, $f'(x) > 0$ for $x < c$, and $f'(x) < 0$ for $x > c$, then $x = c$ is a local maximum.


The Second Derivative Test

- ▶ If $f'(c) = 0$, and $f''(c) > 0$, then $x = c$ is a local minimum.
- ▶ If $f'(c) = 0$, and $f''(c) < 0$, then $x = c$ is a local maximum.

USING THE DERIVATIVE

EXAMPLES


Find all critical points and inflection points of the following function



USING THE DERIVATIVE

EXAMPLES


Find all critical points and inflection points of the following function



USING THE DERIVATIVE

EXAMPLES

Find all critical points and inflection points of the following function




USING THE DERIVATIVE

EXAMPLES

Example

The figure below is a graph of $f'(x)$. Find the x -values that are critical points of the function f itself. Are they local maxima, local minima, or neither?




USING THE DERIVATIVE

EXAMPLES

Example

The figure below is a graph of $f'(x)$. Find the x -values that are critical points of the function f itself. Are they local maxima, local minima, or neither?




We have two critical points ($f'(x) = 0$):

USING THE DERIVATIVE

EXAMPLES

Example

The figure below is a graph of $f'(x)$. Find the x -values that are critical points of the function f itself. Are they local maxima, local minima, or neither?



We have two critical points ($f'(x) = 0$):


- One at $c = -1.5$. Note how $f'(x) < 0$ for $x < -1.5$ and $f'(x) > 0$ for $x > -1.5$. It must be a minimum at $c = -1.5$.

USING THE DERIVATIVE

EXAMPLES

Example

The figure below is a graph of $f'(x)$. Find the x -values that are critical points of the function f itself. Are they local maxima, local minima, or neither?



We have two critical points ($f'(x) = 0$):

- ▶ One at $c = -1.5$. Note how $f'(x) < 0$ for $x < -1.5$ and $f'(x) > 0$ for $x > -1.5$. It must be a local minimum at $c = -1.5$.
- ▶ Another at $c = 1$. Note how $f'(x) > 0$ both before and after $c = 1$. This point is neither maximum nor minimum.

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

We need to evaluate $f''(1)$.

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

We need to evaluate $f''(1)$.

$$f'(x) = 4x^3 - 21x^2 + 17$$

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

We need to evaluate $f''(1)$.

$$f'(x) = 4x^3 - 21x^2 + 17$$

$$f''(x) = 12x^2 - 42x$$

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

We need to evaluate $f''(1)$.

$$f'(x) = 4x^3 - 21x^2 + 17$$

$$f''(x) = 12x^2 - 42x$$

$$f''(1) = 12 - 42 = -30 < 0$$

USING THE DERIVATIVE

EXAMPLES

Example

The function $f(x) = x^4 - 7x^3 + 17x$ has a critical point at $x = 1$. Use the second derivative test to identify it as a local maximum or local minimum.

We need to evaluate $f''(1)$.

$$f'(x) = 4x^3 - 21x^2 + 17$$

$$f''(x) = 12x^2 - 42x$$

$$f''(1) = 12 - 42 = -30 < 0$$

Therefore, $x = 1$ is a local maximum of f .

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function


$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

We have $f'(x) = 6x^2 + 6x - 180$; therefore, the critical points are

$$x = \frac{-6 \pm \sqrt{36 - 4 \cdot 6 \cdot (-180)}}{12} = \{-6, 5\}$$

To decide whether they are local max or min, we will make use of a sign chart:



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180$$

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

$$f''(x) = 0$$

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.

To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

$$f''(x) = 0, \quad 12x + 6 = 0$$

USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.


To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

$$f''(x) = 0, \quad 12x + 6 = 0, \quad x = \frac{-6}{12} = -0.5$$

To decide whether this is an actual inflection point, we need another sign chart!



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.


To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

$$f''(x) = 0, \quad 12x + 6 = 0, \quad x = \frac{-6}{12} = -0.5$$

To decide whether this is an actual inflection point, we need another sign chart!



USING THE DERIVATIVE

EXAMPLES

Example

Use the first derivative to find all critical points, and use the second derivative to find all inflection points of the function

$$f(x) = 2x^3 + 3x^2 - 180x + 3.$$

Identify each critical point as a local maximum, a local minimum, or neither.


To look for inflection points, we need to compute the second derivative:

$$f(x) = 2x^3 + 3x^2 - 180x + 3, \quad f'(x) = 6x^2 + 6x - 180, \quad f''(x) = 12x + 6$$

The inflection points are among the zeros of the second derivative:

$$f''(x) = 0, \quad 12x + 6 = 0, \quad x = \frac{-6}{12} = -0.5$$

To decide whether this is an actual inflection point, we need another sign chart!



USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$f'(x) = 2x + a$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$f'(x) = 2x + a$$

$$f'(6) = 0, \qquad 2 \cdot 6 + a = 0$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$f'(x) = 2x + a$$

$$f'(6) = 0, \qquad 2 \cdot 6 + a = 0, \qquad a = -12$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$\begin{aligned} f'(x) &= 2x + a \\ f'(6) &= 0, \qquad 2 \cdot 6 + a = 0, \qquad a = -12 \end{aligned}$$

Two conditions are enough to find the value of two unknowns:

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$\begin{aligned} f'(x) &= 2x + a \\ f'(6) &= 0, \qquad 2 \cdot 6 + a = 0, \qquad a = -12 \end{aligned}$$

Two conditions are enough to find the value of two unknowns:

$$\begin{cases} 6a + b = -34 \\ a = -12 \end{cases}$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

If the point $(6, 2)$ is in the graph of f , it must be $f(6) = 2$. This gives us a first condition on the unknown parameters a and b :

$$f(6) = 2, \qquad 6^2 + 6a + b = 2, \qquad 6a + b = -34$$

If the point $(6, 2)$ is to be a minimum, in particular it must be a critical point: this gives a second condition on a and b : $f'(6) = 0$

$$f'(x) = 2x + a$$

$$f'(6) = 0, \qquad 2 \cdot 6 + a = 0, \qquad a = -12$$

Two conditions are enough to find the value of two unknowns:

$$\begin{cases} 6a + b = -34 \\ a = -12 \end{cases} \qquad \begin{cases} a = -12 \\ b = -34 + 6 \cdot 12 = 38 \end{cases}$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

The equation of the parabola is then $f(x) = x^2 - 12x + 38$. It is easy to check that $(6, 2)$ is indeed a minimum (just in case!)

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

The equation of the parabola is then $f(x) = x^2 - 12x + 38$. It is easy to check that $(6, 2)$ is indeed a minimum (just in case!)

$$f(x) = x^2 - 12x + 38$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

The equation of the parabola is then $f(x) = x^2 - 12x + 38$. It is easy to check that $(6, 2)$ is indeed a minimum (just in case!)

$$f(x) = x^2 - 12x + 38, \quad f'(x) = 2x - 12$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

The equation of the parabola is then $f(x) = x^2 - 12x + 38$. It is easy to check that $(6, 2)$ is indeed a minimum (just in case!)

$$f(x) = x^2 - 12x + 38, \quad f'(x) = 2x - 12, \quad f''(x) = 2$$

USING THE DERIVATIVE

EXAMPLES

Example

Find constant a and b so that the minimum of the parabola $f(x) = x^2 + ax + b$ is at the point $(6, 2)$.

The equation of the parabola is then $f(x) = x^2 - 12x + 38$. It is easy to check that $(6, 2)$ is indeed a minimum (just in case!)

$$f(x) = x^2 - 12x + 38, \quad f'(x) = 2x - 12, \quad f''(x) = 2$$

If $(6, 2)$ is to be a minimum, it should be $f''(6) > 0$. This is obviously satisfied. We are good.

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8,$$

$$a(5 - b \ln 5) = 8$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right)$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0, \qquad a\left(1 - \frac{b}{5}\right) = 0$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0, \qquad a\left(1 - \frac{b}{5}\right) = 0$$

Let us solve for a and b . The second condition looks easier: it can only be $a = 0$ or $b = 5$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0, \qquad a\left(1 - \frac{b}{5}\right) = 0$$

Let us solve for a and b . The second condition looks easier: it can only be $a = 0$ or $b = 5$

- If $a = 0$, then the first condition is not satisfied, so we need to disregard this option.

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0, \qquad a\left(1 - \frac{b}{5}\right) = 0$$

Let us solve for a and b . The second condition looks easier: it can only be $a = 0$ or $b = 5$

- ▶ If $a = 0$, then the first condition is not satisfied, so we need to disregard this option.
- ▶ If $b = 5$, the first condition turns into

$$a(5 - 5 \ln 5) = 8$$

USING THE DERIVATIVE

EXAMPLES

Example

For what values of a and b does $f(x) = a(x - b \ln x)$ have a local extremum at the point $(5, 8)$?

We need two conditions, because we have two unknowns. The first comes from the fact that the point $(5, 8)$ is in the graph of f :

$$f(5) = 8, \qquad a(5 - b \ln 5) = 8$$

The second comes from the fact that $(5, 8)$ is a critical point: $f'(5) = 0$

$$f'(x) = a\left(1 - \frac{b}{x}\right), \qquad f'(5) = 0, \qquad a\left(1 - \frac{b}{5}\right) = 0$$

Let us solve for a and b . The second condition looks easier: it can only be $a = 0$ or $b = 5$

- ▶ If $a = 0$, then the first condition is not satisfied, so we need to disregard this option.
- ▶ If $b = 5$, the first condition turns into

$$a(5 - 5 \ln 5) = 8, \qquad a = \frac{8}{5 - 5 \ln 5} \approx -2.625369982$$