Doubling time, half-life

Francisco Blanco-Silva

University of South Carolina



- ► Functions
 - x and y intercepts (f(x) = 0, f(0))

$$\Delta y = f(b) - f(a)$$

Average Rate of Change from x = a to x = b

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

▶ Relative Change from x = a to x = b

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

► Linear Functions:

$$f(x) = b + mx$$

Exponential Functions $P_0a^t = P_0(1+r)^t = P_0e^{kt}$

► Functions

- x and y intercepts (f(x) = 0, f(0))
- ightharpoonup Change from x = a to x = b

$$\Delta y = f(b) - f(a)$$

 Average Rate of Change from x = a to x = b

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

▶ Relative Change from x = a to x = b

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

- ► Linear Functions: f(x) = b + mx
- ► Exponential Functions $P_0a^t = P_0(1+r)^t = P_0e^{kt}$

$$a = 1 + r$$
 $r = a - 1$

► Functions

- x and y intercepts (f(x) = 0, f(0))

$$\Delta y = f(b) - f(a)$$

Average Rate of Change from x = a to x = b

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

▶ Relative Change from x = a to x = b

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

- ► Linear Functions: f(x) = b + mx
- Exponential Functions $P_0a^t = P_0(1+r)^t = P_0e^{kt}$

$$a = 1 + r$$
 $r = a - 1$

$$k = \ln a$$
 $a = e^k$

► Functions

- ► x and y intercepts (f(x) = 0, f(0))

$$\Delta y = f(b) - f(a)$$

 Average Rate of Change from x = a to x = b

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

▶ Relative Change from x = a to x = b

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

- ► Linear Functions:
 - f(x) = b + mx
- Exponential Functions $P_0a^t = P_0(1+r)^t = P_0e^{kt}$

$$a = 1 + r \qquad r = a - 1$$

$$k = \ln a$$
 $a = e^k$
 $r = e^k - 1$ $k = \ln(1 + r)$

HOW DO WE RECOGNIZE IT?

What do we know?

▶ If the function is in the form $P = P_0 a^t$, a > 1 gives exponential growth, and a < 1 gives exponential decay.

HOW DO WE RECOGNIZE IT?

▶ If the function is in the form $P = P_0 a^t$, a > 1 gives exponential growth, and a < 1 gives exponential decay.

Doubling time, half-life

► If the function is in the form $P = P_0(1 + r)^t$, r > 0 gives exponential growth, and r < 0 gives exponential decay.

HOW DO WE RECOGNIZE IT?

▶ If the function is in the form $P = P_0 a^t$, a > 1 gives exponential growth, and a < 1 gives exponential decay.

Doubling time, half-life

- ▶ If the function is in the form $P = P_0(1+r)^t$, r > 0 gives exponential growth, and r < 0 gives exponential decay.
- ► If the function is in the form $P = P_0 e^{kt}$, k > 0 gives exponential growth, and k < 0 gives exponential decay.

EXAMPLES

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

EXAMPLES

Example

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to find a function that models this situation. The problem suggests that we need an exponential function, for which we know its continuous decay rate (k = -0.04) and its initial value ($P_0 = 2.4$)

EXAMPLES

Example

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to find a function that models this situation. The problem suggests that we need an exponential function, for which we know its continuous decay rate (k = -0.04) and its initial value ($P_0 = 2.4$)

$$P = P_0 e^{kt} = 2.4 e^{-0.04t}$$
 (in milirems/hour) $t = \text{hours after}$

EXAMPLES

Example

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to find a function that models this situation. The problem suggests that we need an exponential function, for which we know its continuous decay rate (k = -0.04) and its initial value ($P_0 = 2.4$)

$$P = P_0 e^{kt} = 2.4 e^{-0.04t}$$
 (in milirems/hour) $t = \text{hours after}$

The first question is asking for the radiation level after 24 hours:

$$P(24) = 2.4e^{-0.04 \times 24} = 0.9189429264$$
 milirems/hour

EXAMPLES

What do we know?

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to solve for t in the equation P(t) = 0.6

$$P(t) = 0.6$$

$$2.4e^{-0.04t} = 0.6$$

$$e^{-0.04t} = \frac{0.6}{2.4} = 0.25$$

EXAMPLES

Example

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to solve for t in the equation P(t) = 0.6

$$P(t) = 0.6$$

$$2.4e^{-0.04t} = 0.6$$

$$e^{-0.04t} = \frac{0.6}{2.4} = 0.25$$

$$\ln e^{-0.04t} = \ln 0.25$$

EXAMPLES

What do we know?

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to solve for t in the equation P(t) = 0.6

$$P(t) = 0.6$$

$$2.4e^{-0.04t} = 0.6$$

$$e^{-0.04t} = \frac{0.6}{2.4} = 0.25$$

$$\ln e^{-0.04t} = \ln 0.25$$

$$-0.04t = -1.386294361$$

EXAMPLES

The EPA investigated a spill of radioactive iodine. The radiation level at site was 2.4 milirems/hour. The level of radiation decays at a continuous hourly rate of k = -0.04.

- ▶ What was the level of radiation 24 hours later?
- ► Find the number of hours until the level of radiation reached the maximum acceptable limit (0.6 milirems/hour).

We need to solve for t in the equation P(t) = 0.6

-0.04t = -1.386294361

$$P(t) = 0.6$$

 $2.4e^{-0.04t} = 0.6$
 $e^{-0.04t} = \frac{0.6}{2.4} = 0.25$ $t = \frac{-1.386294361}{-0.04} \approx 34.65735903 \text{ hours.}$
 $\ln e^{-0.04t} = \ln 0.25$

Definition

Definition

$$P(t) = 2P_0$$

Definition

$$P(t) = 2P_0$$

$$P_0e^{kt}=2P_0$$

Doubling-time, Half-life

Definition

$$P(t) = 2P_0$$
$$P_0 e^{kt} = 2P_0$$

$$e^{kt}=2$$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$ $P_0 e^{kt} = 2P_0$ $e^{kt} = 2$

Doubling time, half-life

Definition

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$

Doubling-time, Half-life

Definition

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$ $P_0 e^{kt} = 2P_0$ $kt = \ln 2$ $t = \frac{\ln 2}{k}$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0$$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0$$
$$P_0e^{kt} = \frac{1}{2}P_0$$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0$$

$$P_0e^{kt} = \frac{1}{2}P_0$$

$$e^{kt} = \frac{1}{2}$$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0$$
 $\ln e^{kt} = \ln \frac{1}{2}$ $P_0 e^{kt} = \frac{1}{2}P_0$ $e^{kt} = \frac{1}{2}$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0$$
 $\ln e^{kt} = \ln \frac{1}{2}$ $P_0e^{kt} = \frac{1}{2}P_0$ $kt = -\ln 2$ $e^{kt} = \frac{1}{2}$

Definition

The doubling-time of an exponentially increasing quantity is the time required for the quantity to double.

$$P(t) = 2P_0$$
 $\ln e^{kt} = \ln 2$
 $P_0 e^{kt} = 2P_0$ $kt = \ln 2$
 $e^{kt} = 2$ $t = \frac{\ln 2}{k}$

$$P(t) = \frac{1}{2}P_0 \qquad \qquad \ln e^{kt} = \ln \frac{1}{2}$$

$$P_0 e^{kt} = \frac{1}{2}P_0 \qquad \qquad kt = -\ln 2$$

$$e^{kt} = \frac{1}{2} \qquad \qquad t = \frac{-\ln 2}{k}$$

Doubing-time, Half-life

EXAMPLES

Example

The quantity of ozone, Q, is decaying exponentially at a continuous rate of 0.25% per year. What is the half-life of the ozone?

EXAMPLES

Example

The quantity of ozone, *Q*, is decaying exponentially at a continuous rate of 0.25% per year. What is the half-life of the ozone?

We use the formula directly. We only need to know the value of the continuous decay rate, k, which is given to us: k = -0.0025. It is then

$$t = \frac{-\ln 2}{k}$$

EXAMPLES

Example

The quantity of ozone, Q, is decaying exponentially at a continuous rate of 0.25% per year. What is the half-life of the ozone?

We use the formula directly. We only need to know the value of the continuous decay rate, k, which is given to us: k = -0.0025. It is then

$$t = \frac{-\ln 2}{k} = \frac{-\ln 2}{-0.0025} \approx 277.25887224$$
 years.

COMPOUND INTEREST

An amount of money, P_0 , is deposited in an account paying interest at a rate of r per year. Let P be the balance in the account after t years.

- ▶ If the interest is compounded annually, then $P = P_0(1 + r)^t$.
- ▶ If the interest is compounded continuously, then $P = P_0e^{rt}$.

An amount of money, P_0 , is deposited in an account paying interest at a rate of r per year. Let P be the balance in the account after t years.

Doubling time, half-life

- ▶ If the interest is compounded annually, then $P = P_0(1 + r)^t$.
- ▶ If the interest is compounded continuously, then $P = P_0e^{rt}$.

Example

A bank advertises an interest rate of 8% per year. If you deposit \$5,000, how much is in the account 3 years later if the interest is compounded...

- ► Annually?
- ► Continuously?

COMPOUND INTEREST

An amount of money, P_0 , is deposited in an account paying interest at a rate of r per year. Let P be the balance in the account after t years.

- ▶ If the interest is compounded annually, then $P = P_0(1 + r)^t$.
- ▶ If the interest is compounded continuously, then $P = P_0e^{rt}$.

Example

A bank advertises an interest rate of 8% per year. If you deposit \$5,000, how much is in the account 3 years later if the interest is compounded...

- ► Annually?
- ► Continuously?
- ▶ We use the formula $P(t) = P_0(1+r)^t$ here, with r = 0.08, and $P_0 = 5000$. We need to compute P(3)

$$P(3) = 5000(1 + 0.08)^3 = $6298.56$$

An amount of money, P_0 , is deposited in an account paying interest at a rate of r per year. Let P be the balance in the account after t years.

- ▶ If the interest is compounded annually, then $P = P_0(1 + r)^t$.
- ▶ If the interest is compounded continuously, then $P = P_0e^{rt}$.

Example

A bank advertises an interest rate of 8% per year. If you deposit \$5,000, how much is in the account 3 years later if the interest is compounded...

- ► Annually?
- ► Continuously?
- ▶ We use the formula $P(t) = P_0(1+r)^t$ here, with r = 0.08, and $P_0 = 5000$. We need to compute P(3)

$$P(3) = 5000(1 + 0.08)^3 = $6298.56$$

▶ We use the formula $P(t) = P_0 e^{rt}$ for the same values of r and P_0

$$P(3) = 5000e^{0.08 \times 3} = $6356.24575$$

Doubling time, half-life

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

Doubling time, half-life

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$

$$10000e^{0.05t} = 15000$$

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$

$$10000e^{0.05t} = 15000$$

$$e^{0.05t} = \frac{15000}{10000}$$

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$

$$10000e^{0.05t} = 15000$$

$$e^{0.05t} = \frac{15000}{10000}$$

$$\ln e^{0.05t} = \ln 1.5$$

What do we know?

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $10000e^{0.05t} = 15000$ $e^{0.05t} = \frac{15000}{10000}$ $\ln e^{0.05t} = \ln 1.5$

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $10000e^{0.05t} = 15000$ $t = \frac{0.4054651081}{0.05}$ $e^{0.05t} = \frac{15000}{10000}$ $\ln e^{0.05t} = \ln 1.5$

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $t = \frac{0.4054651081}{0.05}$ $t \approx 8.109302162 \text{ years}$ $\ln e^{0.05t} = \ln 1.5$

EXAMPLES

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

We have to solve for t in the equation P(t) = 15000, where $P = P_0 e^{rt}$. They are giving us $P_0 = 10000$ and r = 0.05:

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $t = \frac{0.4054651081}{0.05}$ $t \approx 8.109302162 \text{ years}$ $\ln e^{0.05t} = \ln 1.5$

That is a little bit more than eight years and one month (1.31163 months)

EXAMPLES

What do we know?

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

We have to solve for t in the equation P(t) = 15000, where $P = P_0 e^{rt}$. They are giving us $P_0 = 10000$ and r = 0.05:

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $t = \frac{0.4054651081}{0.05}$ $t \approx 8.109302162 \text{ years}$ $\ln e^{0.05t} = \ln 1.5$

That is a little bit more than eight years and one month (1.31163 months) Or better, eight years and almost six weeks (5.70311 weeks)

What do we know?

Future Value

If \$10,000 is deposited in an account paying interest at a rate of 5% per year, compounded continuously, how long does it take for the balance in the account to reach \$15,000?

We have to solve for t in the equation P(t) = 15000, where $P = P_0 e^{rt}$. They are giving us $P_0 = 10000$ and r = 0.05:

$$P(t) = 15000$$
 $0.05t = 0.4054651081$ $t = \frac{0.4054651081}{0.05}$ $t \approx 8.109302162 \text{ years}$ $\ln e^{0.05t} = \ln 1.5$

That is a little bit more than eight years and one month (1.31163 months) Or better, eight years and almost six weeks (5.70311 weeks) Or even better, eight years almost 40 days (39.9218 days)

EXAMPLES

Future Value

You win the lottery and are offered the choice between \$1 million in four yearly installments of \$250,000 each, starting now, or a lump-sum payment of \$920,000 now.

Doubling time, half-life

Assuming a 6% interest rate per year, compounded continuously, and ignoring taxes, which should you choose?

What do we know?

Future Value

You win the lottery and are offered the choice between \$1 million in four yearly installments of \$250,000 each, starting now, or a lump-sum payment of \$920,000 now.

Doubling time, half-life

Assuming a 6% interest rate per year, compounded continuously, and ignoring taxes, which should you choose?

The following table summarizes the two situations:

	2015	2016	2017	2018
First option	250,000			
		+250,000	+250,000	+250,000
Second option	920,000			

EXAMPLES

	2015	2016	2017	2018
First option	250,000	265459		
		+250,000	+250,000	+250,000
		=515459		
Second option	920,000			

Let us examine the first option.

▶ In one year, the initial amount of \$250,000 gives us $P(1) = 250000e^{0.06 \times 1} = 265459.13675$ after compounding at 6%. Then we add \$250,000 more, for a total of 265459.13675 + 250000 = 515459.13675.

	2015	2016	2017	2018
First option	250,000	265459	547333	
		+250,000	+250,000	+250,000
		=515459	= 797333	
Second option	920,000			

Let us examine the first option.

- ▶ In one year, the initial amount of \$250,000 gives us $P(1) = 250000e^{0.06 \times 1} = 265459.13675$ after compounding at 6%. Then we add \$250,000 more, for a total of 265459.13675 + 250000 = 515459.13675.
- ► After another year, this amount increases to $P(1) = 515459.13675e^{0.06} = 547333.34989$. We add \$250,000 again, to obtain 547333.34989 + 250000 = 797333.34989.

EXAMPLES

	2015	2016	2017	2018
First option	250,000	265459	547333	846637
		+250,000	+250,000	+250,000
		=515459	= 797333	=1096637
Second option	920,000			

Let us examine the first option.

- ▶ In one year, the initial amount of \$250,000 gives us $P(1) = 250000e^{0.06 \times 1} = 265459.13675$ after compounding at 6%. Then we add \$250,000 more, for a total of 265459.13675 + 250000 = 515459.13675.
- ► After another year, this amount increases to $P(1) = 515459.13675e^{0.06} = 547333.34989$. We add \$250,000 again, to obtain 547333.34989 + 250000 = 797333.34989.
- ► After another year, this amounts increases to $P(1) = 797333.34989e^{0.06} = 846637.69106$. At that moment, we add the last \$250,000 to get the final amount of 846637.69106 + 250000 = 1096637.69106

	2015	2016	2017	2018
First option	250,000	265459	547333	846637
_		+250,000	+250,000	+250,000
				=1096637
Second option	920,000			1101439

Let us examine the second option. We only need to worry about how much we have after 3 years:

$$P(3) = 920000e^{0.06 \times 3} = 1101439.974$$

Which option do you prefer?