
Lesson 21: Systems of Differential Equations: Introduction. Reduction to first-order systems.

Francisco Blanco-Silva

University of South Carolina



WHAT DO WE KNOW?

- The concepts of differential equation and initial value problem
- The concept of order of a differential equation.
- ► The concepts of general solution, particular solution and singular solution.
- ► Slope fields
- Approximations to solutions via Euler's Method and Improved Euler's Method

- ► First-Order Differential Equations
 - Separable equations
 - Homogeneous First-Order Equations
 - ► Linear First-Order Equations
 - ▶ Bernoulli Equations
 - General Substitution Methods
 - ► Exact Equations

Coefficients

- ► Second-Order Differential Equations
 - ► Reducible Equations
 - ► General Linear Equations (Intro)
 - ► Linear Equations with Constant
 - ► Characteristic Equation
 - ► Variation of Parameters
 - Undetermined Coefficients

What do we know?

LAPLACE TRANSFORMS

f(x)	$\mathcal{L}{f} = \int_0^\infty e^{-sx} f(x) dx$		f(x)	$\mathcal{L}{f} = \int_0^\infty e^{-sx} f(x) dx$	
1	$\frac{1}{s}$	s > 0	$cf(x) \pm g(x)$	$cF(s) \pm G(s)$	s > max(a, b)
x^p	$\frac{\Gamma(p+1)}{s^{p+1}}$	<i>s</i> > 0	$x^n f(x)$	$(-1)^n F^{(n)}$	s > a
x^n	$\frac{n!}{s^{n+1}}$	s > 0	$e^{\alpha x}f(x)$	$F(s-\alpha)$	$s > a + \alpha$
$e^{\alpha x}$	$\frac{1}{s-\alpha}$	$s > \alpha$	$\frac{f(x)}{x}$	$\int_{s}^{\infty} F(\sigma) d\sigma$	s > a
$\sin \beta x$	$\frac{\beta}{s^2 + \beta^2}$	s > 0	$f \star g$	F(s)G(s)	$s > \max(a, b)$
$\cos \beta x$	$\frac{s}{s^2 + \beta^2}$	s > 0	f'(x)	sF(s) - f(0)	s > a

DEFINITION

Systems of Differential Equations

A system of differential equations is a collection of functions and functional equations of the form

$$\begin{cases} y_1^{(n)} = F_1(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ y_2^{(n)} = F_2(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ \dots \\ y_r^{(n)} = F_r(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \end{cases}$$

order n,r functions

Systems of Differential Equations

DEFINITION

A system of differential equations is a collection of functions and functional equations of the form

$$\begin{cases} y_1^{(n)} = F_1(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ y_2^{(n)} = F_2(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ \dots \\ y_r^{(n)} = F_r(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \end{cases}$$

order n,r functions

Some examples

$$\begin{cases} y_1' = x + y_1 - y_2 \\ y_2' = 2y_1 - 3y_2 \end{cases}$$
first order, two functions

Systems of Differential Equations

DEFINITION

A system of differential equations is a collection of functions and functional equations of the form

$$\begin{cases} y_1^{(n)} = F_1(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ y_2^{(n)} = F_2(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \\ \dots \\ y_r^{(n)} = F_r(x, y_1, y_1', \dots, y_1^{(n-1)}, y_2, y_2', \dots, y_2^{(n-1)}, \dots, y_r, y_r', \dots, y_r^{(n-1)}) \end{cases}$$

order n,r functions

Some examples

$$\begin{cases} y_1' = x + y_1 - y_2 \\ y_2' = 2y_1 - 3y_2 \end{cases}$$
first order, two functions
$$\begin{cases} y_1'' = y_2 + y_3' \\ y_2'' = y_3 - y_1' \\ y_3'' = y_1 + y_2' \end{cases}$$

$$\begin{cases} y_1'' = y_2 + y_3' \\ y_2'' = y_3 - y_1' \\ y_3'' = y_1 + y_2' \end{cases}$$

second order, three functions

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Any differential equation of order n can be transformed into a system of n differential equations of first order:

$$y^{(n)} = F(x, y, y', y'', \dots, y^{(n-1)})$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Any differential equation of order n can be transformed into a system of n differential equations of first order:

$$y^{(n)} = F(x, y, y', y'', \dots, y^{(n-1)})$$

We start by assigning $y_1 = y$ and, to each k-th derivative of y, the function y_{k+1} :

$$y_1 = y$$
, $y_2 = y'$, $y_3 = y''$, ... $y_n = y^{(n-1)}$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Any differential equation of order n can be transformed into a system of n differential equations of first order:

$$y^{(n)} = F(x, y, y', y'', \dots, y^{(n-1)})$$

We start by assigning $y_1 = y$ and, to each k-th derivative of y, the function y_{k+1} :

$$y_1 = y$$
, $y_2 = y'$, $y_3 = y''$, ... $y_n = y^{(n-1)}$

Note that now, $y'_1 = y' = y_2$, $y'_2 = y_3$ and, in general, $y'_k = y_{k+1}$. We have the system we required:

$$\begin{cases} y'_1 = y_2 \\ y'_2 = y_3 \\ \dots \\ y'_{n-1} = y'_n \\ y'_n = y^{(n)} = F(x, y, y', y'', \dots, y^{(n-1)}) = F(x, y_1, y_2, \dots, y_n) \end{cases}$$

Systems of Differential Equations

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

We start with the assignment of functions first. We stop at three (the order of the differential equation)

$$y_1 = y, \qquad y_2 = y', \qquad y_3 = y''$$

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

We start with the assignment of functions first. We stop at three (the order of the differential equation)

$$y_1 = y, \qquad y_2 = y', \qquad y_3 = y''$$

We take derivatives now.

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

We start with the assignment of functions first. We stop at three (the order of the differential equation)

$$y_1=y, \qquad y_2=y', \qquad y_3=y''$$

We take derivatives now.

$$\begin{cases} y_1' = y_2 \end{cases}$$

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

We start with the assignment of functions first. We stop at three (the order of the differential equation)

$$y_1 = y, \qquad y_2 = y', \qquad y_3 = y''$$

We take derivatives now.

$$\begin{cases} y_1' = y_2 \\ y_2' = y_3 \end{cases}$$

EXAMPLES

Transform the differential equation in a first-order system of differential Equations

$$y''' + 3y'' + 2y' - 5y = \sin 2x$$

We start with the assignment of functions first. We stop at three (the order of the differential equation)

$$y_1 = y, \qquad y_2 = y', \qquad y_3 = y''$$

We take derivatives now. Note that $y_3' = y''' = \sin 2x + 5y - 2y' - 3y''$

$$\begin{cases} y_1' = y_2 \\ y_2' = y_3 \\ y_3' = \sin 2x + 5y_1 - 2y_2 - 3y_3 \end{cases}$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

$$(Y_1' = Y_2)$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

$$\begin{cases} Y_1' = Y_2 \\ Y_2' = -3Y_1 + Y_3 \end{cases}$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

$$\begin{cases} Y_1' = Y_2 \\ Y_2' = -3Y_1 + Y_3 \\ Y_3' = Y_4 \end{cases}$$

TRANSFORMATION TO FIRST-ORDER SYSTEMS

Using the same technique, it is also possible to transform any system of differential equations into another of order one.

Transform the following system to First-order

$$\begin{cases} y_1'' = -3y_1 + y_2 \\ y_2'' = 2y_1 - 2y_2 + 40\sin 3x \end{cases}$$

$$\begin{cases} Y_1' = Y_2 \\ Y_2' = -3Y_1 + Y_3 \\ Y_3' = Y_4 \\ Y_4' = 2Y_1 - 2Y_3 + 40\sin 3x \end{cases}$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

Note that

$$y_1^{\prime\prime}=-2y_2^{\prime}$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

Note that

$$y_1'' = -2y_2' = -2(\frac{1}{2}y_1) = -y_1$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

Note that

$$y_1'' = -2y_2' = -2(\frac{1}{2}y_1) = -y_1$$

We have then the second order homogeneous linear equation with constant coefficients $y_1'' + y_1 = 0$, with general solution

$$y_1 = A\cos x + B\sin x$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

Note that

$$y_1'' = -2y_2' = -2(\frac{1}{2}y_1) = -y_1$$

We have then the second order homogeneous linear equation with constant coefficients $y_1'' + y_1 = 0$, with general solution

$$y_1 = A\cos x + B\sin x$$

It must be then

$$y_2 = -\frac{1}{2}y_1'$$

EXAMPLES

It is often possible to solve a first-order system of differential equations by transforming it to a single differential equation.

Transform the following system into a differential equation, and solve it

$$\begin{cases} y_1' = -2y_2 \\ y_2' = \frac{1}{2}y_1 \end{cases}$$

Note that

$$y_1'' = -2y_2' = -2(\frac{1}{2}y_1) = -y_1$$

We have then the second order homogeneous linear equation with constant coefficients $y_1'' + y_1 = 0$, with general solution

$$y_1 = A\cos x + B\sin x$$

It must be then

$$y_2 = -\frac{1}{2}y_1' = \frac{A}{2}\sin x - \frac{B}{2}\cos x$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2 = 2y_1 + y_1'$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2 = 2y_1 + y_1'$$

We have the second-order homogeneous linear equation with constant coefficients

$$y_1'' - y_1' - 2y_1 = 0$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2 = 2y_1 + y_1'$$

We have the second-order homogeneous linear equation with constant coefficients

$$y_1'' - y_1' - 2y_1 = 0$$

The general solution is readily found from the roots of the characteristic equation:

$$r^{2} - r - 2 = 0,$$
 $r = \frac{1 \pm \sqrt{1 - 4 \cdot (-2)}}{2} = \frac{1 \pm 3}{2} = \{2, -1\}$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2 = 2y_1 + y_1'$$

We have the second-order homogeneous linear equation with constant coefficients

$$y_1'' - y_1' - 2y_1 = 0$$

The general solution is readily found from the roots of the characteristic equation:

$$r^{2} - r - 2 = 0,$$
 $r = \frac{1 \pm \sqrt{1 - 4 \cdot (-2)}}{2} = \frac{1 \pm 3}{2} = \{2, -1\}$

We have

$$y_1 = Ae^{-x} + Be^{2x}$$

EXAMPLES

Transform the system into a differential equation, and solve it

$$\begin{cases} y_1' = y_2 \\ y_2' = 2y_1 + y_2 \end{cases}$$

As before, we start by taking derivatives of the easier equation:

$$y_1'' = y_2' = 2y_1 + y_2 = 2y_1 + y_1'$$

We have the second-order homogeneous linear equation with constant coefficients

$$y_1'' - y_1' - 2y_1 = 0$$

The general solution is readily found from the roots of the characteristic equation:

$$r^{2} - r - 2 = 0$$
, $r = \frac{1 \pm \sqrt{1 - 4 \cdot (-2)}}{2} = \frac{1 \pm 3}{2} = \{2, -1\}$

We have

$$y_1 = Ae^{-x} + Be^{2x}$$

 $y_2 = y_1' = -Ae^{-x} + 2Be^{2x}$