


Lesson 9: Rules of Differentiation—Power functions and Polynomials

Francisco Blanco-Silva

University of South Carolina



WHAT DO WE KNOW?

► Functions

- x - and y -**intercepts** ($f(x) = 0, f(0)$)
- **Change** from $x = a$ to $x = b$

$$\Delta y = f(b) - f(a)$$

- **Average Rate of Change** from $x = a$ to $x = b$

$$ARC = \frac{\Delta y}{\Delta x} = \frac{f(b) - f(a)}{b - a}$$

- **Relative Change** from $x = a$ to $x = b$

$$RC = \frac{\Delta y}{f(a)} = \frac{f(b) - f(a)}{f(a)}$$

- **Instantaneous Rate of Change** at $x = a$

$$f'(a)$$

► Linear Functions:

$$f(x) = b + mx$$

► Exponential Functions

$$P_0 a^t = P_0 (1 + r)^t = P_0 e^{kt}$$

► Power Functions

$$kx^p$$

► Polynomials

$$a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n$$

THE DERIVATIVE AS A FUNCTION

In the previous lesson we have seen that the derivative of a function $y = f(x)$ can be seen as a function itself: One that assigns, to each x , the value of the slope of the tangent line at the graph of f at the point $(x, f(x))$.

We denote this new function in two ways

$$f'(x) = \underbrace{\frac{df}{dx}}_{\text{Leibnitz notation}} \quad \begin{array}{l} \leftarrow \text{derivative of } f \\ \leftarrow \text{with respect to the variable } x \end{array}$$

THE DERIVATIVE AS A FUNCTION

In the previous lesson we have seen that the derivative of a function $y = f(x)$ can be seen as a function itself: One that assigns, to each x , the value of the slope of the tangent line at the graph of f at the point $(x, f(x))$.

We denote this new function in two ways

$$f'(x) = \underbrace{\frac{df}{dx}}_{\text{Leibnitz notation}} \quad \begin{array}{l} \leftarrow \text{derivative of } f \\ \leftarrow \text{with respect to the variable } x \end{array}$$

Write the Leibnitz notation for the derivative of the given function and include units.

An employee's pay, P , in dollars, for a week is a function of the number of hours worked, H .

THE DERIVATIVE AS A FUNCTION

In the previous lesson we have seen that the derivative of a function $y = f(x)$ can be seen as a function itself: One that assigns, to each x , the value of the slope of the tangent line at the graph of f at the point $(x, f(x))$.

We denote this new function in two ways

$$f'(x) = \underbrace{\frac{df}{dx}}_{\text{Leibnitz notation}} \quad \begin{array}{l} \leftarrow \text{derivative of } f \\ \leftarrow \text{with respect to the variable } x \end{array}$$

Write the Leibnitz notation for the derivative of the given function and include units.

An employee's pay, P , in dollars, for a week is a function of the number of hours worked, H .

The solution is

$$\frac{dP}{dH}$$

THE DERIVATIVE AS A FUNCTION

In the previous lesson we have seen that the derivative of a function $y = f(x)$ can be seen as a function itself: One that assigns, to each x , the value of the slope of the tangent line at the graph of f at the point $(x, f(x))$.

We denote this new function in two ways

$$f'(x) = \underbrace{\frac{df}{dx}}_{\text{Leibnitz notation}} \quad \begin{array}{l} \leftarrow \text{derivative of } f \\ \leftarrow \text{with respect to the variable } x \end{array}$$

Write the Leibnitz notation for the derivative of the given function and include units.

An employee's pay, P , in dollars, for a week is a function of the number of hours worked, H .

The solution is

$$\begin{array}{l} \frac{dP}{dH} \leftarrow \text{dollars} \\ \quad \quad \leftarrow \text{hours} \end{array}$$

THE DERIVATIVE AS A FUNCTION

In the previous lesson we have seen that the derivative of a function $y = f(x)$ can be seen as a function itself: One that assigns, to each x , the value of the slope of the tangent line at the graph of f at the point $(x, f(x))$.

We denote this new function in two ways

$$f'(x) = \underbrace{\frac{df}{dx}}_{\text{Leibnitz notation}} \quad \begin{array}{l} \leftarrow \text{derivative of } f \\ \leftarrow \text{with respect to the variable } x \end{array}$$

Write the Leibnitz notation for the derivative of the given function and include units.

An employee's pay, P , in dollars, for a week is a function of the number of hours worked, H .

The solution is

$$\begin{array}{l} \frac{dP}{dH} \leftarrow \text{dollars} \\ \quad \quad \leftarrow \text{hours} \end{array}$$

Therefore, the units are **dollars/hour**.

THE DERIVATIVE AS A FUNCTION

Interpretation of the Derivative

The statement $f'(a) = b$ means that if the independent variable x goes up from a to $a + 1$, then the dependent variable goes up or down by $|b|$ units.

THE DERIVATIVE AS A FUNCTION

Interpretation of the Derivative

The statement $f'(a) = b$ means that if the independent variable x goes up from a to $a + 1$, then the dependent variable goes up or down by $|b|$ units.

Example

An economist is interested in how the price of a certain item affects its sales. At a price of p dollars, a quantity q of the item is sold.

$$q = f(p)$$

- ▶ What does it mean $f(160) = 2050$?
- ▶ What does it mean $f'(160) = -25$?
- ▶ What does it mean $f'(30) = 49$?

THE DERIVATIVE AS A FUNCTION

Interpretation of the Derivative

The statement $f'(a) = b$ means that if the independent variable x goes up from a to $a + 1$, then the dependent variable goes up or down by $|b|$ units.

Example

An economist is interested in how the price of a certain item affects its sales. At a price of p dollars, a quantity q of the item is sold.

$$q = f(p)$$

- ▶ What does it mean $f(160) = 2050$?
If the price is \$160, then 2050 items will be sold.
- ▶ What does it mean $f'(160) = -25$?
- ▶ What does it mean $f'(30) = 49$?

THE DERIVATIVE AS A FUNCTION

Interpretation of the Derivative

The statement $f'(a) = b$ means that if the independent variable x goes up from a to $a + 1$, then the dependent variable goes up or down by $|b|$ units.

Example

An economist is interested in how the price of a certain item affects its sales. At a price of p dollars, a quantity q of the item is sold.

$$q = f(p)$$

- ▶ What does it mean $f(160) = 2050$?
If the price is \$160, then 2050 items will be sold.
- ▶ What does it mean $f'(160) = -25$?
If the price goes up from \$160 by \$1 per item, about 25 fewer items will be sold.
- ▶ What does it mean $f'(30) = 49$?

THE DERIVATIVE AS A FUNCTION

Interpretation of the Derivative

The statement $f'(a) = b$ means that if the independent variable x goes up from a to $a + 1$, then the dependent variable goes up or down by $|b|$ units.

Example

An economist is interested in how the price of a certain item affects its sales. At a price of p dollars, a quantity q of the item is sold.

$$q = f(p)$$

- ▶ What does it mean $f(160) = 2050$?
If the price is \$160, then 2050 items will be sold.
- ▶ What does it mean $f'(160) = -25$?
If the price goes up from \$160 by \$1 per item, about 25 fewer items will be sold.
- ▶ What does it mean $f'(30) = 49$?
If the price goes up from \$30 by \$1 per item, about 49 more items will be sold.

THE DERIVATIVE AS A FUNCTION

EXAMPLES

Example

The average value of corn production in Kenya, in Kenyan shillings, of the yearly maize production from an average plot of land is a function $y = f(x)$ of the quantity x of fertilizer used (in kilograms).

- ▶ Interpret the statements $f(5) = 11,500$ and $f'(5) = 350$.
- ▶ Use the statements to estimate $f(6)$ and $f(10)$.
- ▶ Which estimate in the previous part is more reliable?

THE DERIVATIVE AS A FUNCTION

EXAMPLES

Example

The average value of corn production in Kenya, in Kenyan shillings, of the yearly maize production from an average plot of land is a function $y = f(x)$ of the quantity x of fertilizer used (in kilograms).

- ▶ Interpret the statements $f(5) = 11,500$ and $f'(5) = 350$.
 - ▶ Use the statements to estimate $f(6)$ and $f(10)$.
 - ▶ Which estimate in the previous part is more reliable?
-
- ▶ $f(5) = 11,500$ means that if 5 kg of fertilizer are used on an average plot of land, the average value of corn production in Kenya will be 11,500KSh.

THE DERIVATIVE AS A FUNCTION

EXAMPLES

Example

The average value of corn production in Kenya, in Kenyan shillings, of the yearly maize production from an average plot of land is a function $y = f(x)$ of the quantity x of fertilizer used (in kilograms).

- ▶ Interpret the statements $f(5) = 11,500$ and $f'(5) = 350$.
 - ▶ Use the statements to estimate $f(6)$ and $f(10)$.
 - ▶ Which estimate in the previous part is more reliable?
-
- ▶ $f(5) = 11,500$ means that if 5 kg of fertilizer are used on an average plot of land, the average value of corn production in Kenya will be 11,500KSh.
 $f'(5) = 350$ means that, if the amount of fertilizer used on an average plot of land goes up to 6 kg, then the average value of corn production will **increase** by 350KSh more.

THE DERIVATIVE AS A FUNCTION

EXAMPLES

Example

The average value of corn production in Kenya, in Kenyan shillings, of the yearly maize production from an average plot of land is a function $y = f(x)$ of the quantity x of fertilizer used (in kilograms).

- ▶ Interpret the statements $f(5) = 11,500$ and $f'(5) = 350$.
 - ▶ Use the statements to estimate $f(6)$ and $f(10)$.
 - ▶ Which estimate in the previous part is more reliable?
-
- ▶ $f(5) = 11,500$ means that if 5 kg of fertilizer are used on an average plot of land, the average value of corn production in Kenya will be 11,500KSh.
 $f'(5) = 350$ means that, if the amount of fertilizer used on an average plot of land goes up to 6 kg, then the average value of corn production will **increase** by 350KSh more.
 - ▶ We may estimate then $f(6) \approx f(5) + 350 = 11,500 + 350 = 11,850$ KSh, and $f(10) \approx 11,500 + 5 \cdot 350 = 11,500 + 1,750 = 13,250$ KSh

THE DERIVATIVE AS A FUNCTION

EXAMPLES

Example

The average value of corn production in Kenya, in Kenyan shillings, of the yearly maize production from an average plot of land is a function $y = f(x)$ of the quantity x of fertilizer used (in kilograms).

- ▶ Interpret the statements $f(5) = 11,500$ and $f'(5) = 350$.
 - ▶ Use the statements to estimate $f(6)$ and $f(10)$.
 - ▶ Which estimate in the previous part is more reliable?
-
- ▶ $f(5) = 11,500$ means that if 5 kg of fertilizer are used on an average plot of land, the average value of corn production in Kenya will be 11,500KSh.
 $f'(5) = 350$ means that, if the amount of fertilizer used on an average plot of land goes up to 6 kg, then the average value of corn production will **increase** by 350KSh more.
 - ▶ We may estimate then $f(6) \approx f(5) + 350 = 11,500 + 350 = 11,850$ KSh, and $f(10) \approx 11,500 + 5 \cdot 350 = 11,500 + 1,750 = 13,250$ KSh
 - ▶ $f(6) \approx 11,850$ KSh

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

D4 The derivative of a subtraction is the subtraction of the derivatives:

$$h(x) = f(x) - g(x), \quad h'(x) = f'(x) - g'(x)$$

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

D4 The derivative of a subtraction is the subtraction of the derivatives:

$$h(x) = f(x) - g(x), \quad h'(x) = f'(x) - g'(x)$$

D5 The derivative of a scalar times a function is a scalar times the derivative of the function.

$$h(x) = c \cdot f(x), \quad h'(x) = c \cdot f'(x)$$

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

D4 The derivative of a subtraction is the subtraction of the derivatives:

$$h(x) = f(x) - g(x), \quad h'(x) = f'(x) - g'(x)$$

D5 The derivative of a scalar times a function is a scalar times the derivative of the function.

$$h(x) = c \cdot f(x), \quad h'(x) = c \cdot f'(x)$$

D6 The Power Rule

$$f(x) = x^n, \quad f'(x) = nx^{n-1}$$

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

D4 The derivative of a subtraction is the subtraction of the derivatives:

$$h(x) = f(x) - g(x), \quad h'(x) = f'(x) - g'(x)$$

D5 The derivative of a scalar times a function is a scalar times the derivative of the function.

$$h(x) = c \cdot f(x), \quad h'(x) = c \cdot f'(x)$$

D6 The Power Rule

$$f(x) = x^n, \quad f'(x) = nx^{n-1}$$

D7 The derivative of $f(x) = e^x$ is $f'(x) = e^x$.

RULES OF DIFFERENTIATION

D1 The derivative of a constant function is zero.

$$f(x) = c, \quad f'(x) = 0$$

D2 The derivative of $f(x) = x$ is $f'(x) = 1$.

D3 The derivative of a sum is the sum of the derivatives:

$$h(x) = f(x) + g(x), \quad h'(x) = f'(x) + g'(x)$$

D4 The derivative of a subtraction is the subtraction of the derivatives:

$$h(x) = f(x) - g(x), \quad h'(x) = f'(x) - g'(x)$$

D5 The derivative of a scalar times a function is a scalar times the derivative of the function.

$$h(x) = c \cdot f(x), \quad h'(x) = c \cdot f'(x)$$

D6 The Power Rule

$$f(x) = x^n, \quad f'(x) = nx^{n-1}$$

D7 The derivative of $f(x) = e^x$ is $f'(x) = e^x$.

D8 For any $a > 0$, the derivative of $f(x) = a^x$ is $f'(x) = a^x \ln a$.

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f(x) = x$$

$$f(x) = x + \pi$$

$$f(x) = 200x$$

$$f(x) = 45 - 5x$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f(x) = x + \pi$$

$$f(x) = 200x$$

$$f(x) = 45 - 5x$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f(x) = 200x$$

$$f(x) = 45 - 5x$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f(x) = 45 - 5x$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f'(x) = 3 \cdot \frac{1}{2}x^{1/2-1} = \frac{3}{2}x^{-1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f'(x) = 3 \cdot \frac{1}{2}x^{1/2-1} = \frac{3}{2}x^{-1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}} = x^{-7/2}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f'(x) = 3 \cdot \frac{1}{2}x^{1/2-1} = \frac{3}{2}x^{-1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}} = x^{-7/2}$$

$$f'(x) = -\frac{7}{2}x^{-7/2-1} = -\frac{7}{2}x^{-9/2}$$

$$f(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f'(x) = 3 \cdot \frac{1}{2}x^{1/2-1} = \frac{3}{2}x^{-1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}} = x^{-7/2}$$

$$f'(x) = -\frac{7}{2}x^{-7/2-1} = -\frac{7}{2}x^{-9/2}$$

$$f(x) = e^x$$

$$f'(x) = e^x$$

$$f(x) = 2^x$$

RULES OF DIFFERENTIATION

BASIC EXAMPLES

Find the derivative of the following functions:

$$f(x) = 5$$

$$f'(x) = 0$$

$$f(x) = x$$

$$f'(x) = 1$$

$$f(x) = x + \pi$$

$$f'(x) = 1 + 0 = 1$$

$$f(x) = 200x$$

$$f'(x) = 200 \cdot 1 = 200$$

$$f(x) = 45 - 5x$$

$$f'(x) = 0 - 5 \cdot 1 = -5$$

$$f(x) = x^5$$

$$f'(x) = 5x^{5-1} = 5x^4$$

$$f(x) = 45x^7$$

$$f'(x) = 45 \cdot 7x^{7-1} = 315x^6$$

$$f(x) = 2\pi - x^{3/2}$$

$$f'(x) = 0 - \frac{3}{2}x^{3/2-1} = -\frac{3}{2}x^{1/2}$$

$$f(x) = 3\sqrt{x} = 3x^{1/2}$$

$$f'(x) = 3 \cdot \frac{1}{2}x^{1/2-1} = \frac{3}{2}x^{-1/2}$$

$$f(x) = \frac{1}{\sqrt{x^7}} = x^{-7/2}$$

$$f'(x) = -\frac{7}{2}x^{-7/2-1} = -\frac{7}{2}x^{-9/2}$$

$$f(x) = e^x$$

$$f'(x) = e^x$$

$$f(x) = 2^x$$

$$f'(x) = 2^x \ln 2$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$f'(t) = 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$\begin{aligned} h(t) &= (t - 3t^2)(\sqrt{t} + 4) \\ &= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2 \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$\begin{aligned} h(t) &= (t - 3t^2)(\sqrt{t} + 4) \\ &= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2 \\ &= t^{3/2} + 4t - 3t^{5/2} - 12t^2 \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$h'(t) = \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$\begin{aligned} h'(t) &= \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1} \\ &= \frac{3}{2}t^{1/2} + 4 - \frac{15}{2}t^{3/2} - 24t \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$\begin{aligned} h'(t) &= \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1} \\ &= \frac{3}{2}t^{1/2} + 4 - \frac{15}{2}t^{3/2} - 24t \end{aligned}$$

$$\begin{aligned} h(t) &= \frac{4 + \sqrt{t}}{t^5} \\ &= \frac{4}{t^5} + \frac{t^{1/2}}{t^5} \end{aligned}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$\begin{aligned} h'(t) &= \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1} \\ &= \frac{3}{2}t^{1/2} + 4 - \frac{15}{2}t^{3/2} - 24t \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

$$= \frac{4}{t^5} + \frac{t^{1/2}}{t^5}$$

$$= 4t^{-5} + t^{-9/2}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$\begin{aligned} h'(t) &= \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1} \\ &= \frac{3}{2}t^{1/2} + 4 - \frac{15}{2}t^{3/2} - 24t \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

$$= \frac{4}{t^5} + \frac{t^{1/2}}{t^5}$$

$$= 4t^{-5} + t^{-9/2}$$

$$h'(t) = 4 \cdot (-5)t^{-5-1} - \frac{9}{2}t^{-9/2-1}$$

RULES OF DIFFERENTIATION

ADVANCED EXAMPLES

Find the derivative of the following functions:

$$f(t) = t^2 - 3t^6 + 5e^t$$

$$\begin{aligned} f'(t) &= 2t^{2-1} - 3 \cdot 6t^{6-1} + 5 \cdot e^t \\ &= 2t - 18t^5 + 5e^t \end{aligned}$$

$$h(t) = (t - 3t^2)(\sqrt{t} + 4)$$

$$= t\sqrt{t} + 4t - 3t^2\sqrt{t} - 12t^2$$

$$= t^{3/2} + 4t - 3t^{5/2} - 12t^2$$

$$\begin{aligned} h'(t) &= \frac{3}{2}t^{3/2-1} + 4 - 3 \cdot \frac{5}{2}t^{5/2-1} - 12 \cdot 2t^{2-1} \\ &= \frac{3}{2}t^{1/2} + 4 - \frac{15}{2}t^{3/2} - 24t \end{aligned}$$

$$h(t) = \frac{4 + \sqrt{t}}{t^5}$$

$$= \frac{4}{t^5} + \frac{t^{1/2}}{t^5}$$

$$= 4t^{-5} + t^{-9/2}$$

$$\begin{aligned} h'(t) &= 4 \cdot (-5)t^{-5-1} - \frac{9}{2}t^{-9/2-1} \\ &= -20t^{-6} - \frac{9}{2}t^{-11/2} \end{aligned}$$