
# Lesson 5: Separable Equations. Singular Solutions

Francisco Blanco-Silva

University of South Carolina



September 4, 2013

► The concept of a differential equation

$$F(x,y,y',\ldots,y^{(n)})=0$$

► The concept of a differential equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

► The concept of **order** of a differential equation.

► The concept of a differential equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

- ► The concept of **order** of a differential equation.
- ► The concept of a **general solution**

► The concept of a differential equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

Singular Solutions

- ► The concept of **order** of a differential equation.
- ► The concept of a **general solution**
- ► The concepts of an **initial value problem** (IVP) and **particular solution**.

► The concept of a differential equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

- ▶ The concept of **order** of a differential equation.
- ► The concept of a **general solution**
- ► The concepts of an **initial value problem** (IVP) and particular solution.
- Slope fields

► The concept of a differential equation

$$F(x, y, y', \dots, y^{(n)}) = 0$$

Singular Solutions

- ► The concept of **order** of a differential equation.
- ► The concept of a **general solution**
- ► The concepts of an **initial value problem** (IVP) and **particular solution**.
- ► Slope fields
- Approximations to solutions via Euler's Method and Improved Euler's Method

#### **DEFINITION**

The first order equation y' = H(x, y) is called separable if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

#### DEFINITION

The first order equation y' = H(x, y) is called **separable** if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

#### DEFINITION

The first order equation y' = H(x, y) is called separable if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

$$\frac{dy}{dx} = H_1(x)H_2(y)$$

#### DEFINITION

The first order equation y' = H(x, y) is called separable if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

$$\frac{dy}{dx} = H_1(x)H_2(y)$$
$$\frac{dy}{H_2(y)} = H_1(x) dx$$

#### DEFINITION

The first order equation y' = H(x, y) is called separable if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

$$\frac{dy}{dx} = H_1(x)H_2(y)$$

$$\frac{dy}{H_2(y)} = H_1(x) dx$$

$$\int \frac{dy}{H_2(y)} = \int H_1(x) dx$$

#### **DEFINITION**

The first order equation y' = H(x, y) is called separable if we can write H(x, y) as the product of a function of x and a function of y:

$$H(x,y) = H_1(x)H_2(y)$$

We may find general solutions by simple integration:

$$\frac{dy}{dx} = H_1(x)H_2(y)$$

$$\frac{dy}{H_2(y)} = H_1(x) dx$$

$$\int \frac{dy}{H_2(y)} = \int H_1(x) dx$$

Solutions found this way may be expressed explicitly (that is, solving for *y* after integration), or implicitly (without solving for *y*)

#### EXAMPLES

# Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3}$$

$$y(0) = 7$$

**EXAMPLES** 

### Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} \qquad \qquad y(0) = 7$$

We start by re-arranging the equation and taking integrals:

$$\frac{dy}{dx} = \underbrace{6x}^{H_1(x)} \underbrace{(y-1)^{2/3}}_{(y-1)^{2/3}} \qquad \int \frac{dy}{(y-1)^{2/3}} = \int 6x \, dx$$

**EXAMPLES** 

# Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

We start by re-arranging the equation and taking integrals:

$$\frac{dy}{dx} = 6x (y-1)^{2/3} \qquad \int \frac{dy}{(y-1)^{2/3}} = \int 6x \, dx$$

$$3(y-1)^{1/3} = 3x^2 + C$$

**EXAMPLES** 

# Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

We start by re-arranging the equation and taking integrals:

$$\frac{dy}{dx} = 6x (y-1)^{2/3} \qquad \int \frac{dy}{(y-1)^{2/3}} = \int 6x \, dx$$
$$3(y-1)^{1/3} = 3x^2 + C$$

To solve the IVP, we need to force y(0) = 7:

**EXAMPLES** 

# Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

We start by re-arranging the equation and taking integrals:

$$\frac{dy}{dx} = 6x (y-1)^{2/3} \qquad \int \frac{dy}{(y-1)^{2/3}} = \int 6x \, dx$$
$$3(y-1)^{1/3} = 3x^2 + C$$

To solve the IVP, we need to force y(0) = 7:

$$3(7-1)^{1/3} = 3 \cdot 0^2 + C$$
  $C = 3 \cdot 6^{1/3}$ 

**EXAMPLES** 

# Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

We start by re-arranging the equation and taking integrals:

$$\frac{dy}{dx} = 6x (y-1)^{2/3} \qquad \int \frac{dy}{(y-1)^{2/3}} = \int 6x \, dx$$
$$3(y-1)^{1/3} = 3x^2 + C$$

To solve the IVP, we need to force y(0) = 7:

$$3(7-1)^{1/3} = 3 \cdot 0^2 + C$$
  $C = 3 \cdot 6^{1/3}$   
Solution:  $(y-1)^{1/3} = x^2 + 6^{1/3}$ 

**EXAMPLES** 

### Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

Note the form of the solution:

$$(y-1)^{1/3} = x^2 + 6^{1/3}$$

This is **implicit**. If we want to provide an **explicit** solution, we need to solve for *y*:

**EXAMPLES** 

### Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

Note the form of the solution:

$$(y-1)^{1/3} = x^2 + 6^{1/3}$$

This is **implicit**. If we want to provide an **explicit** solution, we need to solve for *y*:

$$y - 1 = (x^2 + 6^{1/3})^3$$

**EXAMPLES** 

### Example

Find a particular solution of the initial value problem

$$y' = 6x(y-1)^{2/3} y(0) = 7$$

Note the form of the solution:

$$(y-1)^{1/3} = x^2 + 6^{1/3}$$

This is **implicit**. If we want to provide an **explicit** solution, we need to solve for *y*:

$$y - 1 = (x^2 + 6^{1/3})^3$$
$$y = 1 + (x^2 + 6^{1/3})^3$$

#### EXAMPLES

### Examples

$$y' = \frac{(x-1)y^5}{x^2(2y^3 - y)}$$

#### EXAMPLES

## Examples

$$y' = \frac{(x-1)y^5}{x^2(2y^3 - y)}$$

$$\frac{dy}{dx} = \frac{x-1}{x^2} \cdot \frac{y^5}{y(2y^2-1)}$$

EXAMPLES

### Examples

$$y' = \frac{(x-1)y^5}{x^2(2y^3 - y)}$$

$$\frac{dy}{dx} = \frac{x-1}{x^2} \cdot \frac{y^5}{y(2y^2 - 1)}$$
$$\frac{2y^2 - 1}{y^4} dy = (x^{-1} - x^{-2}) dx$$

#### EXAMPLES

### Examples

$$y' = \frac{(x-1)y^5}{x^2(2y^3 - y)}$$

$$\frac{dy}{dx} = \frac{x-1}{x^2} \cdot \frac{y^5}{y(2y^2 - 1)}$$
$$\frac{2y^2 - 1}{y^4} dy = (x^{-1} - x^{-2}) dx$$
$$\int (2y^{-2} - y^{-4}) dy = \int (x^{-1} - x^{-2}) dx$$

#### EXAMPLES

### Examples

$$y' = \frac{(x-1)y^5}{x^2(2y^3 - y)}$$

$$\frac{dy}{dx} = \frac{x-1}{x^2} \cdot \frac{y^5}{y(2y^2 - 1)}$$

$$\frac{2y^2 - 1}{y^4} dy = (x^{-1} - x^{-2}) dx$$

$$\int (2y^{-2} - y^{-4}) dy = \int (x^{-1} - x^{-2}) dx$$

$$-2y^{-1} + \frac{1}{3}y^{-3} = \ln|x| + x^{-1} + C$$

#### **EXAMPLES**

Sometimes, equations that don't look separable may be so, if manipulated accordingly.

# Example

$$y' = 1 + x + y + xy$$

#### **EXAMPLES**

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

# Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

#### **EXAMPLES**

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

# Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

$$y' = 1 + x + y + xy$$

#### **EXAMPLES**

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

### Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

$$y' = 1 + x + y + xy$$
  
 $y' = 1 + x + y(1 + x)$ 

#### **EXAMPLES**

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

### Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

$$y' = 1 + x + y + xy$$
  

$$y' = 1 + x + y(1 + x)$$
  

$$y' = (1 + y)(1 + x)$$

#### **EXAMPLES**

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

### Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

$$y' = 1 + x + y + xy$$
  
 $y' = 1 + x + y(1 + x)$   
 $y' = (1 + y)(1 + x)$ 

$$\frac{dy}{1 + y} = (1 + x) dx$$

#### EXAMPLES

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

### Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

#### EXAMPLES

Sometimes, equations that *don't look separable* may be so, if manipulated accordingly.

# Example

Find a general solution (implicit) of the equation

$$y' = 1 + x + y + xy$$

$$y' = 1 + x + y + xy$$
  
 $y' = 1 + x + y(1 + x)$   
 $y' = (1 + y)(1 + x)$ 

$$\frac{dy}{1+y} = (1+x) dx$$

$$\int \frac{dy}{1+y} = \int (1+x) dx$$

$$\ln|1+y| = x + \frac{1}{2}x^2 + C$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \cot x dx$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \cot x dx$$

$$\int \frac{dy}{y} = \int \frac{\cos x}{\sin x} dx$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \cot x dx$$

$$\int \frac{dy}{y} = \int \frac{\cos x}{\sin x} dx$$

$$\ln|y| = \ln|\sin x| + C$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \cot x dx$$

$$\int \frac{dy}{y} = \int \frac{\cos x}{\sin x} dx$$

$$\ln|y| = \ln|\sin x| + C$$

$$e^{\ln|y|} = e^{\ln|\sin x|} \underbrace{e^{C}}_{A}$$

**EXAMPLES** 

# Example

$$\tan x \, \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \cot x dx$$

$$\int \frac{dy}{y} = \int \frac{\cos x}{\sin x} dx$$

$$\ln|y| = \ln|\sin x| + C$$

$$e^{\ln|y|} = e^{\ln|\sin x|} \underbrace{e^{C}}_{A}$$

$$|y| = A|\sin x|$$

# SINGULAR SOLUTIONS

#### MOTIVATION AND DEFINITION

Let us examine the previous equations again, with their found general solutions:

$$y' = 6x(y-1)^{2/3} (y-1)^{1/3} = x^2 + C$$

$$y' = \frac{x-1}{x^2} \cdot \frac{y^5}{2y^3 - y} \frac{1}{3}y^{-3} - 2y^{-1} = \ln|x| + \frac{1}{x} + C$$

$$y' = (1+x)(1+y) \ln|1+y| = x + \frac{1}{2}x^2 + C$$

$$\tan x \frac{dy}{dx} = y |y| = A|\sin x|$$

There are solutions to these equations that cannot be found by integration. They can only be found by inspection of the equations. The slope fields of the equations also reveal these so called singular solutions. Can you find them?

# SINGULAR SOLUTIONS

#### MOTIVATION AND DEFINITION

Let us examine the previous equations again, with their found general solutions:

$$y' = 6x(y-1)^{2/3} (y-1)^{1/3} = x^2 + C y = 1$$

$$y' = \frac{x-1}{x^2} \cdot \frac{y^5}{2y^3 - y} \frac{1}{3}y^{-3} - 2y^{-1} = \ln|x| + \frac{1}{x} + C y = 0$$

$$y' = (1+x)(1+y) \ln|1+y| = x + \frac{1}{2}x^2 + C y = -1$$

$$\tan x \frac{dy}{dx} = y |y| = A|\sin x| y = 0$$

There are solutions to these equations that cannot be found by integration. They can only be found by *inspection* of the equations. The slope fields of the equations also reveal these so called singular solutions. Can you find them?