

## MATH 141 HW 16 (6589721)

Due: Thu Nov 20 2014 11:59 PM EST

Question

[1](#)
[2](#)
[3](#)
[4](#)
[5](#)
[6](#)
[7](#)
[8](#)
[9](#)
[10](#)
[11](#)
[12](#)
[13](#)
[14](#)
[15](#)
[16](#)
[17](#)
[18](#)
[19](#)
[20](#)
[21](#)
[22](#)
[23](#)
[24](#)
[25](#)
[26](#)
[27](#)
[28](#)
[29](#)
[30](#)
[31](#)
[32](#)
[33](#)
[34](#)
[35](#)

## 1. Question Details

SCalcET6 5.2.033. [698729]


The graph of  $f$  is shown. Evaluate each integral by interpreting it in terms of areas.

$$a = 16$$

$$b = 32$$

$$c = 48$$

$$d = 64$$



$$(a) \int_0^{16} f(x) dx = \boxed{\phantom{000}} \quad \text{🔑 256}$$

$$(b) \int_0^{40} f(x) dx = \boxed{\phantom{000}} \quad \text{🔑 640}$$


$$(c) \int_{40}^{56} f(x) dx = \boxed{\phantom{000}} \quad \text{🔑 -192}$$

$$(d) \int_0^{72} f(x) dx = \boxed{\phantom{000}} \quad \text{🔑 128}$$

## 2. Question Details

SCalcET6 5.2.034.MI. [1816769]

The graph of  $g$  consists of two straight lines and a semicircle. Use it to evaluate each integral.



(a)  $\int_0^6 g(x) \, dx$

$\boxed{36}$

(b)  $\int_6^{18} g(x) \, dx$

$\boxed{-18\pi}$

(c)  $\int_0^{21} g(x) \, dx$

$\boxed{40.5 - 18\pi}$

## 3. Question Details

SCalcET6 5.2.035.MI. [1386386]

Evaluate the integral by interpreting it in terms of areas.

$$\int_0^3 \left( \frac{7}{2}x - 7 \right) dx$$

$\boxed{-\frac{21}{4}}$

## 4. Question Details

SCalcET6 5.2.038.MI. [1816965]

Evaluate the integral by interpreting it in terms of areas.

$$\int_{-3}^3 (4 - 2x) \, dx$$

$\boxed{24}$

## 5. Question Details

SCalcET6 5.3.047. [1817356]

Use a graph to give a rough estimate of the area of the region that lies beneath the given curve. Then find the exact area.

$$y = \sqrt[5]{x}, \quad 0 \leq x \leq 32$$

  160/3

## 6. Question Details

SCalcET6 5.3.048. [1817214]

Use a graph to give a rough estimate of the area of the region that lies beneath the given curve. Then find the exact area.

$$y = x^{-3}, \quad 2 \leq x \leq 5$$

  21/200

## 7. Question Details

SCalcET6 5.3.049. [1290674]

Find the area of the region that lies beneath the given curve.

$$y = \sin(x), \quad 0 \leq x \leq \pi$$

 2

## 8. Question Details

SCalcET6 5.3.053. [1291220]

Find the derivative of the function.

$$g(x) = \int_{2x}^{7x} \frac{u^2 - 3}{u^2 + 3} du$$

  $-2 \cdot \frac{4x^2 - 3}{4x^2 + 3} + 7 \cdot \frac{49x^2 - 3}{49x^2 + 3}$ 

## 9. Question Details

SCalcET6 5.3.054. [1290381]

Find the derivative of the function.


$$g(x) = \int_{\tan(x)}^{4x^2} \frac{1}{\sqrt{5+t^3}} dt$$

  $-\frac{(\sec(x))^2}{\sqrt{5+(\tan(x))^3}} + \frac{8x}{\sqrt{5+64x^6}}$

## 10. Question Details

SCalcET6 5.3.003. [872785]

Let  $g(x) = \int_0^x f(t)dt$ , where  $f$  is the function whose graph is shown.



(a) Evaluate  $g(0)$ ,  $g(5)$ ,  $g(10)$ ,  $g(15)$  and  $g(30)$ .

$$g(0) = \boxed{\phantom{000}} \quad \boxed{0}$$

$$g(5) = \boxed{\phantom{000}} \quad \boxed{50}$$

$$g(10) = \boxed{\phantom{000}} \quad \boxed{125}$$

$$g(15) = \boxed{\phantom{000}} \quad \boxed{175}$$

$$g(30) = \boxed{\phantom{000}} \quad \boxed{75}$$

(b) On what interval is  $g$  increasing?

$$(\boxed{\phantom{000}} \quad \boxed{0}, \boxed{\phantom{000}} \quad \boxed{15})$$

(c) Where does  $g$  have a maximum value?

$$x = \boxed{\phantom{000}} \quad \boxed{15}$$

(d) Sketch a rough graph of  $g$ . (Do this on paper. Your instructor may ask you to turn in this work.)

## 11. Question Details

SCalcET6 5.3.007.MI. [1387698]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$g(x) = \int_1^x \frac{3}{t^3 + 1} dt$$

$$g'(x) = \boxed{\phantom{000}} \quad \boxed{\frac{3}{x^3 + 1}}$$

## 12. Question Details

SCalcET6 5.3.008. [1816419]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$g(x) = \int_2^x e^{4t^2 - 5t} dt$$

$$g'(x) = \boxed{\phantom{000}} \quad \boxed{e^{4x^2 - 5x}}$$

## 13. Question Details

SCalcET6 5.3.009. [1817010]

Use Part 1 of the [Fundamental Theorem of Calculus](#) to find the derivative of the function.

$$g(y) = \int_3^y t^2 \sin 2t \, dt$$

$$g'(y) = \boxed{y^2 \sin(2y)}$$

## 14. Question Details

SCalcET6 5.3.010. [1817149]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$g(r) = \int_2^r \sqrt{x^2 + 2} \, dx$$

$$g'(r) = \boxed{\sqrt{r^2 + 2}}$$

## 15. Question Details

SCalcET6 5.3.012.MI. [1386615]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$G(x) = \int_x^2 \cos \sqrt{t} \, dt$$

$$G'(x) = \boxed{-\cos(\sqrt{x})}$$

## 16. Question Details

SCalcET6 5.3.013.MI. [1387359]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$h(x) = \int_2^{1/x} \arctan 5t \, dt$$

$$h'(x) = \boxed{-\frac{\arctan\left(\frac{5}{x}\right)}{x^2}}$$

## 17. Question Details

SCalcET6 5.3.014. [1290601]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$h(x) = \int_2^{x^2} \sqrt{6 + t^3} \, dt$$

$$h'(x) = \boxed{2x\sqrt{6 + x^6}}$$

## 18. Question Details

SCalcET6 5.3.015. [1816284]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$y = \int_5^{\tan x} \sqrt{6t + \sqrt{t}} \, dt$$

$$y' = \boxed{\sqrt{6 \tan(x) + \sqrt{\tan(x)}} \sec^2(x)}$$

## 19. Question Details

SCalcET6 5.3.016. [1291391]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$y = \int_1^{\cos(x)} (5 + v^5)^6 \, dv$$

$$y' = \boxed{-(5 + (\cos(x))^5)^6 \sin(x)}$$

## 20. Question Details

SCalcET6 5.3.017. [1816635]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$y = \int_{2-3x}^3 \frac{u^3}{1+u^2} \, du$$

$$y' = \boxed{\frac{3(2-3x)^3}{1+(2-3x)^2}}$$

## 21. Question Details

SCalcET6 5.3.018. [1817418]

Use Part 1 of the Fundamental Theorem of Calculus to find the derivative of the function.

$$y = \int_{e^x}^0 2 \sin^2 t \, dt$$

$$y' = \boxed{-2e^x \sin^2(e^x)}$$

## 22. Question Details

SCalcET6 5.3.019. [1817104]

Evaluate the integral.

$$\int_{-2}^3 (x^3 - 6x) \, dx$$


 5/4

## 23. Question Details

SCalcET6 5.3.020. [698756]

Evaluate the integral.

$$\int_{-3}^6 7 dx$$

  63

## 24. Question Details

SCalcET6 5.3.021. [1816170]

Evaluate the integral.

$$\int_2^6 (3 - 2t + 3t^2) dt$$

  188

## 25. Question Details

SCalcET6 5.3.022.MI. [1816672]

Evaluate the integral.

$$\int_0^1 \left( 3 + \frac{3}{4}u^4 - \frac{4}{5}u^9 \right) du$$

  307/100

## 26. Question Details

SCalcET6 5.3.023.MI. [1387391]

Evaluate the integral.

$$\int_0^1 x^{2/5} dx$$

  5/7

## 27. Question Details

SCalcET6 5.3.024. [656986]

Evaluate the integral.

$$\int_1^{32} \sqrt[5]{x} dx$$

  105/2

## 28. Question Details

SCalcET6 5.3.025. [698640]

Evaluate the integral.

$$\int_1^4 \frac{5}{t^4} dt$$

 105/64

## 29. Question Details

SCalcET6 5.3.026. [1816879]

Evaluate the integral.

$$\int_{5\pi}^{6\pi} \cos \theta \, d\theta$$

 0

## 30. Question Details

SCalcET6 5.3.027. [872789]

Evaluate the integral.

$$\int_0^1 x(1+x^5) \, dx$$

 9/14

## 31. Question Details

SCalcET6 5.3.028. [698708]

Evaluate the integral.

$$\int_0^1 (4 + 6x\sqrt{x}) \, dx$$

 32/5

## 32. Question Details

SCalcET6 5.3.029.MI. [1816347]

Evaluate the integral.

$$\int_1^4 \frac{x-5}{\sqrt{x}} \, dx$$

 -16/3



33. Question Details

SCalcET6 5.3.030. [1816731]

Evaluate the integral.

$$\int_0^4 (y - 3)(2y + 1) dy$$

  -28/3

34. Question Details

SCalcET6 5.3.033.MI. [1816434]

Evaluate the integral.

$$\int_2^3 (2 + 2y)^2 dy$$


  148/3

35. Question Details

SCalcET6 5.3.035. [1816281]

Evaluate the integral.

$$\int_1^9 \frac{1}{2x} dx$$


   $\frac{\ln(9)}{2}$ 

36. Question Details

SCalcET6 5.3.036. [1816133]

Evaluate the integral.

$$\int_0^3 3^x dx$$


   $\frac{26}{\ln(3)}$ 

37. Question Details

SCalcET6 5.3.040. [2792791]

Evaluate the integral.

$$\int_3^4 \frac{2 + u^2}{u^3} du$$

   $\frac{7}{144} + \ln\left(\frac{4}{3}\right)$

## 38. Question Details

SCalcET6 5.3.055. [1291181]

Find the derivative of the function.

$$g(x) = \int_{\sqrt{x}}^{x^2} \sqrt{t} \sin(t) dt$$

$$\boxed{-\frac{\sin(\sqrt{x})}{2\sqrt{x}} + 2x^2 \sin(x^2)}$$

## 39. Question Details

SCalcET6 5.3.056. [1290003]

Find the derivative of the function.

$$y = \int_{4 \cos(x)}^{2x} \cos(u^4) du$$

$$\boxed{4 \sin(x) \cos(256 (\cos(x))^4) + 2 \cos(16x^4)}$$

## Assignment Details

Name (AID): **MATH 141 HW 16 (6589721)**Submissions Allowed: **100**Category: **Homework**

Code:

Locked: **Yes**Author: **Blanco-Silva, Francisco** ( [blanco@math.sc.edu](mailto:blanco@math.sc.edu) )Last Saved: **Nov 19, 2014 08:23 AM EST**Permission: **Protected**Randomization: **Person**Which graded: **Last**

## Feedback Settings

Before due date

Question Score

Assignment Score

Publish Essay Scores

Question Part Score

Mark

Response

Save Work

After due date

Question Score

Assignment Score

Publish Essay Scores

Key

Question Part Score

Solution

Mark

Add Practice Button

Help/Hints

Response